

Dynamics of D-brane Black Holes

Steven S. Gubser¹

Abstract

We explore the interplay between black holes in supergravity and quantum field theories on the world-volumes of D-branes. Each sheds light on the other in various ways. The world-volume description provides a statistical mechanical picture of black hole entropy and a manifestly unitary account of Hawking radiation. The black hole description elucidates the nature of the couplings between the world-volume theory and supergravity in the bulk. Finally, the computation of Green's functions in the world-volume theory can be mapped in a large N limit to a classical minimization problem in supergravity.

A brief summary of black hole entropy calculations for D-brane black holes is followed by a detailed study of particle absorption by black holes whose string theory description involves D-branes intersecting along a string. A two-dimensional conformal field theory with large central charge describes the low-energy excitations of this string. In the string theory description, Hawking radiation is simply the time-reversal of particle absorption. The greybody factors (deviations in the power spectrum from Planck's law) are characteristic of conformal field theory at finite temperature.

Particle absorption by extremal three-branes is examined next, with particular attention to the implications for supersymmetric gauge theory in four dimensions. A fascinating duality between supergravity and gauge theory emerges from the study of these processes. Fields of supergravity are dual to local operators in the gauge theory. A non-renormalization theorem of $\mathcal{N} = 4$ gauge theory helps explain why certain aspects of the duality can be explored perturbatively. Anomalous dimensions of a large class of local operators in the gauge theory are shown to become large at strong 't Hooft coupling, signaling a possible simplification of $\mathcal{N} = 4$ gauge theory in this limit.

A dissertation presented to the faculty of Princeton University
in candidacy for the degree of Doctor of Philosophy

Recommended for acceptance by the Department of Physics

June 1998

©Copyright by Steven Scott Gubser, 1998. All rights reserved.

¹ssgubser@born.harvard.edu

Acknowledgements

The research summarized in this thesis was supported in part by DOE Grant No. DE-FG02-91ER40671, by the NSF Presidential Young Investigator Award PHY-9157482, by the James S. McDonnell Foundation under Grant No. 91-48, and by the Hertz Foundation.

I would like to thank my collaborators: Curt Callan, Aki Hashimoto, Michael Krasnitz, Juan Maldacena, Amanda Peet, Alexander Polyakov, Arkady Tseytlin, and particularly my advisor, Igor Klebanov. Without them not a tenth of the research in this dissertation would have been realized.

Thanks to all my teachers, particularly E. Lieb, P. Meyers, H. Osborn, B. van Fraassen, A. Eisenkraft, P. Ryan, P. Sarnak, S. Sondhi, and D. Gross.

I am grateful to Laurel Lerner for her encouragement, both on matters academic and pianistic.

Finally, thanks to my brother Charles, for leading the way, to my grandparents, for many years of encouragement, and to my parents.

Contents

1	Introduction	4
2	Entropy considerations	11
2.1	Thermodynamics of coincident branes	11
2.2	Embedding of three-branes and the effective string	13
3	D-brane approach to fixed scalars	16
3.1	Introduction	16
3.2	Field theory effective action considerations	17
3.2.1	$D = 5$ case	17
3.2.2	$D = 4$ case	22
3.3	Effective string couplings	24
3.4	Semi-classical description of absorption	27
3.5	D-brane absorption cross-sections and emission rates	30
3.6	Conclusions	34
4	Partial wave absorption by effective strings	36
4.1	Introduction	36
4.2	The semi-classical computation	37
4.3	The effective string analysis	39
4.4	Limitations on partial wave absorption	46
4.5	Conclusion	47
5	Photons and fermions falling into four-dimensional black holes	49
5.1	Introduction	49
5.2	Minimally coupled photons in $\mathcal{N} = 4$ supergravity	50
5.3	Separable equations for gauge fields	52
5.4	Semi-classical absorption probabilities	54
5.4.1	Probabilities from energy fluxes	55
5.4.2	Matching solutions	56
5.5	The axion and minimally coupled fermions	58
5.5.1	The axion	58
5.5.2	Minimally coupled fermions	59
5.6	The effective string model	61

5.7	Conclusion	66
6	Absorption by non-dilatonic branes	71
6.1	Introduction	71
6.2	Spacetime effective actions and perturbations around non-dilatonic p -brane solutions .	73
6.2.1	Perturbations of three-branes	73
6.2.2	Perturbations of two-branes and five-branes	75
6.3	The self-dual three-brane	75
6.3.1	Classical absorption	75
6.3.2	Universality of the absorption cross-section for minimally coupled scalars . . .	76
6.3.3	Higher partial waves	79
6.4	Absorption cross-sections for M-branes	81
6.5	Conclusions	84
7	Cross-sections and Schwinger terms in the world-volume	86
7.1	Introduction	86
7.2	D-brane approach to absorption	88
7.3	Extension to other branes	91
8	Green's functions from supergravity	94
8.1	Introduction	94
8.2	Green's functions from the supergravity action	96
8.2.1	Preliminary: symmetries and boundary conditions	97
8.2.2	The central charge from the stress tensor two-point function	98
8.3	Massive string states and anomalous dimensions	101
8.4	Conclusions	103
9	Concluding Remarks	104
	Bibliography	106

Chapter 1

Introduction

The usual conception of how string theory (see for example [1]) makes contact with the physical world is that string theory is compactified from ten to four dimensions on a manifold of size on the order 10^{-32} cm—roughly as small compared to a proton as a proton is to Mount Everest. All the “fundamental” particles of the Standard Model of particle physics are supposed to be realized as different excitation modes of strings. The most powerful of particle accelerators probe distances only about a sixth of the way (on a logarithmic scale) between the proton’s size and the string scale, so it is difficult to imagine how the “stringy” character of elementary particles could ever be directly observed. There is however an extensive literature investigating the phenomenological implications of string theory and how contact may yet be made with experiment (for a recent review see [2]).

String theory is also a quantum theory of gravity. In fact, it is the only consistent quantum theory to date which contains gravity in any dynamical sense. (In two and three dimensions, theories of quantum gravity can be formulated, but they are non-dynamical in the sense that gravitons do not propagate). Another avenue, then, for connecting string theory with the physical world is via quantum gravitational effects. As an example one might consider Hawking radiation from black holes. This is an essentially quantum mechanical effect: whereas black holes classically cannot radiate (due to the causal structure of their geometry), quantum mechanically they radiate as blackbodies. For many years a fundamental, microscopic description of this effect was almost as much a mystery in string theory as in General Relativity. However, recent advances in the description of string-theoretic solitons have enabled string theorists to give an account of Hawking radiation for a certain limited class of black holes. It is worth emphasizing that we are still very far from experiment, particularly since the black holes amenable to treatment in string theory in general carry a large electric charge. Astrophysically, such a charge would quickly be neutralized by infalling matter.

The recent developments in the understanding of dualities in string theory (for a review see [3]) are very much at the heart of progress in the string-theoretic description of black holes. A particularly prominent role has been played by the so-called D-branes [4], which are objects derived via a detailed understanding of T-duality in open and closed string theories. Closed string theories require that strings cannot have ends, whereas open strings by definition do have ends. However, the full consistency of closed string theories under duality symmetries requires the existence of D-branes, which can be thought of as special surfaces on which strings are allowed to end. But they are more than this mathematical characterization suggests: they are dynamical objects in themselves, with a definite tension which at weak coupling is large compared to the tension of fundamental strings. The large tension of D-branes suggests that they may be suitable objects from which to build black holes. The development of this idea and its implications for the description of dynamical effects like Hawking radiation are two of the main themes of this thesis.

The link between extremal or near-extremal black holes and the world-volume theories describing

excitations of branes has been explored in a now-voluminous literature. Various aspects of the general program of understanding this link have been initiated in [5, 6] (entropy of effective string models), [7, 8] (Hawking radiation processes), and [9, 10] (absorption into extremal three-branes). A tangentially related subject, the probing of sub-string-scale geometries using D-branes, grew from the initial work of [11] into the vast subject of matrix theory [12].

The excitations of D-branes at low energies are described by supersymmetric gauge theories [13]. This observation opens up yet another avenue for connecting string theory with mainstream physics: if on one hand D-branes describe black holes, and on the other hand are characterized at low energies by gauge theory, then perhaps some information about gauge theory is encoded in the supergravity solutions of the corresponding black holes. The realization that there are concrete ways of developing this correspondence is at present one of the main sources of excitement in the string theory community. There are even hints that supersymmetry may no longer be an essential ingredient in the gravity–gauge theory correspondence. Earlier work on extracting gauge theory physics from intersecting brane configurations [14] relies very little on supergravity, but has yielded some remarkable insights into the geometric character of supersymmetric gauge theory (for a review see [15]).

In the past few months, with the appearance of the papers [16, 17, 18, 19, 20], our entire understanding of the link through string theory and D-branes between supergravity and gauge theory has evolved rapidly. The essential ingredients in the new approach are the following two insights: 1) information about the infrared behavior of gauge theories is encoded in the near-horizon geometry of black branes in supergravity; and 2) perturbations to the gauge theory can be implemented by constraints on the supergravity fields on a hypersurface which surrounds the brane at a distance close to where the near-horizon geometry merges with asymptotically flat space. The second insight, achieved in various forms in [17, 18, 19, 20], has been termed “holography” by Witten. Although this term (originally proposed by ’t Hooft and Susskind as applicable to black holes) has been used in a somewhat murky fashion in the string theory literature, its current application is appropriate in the following sense. The dynamics of the gauge theory is mirrored in the supergravity theory subjected to boundary conditions on the hypersurface, which is of one lower dimension than the ambient space (but of the same dimension as the world-volume of the gauge theory). Thus one views the gauge theory as a hologram of the supergravity.

All but the last two chapters of this thesis are devoted to the pre-holography era of black holes in string theory. Chapter 2 presents a brief investigation of the entropy of black holes built out of D-branes. The focus is on the two examples which have been most intensively studied in the literature: D3-branes in ten-dimensional spacetime, and three-charge black holes with regular horizon in five-dimensional spacetime. The remarkable insight that the D-brane approach yields is that the entropy of certain black holes can be explained in terms of the statistical mechanics of the world-volume theory of D-branes.

The remainder of the thesis is split into two parts. The first part, chapters 3-5, is based on the papers [21, 22, 23]. The focus is on the string theory description of Hawking radiation and particle absorption by near-extremal black holes in four and five dimensions. The string theory model for these black holes is an “effective string” whose excitations are described by a conformal field theory (CFT). The influential paper [5] includes a computation of the central charge of this conformal field theory via a moduli space argument. In chapter 2 we develop a more concise but limited version of this argument using index theorems. Here the effective string is realized as a D3-brane with two of its spatial dimensions wrapped around a four-dimensional manifold. A more familiar description, developed initially in [6], is to view the effective string as some number of D1-branes embedded in D5-branes (whose four extra dimensions are again wrapped around a four-dimensional manifold). A black hole in five dimensions can be obtained by wrapping the effective string around a circle. Black holes in four dimensions have also been treated via effective strings which can be constructed within string theory [24, 25], but which arise most naturally intersecting five-branes in M-theory [26, 27].

In the extensive recent literature comparing black hole entropy with microscopic state counts,

black holes which admit an effective string description have always proven the most reliably tractable. Agreements between the Bekenstein-Hawking entropy and the statistical mechanical entropy are no longer considered miraculous in the case of supersymmetric black holes, because the two calculations are different ways of counting states that preserve supersymmetry. These so-called BPS states are generally protected against mass renormalizations by their short multiplet structure under the supersymmetry algebra. More surprising are successful comparisons of near-extremal black hole entropy with statistical mechanical analyses, because in such cases there is no supersymmetry to prevent microstates from changing their masses as one proceeds from the region of weak coupling where the statistical mechanical analysis is performed to the region of strong coupling where supergravity is applicable. In chapter 2 we will see that such agreements are indeed rather the exception than the rule. However, comparisons of the entropy of near-extremal effective string black holes have succeeded in certain limits [6]. As it turns out, the central charge alone determines the effective string model's entropy in the relevant limits, usually termed the "dilute gas regime."¹ Thus the status of the central charge as a non-renormalized quantity (namely a measure of the density of BPS states) basically guarantees the success of the entropy comparisons in the dilute gas regime.

The original realization [5] of the effective string CFT was as a supersymmetric non-linear sigma model whose target space was a symmetric power of a four dimensional manifold. It is straightforward to extract the central charge and proceed with entropy calculations, but it has proven more difficult to determine what are the primary fields and how they couple to the fields of supergravity. Precisely these questions can be explored via calculations of absorption and Hawking emission by black holes.

In the semi-classical supergravity description of black holes, Hawking radiation and particle absorption are radically different phenomena: absorption is a classical effect, while Hawking radiation is strictly quantum mechanical. To be precise, Hawking's formula (see for example [28]) relating the emission rate $d\Gamma$ of massless particles into an element of wave-number phase space d^3k to the absorption cross-section σ is

$$d\Gamma = \hbar c \frac{\sigma}{e^{E/k_B T} - 1} \frac{d^3k}{(2\pi)^3} \quad (1.1)$$

where the temperature is given in terms of the surface gravity κ at the horizon:

$$T = \frac{\hbar}{c} \frac{\kappa}{2\pi k_B} . \quad (1.2)$$

Planck's constant is involved in two essential ways, and both the Hawking rate and the Hawking temperature vanish as $\hbar \rightarrow 0$. From here on we will adopt units in which $\hbar = c = k_B = 1$.

In string theory the relation between absorption and Hawking radiation is simply time-reversal. Absorption into an extremal black hole is mediated by S -matrix elements which can be calculated from a knowledge of couplings between supergravity and the effective string. Hawking radiation is impossible for an extremal black hole, but the effective string description extends easily to the near-extremal case, and then the same S -matrix element that describes particle absorption also mediates Hawking radiation. Time-reversal symmetry is broken only by the kinematics of the thermal field theory on the effective string. Absorption and emission processes are related by detailed balance, so that an effective string black hole can be in equilibrium with a thermal sea of particles in the ambient space (it should be remarked at this point that unlike Schwarzschild black holes, near-extremal black holes generically have positive specific heat, so thermal equilibrium is possible). Indeed, in the statistical mechanical string theory picture, (1.1) is nothing more than the statement of detailed balance. A concrete illustration of this point will be developed in chapter 3 in the context of fixed scalars.

Supergravities in dimensions less than ten with extended supersymmetry emerge from the low-energy limit of string theory, and in general have a large number of scalar fields [29]. For the purpose

¹This is only true once the proper identification of charges and angular momenta has been made in the effective string description.




Figure 1.1: a) Particle absorption by the effective string; b) The string disk diagram for absorption.

of studying effective string black holes, we will primarily be interested in maximally supersymmetric theories in four or five dimensions, which admit extremal black holes with regular horizons ([30] and references therein). Some of the scalar fields can be regarded as moduli of the compactified geometry that takes the theory down from ten dimensions to four or five. In general, small fluctuations of these scalars obey the curved space wave equation (the massless Klein-Gordon equation), and are termed “minimal scalars.” Their absorption into black holes can be described in the effective string language as a massless closed string state hitting a set of intersecting D-branes and turning into a pair of open strings that run in opposite directions along the $1 + 1$ -dimensional intersection manifold, as depicted in figure 1.1a). In principle these absorption amplitudes can be computed via a string disk diagram, figure 1.1b), with one insertion in the bulk (the closed string vertex) and two on the boundary (the open string vertices). Calculations along these lines have been explored [31], but difficulties arise due to the complicated nature of the D-brane bound state which constitutes the effective string. It proves more insightful to propose as a low-energy effective action the Dirac-Born-Infeld (DBI) action [32]. Because this action includes among its fields the induced metric on the effective string, the dilaton, and other fields of supergravity (in a manner recently systematized in [33, 34, 35, 36] for certain simpler brane configurations than the effective string), there is an implicit prescription for calculating the coupling between supergravity fields and effective string excitations.

In view of the extensive literature on minimal scalars, our primary focus will be on another class, the so-called “fixed scalars” ([21] and references therein) whose absorption cross-section at low energies (and consequently Hawking emission) is suppressed by powers of the energy relative to that of the minimal scalars. In the effective string description, fixed scalars couple to no less than four excitations on the effective string. In figure 1.1b), one would have four rather than two boundary insertions. In the DBI effective action, the coupling turns out to be quartic rather than quadratic in the elementary fields. Morally speaking, one can view the suppression of the absorption cross-section as due to the amount of phase space available at low energies. From the point of view of supergravity, the suppression is due to a potential barrier, which might also be thought of as a variable mass term in the Klein-Gordon equation.

The standard WKB method for barrier penetration does not apply to the calculation of the low-energy cross-section because the wavelength is longer than the width of the barrier. Instead, when the linearized equations of motion are not exactly soluble, one can use approximation techniques first developed in [37]. A substantial part of chapters 3-6 is devoted to the application of these techniques to various extremal and near-extremal black holes. In particular, chapter 4 is devoted to an analysis of partial waves of a minimal scalar falling into near-extremal black holes in five dimensions, while

chapter 6 focuses on partial waves of minimal scalars falling into D3-branes in ten dimensions. Aside from a derivation of a version of the Optical Theorem in arbitrary dimensions, the semi-classical calculations in these chapters are fairly nondescript. The real interest lies in the relevance to a microscopic string theory description.

The description of partial wave absorption in the effective string model is somewhat problematic because of our limited understanding of the underlying conformal field theory. It is clear [38] that angular momentum is carried by the fermionic excitations of the effective string. The Pauli Exclusion principle would then seem to imply that only a finite number of partial waves are capable of coupling to the effective string. The analysis of chapter 4 suggests that this number is large and possibly related to the principle of cosmic censorship.

In chapter 5, an attempt is made to move a few steps closer to astrophysical reality by considering particles with spin (photons or chiral fermions) falling into a black hole in four dimensions. The effective string description is again found to be in remarkable agreement with the functional form of the cross-sections. Supersymmetry still plays an essential role in the discussion, since the black hole under consideration is realized in pure $\mathcal{N} = 4$ supergravity. This is not a theory which descends in any simple way from string theory, so the effective string model lacks a microscopic justification (in contrast to the case of the D1-D5 bound state in five dimensions or the four-charge black hole in $\mathcal{N} = 8$ four-dimensional supergravity). However, taking it as a useful effective description in terms of conformal field theory we explore the properties of conformal Green's functions at finite temperature. The powerful constraints of conformal invariance essentially dictate the energy dependence of the cross-section, also termed the greybody factor.

The second part of the thesis, chapters 6-8, is based on the papers [10, 39, 18], and is concerned primarily with the relation via string theory of ten-dimensional type IIB supergravity with four-dimensional $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) theory. The supergravity theory is the low-energy limit of type IIB string theory, which is one of the two maximally supersymmetric string theories in ten dimensions. One of the charged branes in this theory is the D3-brane, which (following [4]) one views as a $3 + 1$ -dimensional hypersurface (usually flat) in $9 + 1$ -dimensional spacetime, on which strings are allowed to end. One of the early discoveries [13] was that the low-energy effective theory describing excitations of N coincident D3-branes is $\mathcal{N} = 4$ SYM theory with gauge group $U(N)$. It has long been hoped that this theory, on account of its supersymmetry, conformal symmetry, and finiteness, would provide one of the simplest examples of an interacting quantum field theory in $3 + 1$ dimensions. However, the theory has proven notoriously difficult to solve in any sense similar to the exactly solvable quantum field theories in $1 + 1$ dimensions—for example rational conformal field theories [40, 41, 42].² For this reason, any handle one can get on $\mathcal{N} = 4$ SYM via D3-branes is of tremendous interest, independent of black hole physics.

In fact, the same supergravity methods used to explore low-energy absorption processes in the case of four- and five-dimensional black holes apply equally to the case of the extremal D3-brane. In chapter 6 these methods are compared with tree-level calculations in the world-volume theory. It is shown that the energy dependence of absorption cross-sections is predicted reliably by the gauge theory;³ however it is only for low partial waves (the s -wave or the p -wave) for the infalling particle that we have been able to obtain the properly normalized cross-section through tree-level calculations. In a sense even this level of agreement between gauge theory and supergravity is surprising, because the gauge theory calculations are valid only when the 't Hooft coupling $g_{YM}\sqrt{N}$ is small, whereas the supergravity calculations are reliable only when the spacetime curvatures are small. These turn out to be exactly opposite limits. In chapter 7 we advance an explanation for why the tree-level gauge theory computation for the s -wave was in fact reliable. The absorption cross-sections are related to

²See however [43] for recent progress on extracting correlation functions of $\mathcal{N} = 4$ SYM using superspace techniques.

³In light of recent developments [18, 19] relating Kaluza-Klein masses in the near-horizon geometry to the conformal dimensions of gauge theory operators, this agreement illustrates how the dimensions of operators coupling to arbitrary partial waves can be computed successfully at weak coupling.




Figure 1.2: The throat-brane correspondence.

the cuts in two-point functions in the gauge theory: to be precise,

$$\lim_{\kappa \rightarrow 0} \frac{i\omega}{\kappa^2} \sigma = \text{Disc } \Pi_2^{\text{SYM}}(p^2) \Big|_{p^\mu = (\omega, 0, 0, 0)} . \quad (1.3)$$

In (1.3), κ is the gravitational constant ($2\kappa^2 = 16\pi G_N$), Π_2^{SYM} is a Green's function in the gauge theory, and $p^\mu = (\omega, 0, 0, 0)$ is imposed to reflect the four-dimensional components (parallel to the D3-brane) of the ten-dimensional momentum p^M of the infalling particle. The “decoupling” limit $\kappa \rightarrow 0$ is taken in order to extract from the cross-section the part which is relevant to the renormalizable gauge theory. This corresponds to the leading dependence of σ on the energy ω . In principle, the full cross-section contains information about the full (non-renormalizable) theory describing fluctuations of the D3-branes; this is thought to be a non-abelian extension of the Dirac-Born-Infeld (DBI) action [44]. While the dynamics of the DBI theory may be reflected in a very complicated two-point function Π_2^{DBI} , certain two-point functions in the renormalizable gauge theory are quite simple and well-understood on account of its conformal symmetry. The example considered in chapter 7 is the two-point function of the stress-energy tensor, which pertains to the absorption of gravitons polarized parallel to the brane. This two-point function is in fact related to a conformal anomaly, and as such can be computed reliably at one loop. The corresponding statement regarding the graviton cross-section is that it may be computed reliably at tree level: all radiative corrections cancel.

In hindsight the formula (1.3) should have immediately suggested the far-reaching extension of the gauge theory–supergravity correspondence which we formulate in chapter 8. As a preliminary to the statement of this extension, it is perhaps useful to review a heuristic picture of the throat-brane correspondence developed in conversations with Prof. C. Callan. String theory offers two different descriptions of three-branes, each valid in a different regime of couplings. The first is the low-energy supergravity solution, pictured schematically in figure 1.2a). The near-horizon region (pictured as a long throat in the figure) has the geometry of anti-de Sitter space, AdS_5 , times a sphere S^5 of constant radius. It is joined in the “mouth region” (the vicinity of the dotted line) to the asymptotically flat region. The second description, figure 1.2b) is simply a cluster of N coincident D3-branes in flat space. Because these two pictures describe the same physical system, the response to perturbations (i.e. infalling particles from asymptotically flat infinity) should be the same. Experience with the linearized equations of motion in supergravity teaches us that perturbations which are the same far from the brane are also the same to leading order in small energies when propagated in the curved geometry down to the dotted line as when propagated to the location of the D-branes in flat space. Now imagine solving supergravity in the throat subject to boundary conditions imposed on the dotted line. In view of the physical equivalence of a) and b), that solution should encode the same physics as the D-brane world-volume gauge theory, when subjected to the same external perturbations.

The claim can be made precise in the following fashion. Consider a field ϕ in supergravity which couples to a local operator $\mathcal{O}$ in the world-volume. For the sake of having a mathematically well-defined problem, consider Wick-rotating both the gauge theory and the supergravity to a Euclidean

signature (this is a well-defined operation on the supergravity side because the near-horizon metric is conformal to flat space). Let us define $K[\phi_0]$ as the minimum of the supergravity action $S[\phi]$ in the throat, subject to Dirichlet boundary conditions $\phi = \phi_0$ imposed at the mouth. In the low-energy limit, this is equivalent to imposing asymptotic boundary conditions on supergravity fields in anti-de Sitter spacetime. Then the extension of (1.3) can be formulated as an identification of the free energy of the gauge theory with the supergravity action:

$$\begin{aligned} Z_{\text{YM}}[\phi_0] &= e^{-\beta F_{\text{YM}}[\phi_0]} = \left\langle \exp \left(- \int d^4x \phi_0 \mathcal{O} \right) \right\rangle \\ &= e^{-K[\phi_0]} = \underset{\phi=\phi_0 \text{ on } \partial \text{AdS}}{\text{maximum}} e^{-S[\phi]} . \end{aligned} \tag{1.4}$$

The main caveat to this identification is that the validity of classical supergravity requires that the number of D3-branes be large so that curvatures are small. More precisely the 't Hooft coupling $g_{\text{YM}}\sqrt{N}$ must be large. It is possible that quantum loop corrections to the supergravity action (obtained by replacing the maximum in the last expression by a formal path integral, evaluated via the saddle point approximation) can provide at least the leading $1/N^2$ corrections to the large N behavior of F_{YM} . A truly satisfactory treatment of such effects and the (possibly more important) α' corrections must be carried out via the full type IIB string theory in the near-horizon background geometry. Unfortunately, this geometry is supported by a Ramond-Ramond five-form flux, and it is not known how to incorporate such a field into the usual non-linear sigma model approach to string theory.

From the point of view of quantum field theory, however, an exciting consequence of (1.4) is that in the strong coupling limit, $\mathcal{N} = 4$ gauge theory seems to simplify to something which can be captured by a classical minimization problem.⁴ In particular, as will be suggested in chapter 8, all operators in the gauge theory except chiral primaries and their descendants (the operators corresponding to fields in the supergravity theory) may be expected to acquire a large mass dimension in the strong coupling limit. It is conceivable that they even drop out of the operator algebra in this limit, leaving us with a greatly reduced set of finite gauge-invariant Green's functions. It would be the realization of the dreams of decades to obtain simple expressions for these Green's functions, or at least simple equations for them.

⁴The following ideas arose in a discussion with Prof. T. Banks.

Chapter 2

Entropy considerations

A large segment of the recent string theory literature, starting with [5], is devoted to comparisons of the Bekenstein-Hawking entropy for a black hole or brane with entropies derived by counting states in some microscopic theory proposed to describe the black hole degrees of freedom. Finding the Bekenstein-Hawking entropy is straightforward, since it is one quarter of the horizon area in Planck units. As a rule, the microscopic calculations rely on either conformal invariance or the assumption of weak coupling to aid in the counting.

In section 2.1 we calculate the Bekenstein-Hawking entropy of clusters of coincident Dp -branes and compare with the free field theory result derived from the matter content of the world-volume theory. These results represent an extension of the work of [45] and have also appeared in various forms in the literature [46, 47, 48, 49].

In section 2.2 we exhibit a derivation of the effective string model of five-dimensional black holes starting from a D3-brane wrapped around a two-cycle of a four-dimensional compact manifold. The methods are similar to those developed in [27] to study effective string descriptions of four-dimensional black holes. Index theorems relating the dimensions of cohomology groups to integrals of characteristic classes are used to calculate the central charge of the CFT characterizing fluctuations of the effective string.

2.1 Thermodynamics of coincident branes

Coincident near-extremal black p -branes are described by the solution [50, 47]

$$\begin{aligned} ds^2 &= -A(r)dt^2 + B(r)dr^2 + C(r)r^2 d\Omega_{n+1}^2 + D(r)dy^i dy^i \\ &= -\frac{h}{\sqrt{f}}dt^2 + \frac{\sqrt{f}}{h}dr^2 + \sqrt{f}r^2 d\Omega_{n+1}^2 + \frac{1}{\sqrt{f}}dy^i dy^i \\ e^{2\phi} &= f^{(n-4)/2}, \end{aligned} \tag{2.1}$$

where

$$h = 1 - \frac{r_0^n}{r^n} \quad f = 1 + \frac{r_0^n \sinh^2 \alpha}{r^n}. \tag{2.2}$$

In (2.1), $ds^2 = G_{\mu\nu}dx^\mu dx^\nu$ is the string metric, related to the Einstein metric $g_{\mu\nu}$ by $g_{\mu\nu} = e^{-\phi/2}G_{\mu\nu}$. The y^i are the p spatial coordinates of the p -brane world-volume. For convenience we set $n = 7 - p$.

Newton's constant in uncompactified type II string theory is [51]

$$16\pi G = 2\kappa^2 = (2\pi)^7 g^2 (\alpha')^4. \tag{2.3}$$

The right hand side is the value at infinity, where the dilaton ϕ vanishes. With varying ϕ , the string coupling is really ge^ϕ . The Planck length ℓ_{Pl} therefore acquires a factor $e^{\phi/4}$. We have to include this factor when we calculate the Bekenstein-Hawking entropy: that entropy is roughly one “string bit” per unit Planck area ℓ_{Pl}^8 . More precisely,

$$S_{\text{cl}} = \frac{A_{\text{str}}}{4Ge^{2\phi}} = \frac{4\pi\Omega_{n+1}}{2\kappa^2} V r_0^{n+1} \cosh \alpha , \quad (2.4)$$

where A_{str} is measured using the string metric and Ω_{n+1} is the volume of the sphere S^{n+1} :

$$\Omega_m = \text{Vol } S^m = 2 \frac{\pi^{\frac{m+1}{2}}}{\Gamma(\frac{m+1}{2})} . \quad (2.5)$$

The mass of black p -branes was computed in general by Lu in [52]. To apply his results it is necessary to use the Einstein metric. The result is

$$M_{\text{ADM}} = \frac{\Omega_{n+1}V}{2\kappa^2} r_0^n \left(\frac{n+2}{2} + \frac{n}{2} \cosh 2\alpha \right) . \quad (2.6)$$

The non-extremal energy is defined as the difference between this mass and its extremal limit:

$$E_{\text{cl}} = M_{\text{ADM}} - M_0 = M_0 \left(\frac{n+2}{n} \text{csch } 2\alpha + \coth 2\alpha - 1 \right) \quad (2.7)$$

where

$$M_0 = \frac{Q_p V}{(2\pi)^p g \alpha^{\frac{p+1}{2}}} . \quad (2.8)$$

Lastly, one must consider the charge of the p -brane, which is quantized in a manner consistent with the Dirac quantization condition [50]. For BPS states (extremal branes), the charge μ_p is related to the tension τ_p :

$$2\kappa^2 \tau_p^2 = \mu_p^2 = (2\pi)^{7-2p} (\alpha')^{3-p} . \quad (2.9)$$

For non-extremal branes, one has the relation [47]

$$\frac{1}{2} r_0^n \sinh 2\alpha = \frac{2\kappa^2}{n\Omega_{n+1}} Q_p \tau_p = \frac{Q_p \sqrt{2\kappa^2 (2\pi)^{7-2p} (\alpha')^{3-p}}}{n\Omega_{n+1}} = \frac{(2\pi)^n}{n\Omega_{n+1}} \sqrt{\alpha'}^n g Q_p . \quad (2.10)$$

Given S and E , we can compute T from thermodynamics:

$$T_{\text{cl}} = \left(\frac{\partial E_{\text{cl}}}{\partial S_{\text{cl}}} \right)_{Q,V} = \frac{n}{4\pi r_0 \cosh \alpha} . \quad (2.11)$$

This value precisely matches with Hawking’s formula $T = \frac{\kappa}{2\pi}$ where κ is the surface gravity, computed according to the formula

$$\kappa = \frac{\partial A / \partial r}{2\sqrt{AB}} \Big|_{r=r_0} , \quad (2.12)$$

where A and B are the metric components appearing in (2.1). Note that (2.12) gives the same answer for the string metric as for the Einstein metric. This will always happen unless the dilaton does something singular on the horizon, which would seem unphysical because the horizon is supposed to be locally undetectable to an infalling observer.

By straightforward state counting, a free gas of $8Q_p^2$ species of bosons and an equal number of fermions at temperature T occupying a volume V can be shown to have free energy

$$F_{\text{free}} = -8Q_p^2 \frac{\Omega_{p-1}}{(2\pi)^p} \Gamma(p) \zeta(p+1) (2 - 2^{-p}) V T^{p+1} . \quad (2.13)$$

It follows that the energy and entropy are given by

$$E_{\text{free}} = -pF_{\text{free}} \quad S_{\text{free}} = -\frac{p+1}{T}F_{\text{free}} . \quad (2.14)$$

The $p = 3$ case agrees with the results of [45]. Note however that here we are taking V and T as the independent variables which are identified between GR and D-branes.

The quantity which can most directly be compared with field theory is the free energy:

$$F_{\text{cl}} = E_{\text{cl}} - T_{\text{cl}}S_{\text{cl}} = -M_0 \left(\frac{n-2}{n} \text{csch } 2\alpha - \coth 2\alpha + 1 \right) . \quad (2.15)$$

If we define χ by $F_{\text{cl}} = F_{\text{free}}\chi$, then

$$\chi = c \left[\frac{gQ_p}{(\sqrt{\alpha'}T)^{n-4}} \right]^{-\frac{n-4}{n-2}} \quad (2.16)$$

where c is a pure number:

$$c = \frac{4^{\frac{4n-3}{n-2}} (n-2) n^{\frac{-3n}{n-2}} \pi^{\frac{n^2-3n+10}{2(n-2)}} \Gamma\left(\frac{7-n}{2}\right) \Gamma\left(\frac{n+2}{2}\right)^{\frac{2}{n-2}}}{2^8 - 2^n \Gamma(7-n)\zeta(8-n)} . \quad (2.17)$$

When $n = 4$ (i.e. for the three-brane), $\chi = c = 3/4$, a result first obtained in [45, 53]. Results have been obtained in [46, 49]. For $n \neq 4$, the ratio in square brackets in (2.16) is precisely the 't Hooft coupling of the world-volume gauge theory multiplied by the power of the temperature needed to make it dimensionless. This has led to the conjecture [48] that the semi-classical result F_{cl} constitutes a prediction of gravity about the entropy of maximally supersymmetric gauge theory at strong coupling. More generally, one would expect that the exact free energy interpolates between F_{free} and F_{cl} as one proceeds from weak to strong coupling. It remains a mystery how one might make any concrete check on this interesting speculation.

2.2 Embedding of three-branes and the effective string

In [27], methods of algebraic geometry were used to make an exact count of the degrees of freedom on the effective string that underlies microscopic models of four-dimensional black holes [25, 26]. These powerful methods can be adapted to the analogous computation for five-dimensional black holes, where the effective string [5, 6] is a left-right symmetric conformal field theory with $(4, 4)$ supersymmetry. When the effective string is realized as Q_1 D1-branes bound to Q_5 D5-branes wound around T^5 , it was argued in [6, 54] that the number of left-moving bosons (real scalars) was $N_L^B = 4Q_1Q_5$. The central charge consequently was found to be $c = 6Q_1Q_5$, and the microscopic entropy of the conformal field theory (CFT) at level (N_L, N_R) ,

$$S = 2\pi\sqrt{Q_1Q_5}(\sqrt{N_L} + \sqrt{N_R}) , \quad (2.18)$$

agreed with the Bekenstein-Hawking entropy of the geometry with the same charges.

The purpose of this section is to obtain the formula $N_L^B = 4Q_1Q_5$ from a T-dual picture where the D1-D5 system is replaced by two sets of intersecting D3-branes. We will restrict ourselves to BPS brane configurations, with the understanding that once the central charge of the effective string CFT is established the general dilute-gas formula (2.18) follows immediately for near-extremal black holes as well as extremal ones. The extremal entropy calculation can be regarded as a subcase of [5] where the analysis is particularly simple.

The D3-brane must span the entire S^1 which the effective string wraps, so its embedding in the transverse T^4 is specified by a two-cycle. For a more general manifold M of complex dimension 2, the embedding of the D3-brane is a complex curve P . In order to preserve some supersymmetry, M must be either T^4 or K3, and P must be a holomorphic cycle. Using the same letter P to denote the dual of P in integral cohomology, the requirement of holomorphicity is $\bar{\partial}P = 0$. Carrying over the assumptions of [27] that M is much larger than the string scale, that S^1 is much larger yet, and that P is a very ample divisor of M , one is left with the fairly simple problem of determining the moduli of the effective string from Kaluza-Klein (KK) reduction of the degrees of freedom on a D3-brane wrapping a given homology cycle.

The real scalar bosons on the effective string come from two sources: the moduli of the D3-brane embedding and the D3-brane gauge field, KK-reduced to 1 + 1 dimensions. Explicitly,

$$N_L^B = d_P + b_1 , \quad (2.19)$$

where d_P is the number of real moduli of the embedding of P in M and b_1 is the first Betti number of P . The number of right-moving bosons is of course the same since the effective string in this case is left-right symmetric. Note in particular that b_1 , not the numbers $b_1^\pm$ of self-dual and anti-self-dual one-forms in $H^1(P)$, enter into the formulas for N_L^B and N_R^B . This is in contrast to the M5-brane case considered in [27]. The difference is that the field strength of the gauge field on the D3-brane is not required to be self-dual, whereas for the M5-brane it is.

Following the methods of [27], let us now compute d_P and b_1 . It is hoped that this simple application of the powerful techniques of algebraic geometry will lead to a more detailed understanding of the effective string.

To compute d_P one needs to give a description of the moduli space of embeddings of P in M . The holomorphic cycle P cannot be specified as the vanishing locus of a holomorphic function on M because holomorphic functions on M are constant. But it is possible to do something almost as good: one can specify P as the vanishing locus of a holomorphic section s of a holomorphic line bundle $\mathcal{L}$ over M . The convenience of this description is two-fold. First, the Chern class of $\mathcal{L}$ is $c(\mathcal{L}) = 1 + P$. And second, the moduli space is the (complex) projectivization of the Dolbeault cohomology space $H^0(M, \mathcal{L})$ [27]. This is because the zeroth Dolbeault space specifies precisely the different ways of picking a section s with $\bar{\partial}s = 0$; the projectivization is necessary because λs for nonzero complex λ specifies the same surface P as does s . Using an index theorem, one obtains

$$\begin{aligned} \sum_{i=0}^2 (-1)^i \dim H^i(M, \mathcal{L}) &= \int_M e^{c_1(\mathcal{L})} \text{td}(M) \\ &= \int_M \left(1 + P + \frac{1}{2}P^2\right) \left(1 + \frac{1}{12}c_2(M)\right) \\ &= \int_M \left(\frac{1}{2}P^2 + \frac{1}{12}c_2(M)\right) \\ &= D + \frac{1}{12}C_2(M) . \end{aligned} \quad (2.20)$$

The validity of the second line depends on $c_1(M) = 0$, which is guaranteed for K3 by SU(2) holonomy. Since M is a complex two-manifold, $C_2(M)$ is the Euler number $\chi(M)$: for T^4 this is 0 while for K3 it is 24. A detailed exposition of the properties of K3 can be found in [55], and [56] contains all the information regarding characteristic classes needed for the present discussion. In the third line of (2.20), D is the intersection number of P with itself. Given an integral basis α_A of $H^2(M, \mathbf{Z})$, the intersection form is

$$D_{AB} = \frac{1}{2} \int_M \alpha_A \wedge \alpha_B . \quad (2.21)$$

Expanding the cycle $P = \sum_A p^A \alpha_A$ in the dual basis of the homology group, one obtains $D = \frac{1}{2} D_{AB} p^A p^B$. For D3-brane configurations on T^4 T-dual to Q_1 D1-branes bound to Q_5 D5-branes, one

has $P = Q_1\alpha_1 + Q_5\alpha_5$ (as before, in homology) where α_1 and α_5 are two tori whose cross product is the full T^4 : the x^1 - x^2 torus and x^3 - x^4 torus, for example. Then $D = Q_1Q_5$.

Finally, one can use the observation [27] that all Dolbeault numbers but the zeroth vanish when P is very ample to obtain

$$d_p = 2D + \frac{1}{6}\chi(M) - 2. \quad (2.22)$$

The task of computing b_1 is conceptually more straightforward than d_p , since $b_1 = 2 - \chi(P)$ where $\chi(P)$ is the Euler characteristic of P . The Euler character of P is $c_1(P)$, and using the factorization property of Chern classes one can argue that $c_1(P) = -P$. Heuristically, the derivation of this equality rests on identifying fibers of $\mathcal{L}$ over a point of P as the part of M 's tangent space normal to P . Roughly speaking, when we take the tensor sum of normal and tangent bundles to P , we obtain the tangent bundle of M restricted to P , whose Chern class is simply 1. Chern classes multiply over tensor sums, so $c(P)c(\mathcal{L}) = 1$ and consequently $c(P) = 1 - P$. The subtleties we have ignored in this explanation are concisely dealt in [27] via an exact sequence argument. Now one can compute

$$\chi(P) = \int_P c_1(P) = - \int_P P = - \int_M P^2 = -2D \quad (2.23)$$

and obtain at last

$$b_1 = 2D + 2. \quad (2.24)$$

Putting everything together,

$$N_L^B = \frac{2}{3}c_L = 4D + \frac{1}{6}\chi(M) = \begin{cases} 4D & \text{for } M = T^4 \\ 4D + 4 & \text{for } M = K3. \end{cases} \quad (2.25)$$

Correction terms like $\frac{1}{6}\chi M$, subleading for large intersection number, were argued in [27] to correspond to supergravity loop corrections.

Chapter 3

D-brane approach to fixed scalars

3.1 Introduction

Many conventional wisdoms of general relativity are being reconsidered in the context of string theory simply because the string effective actions for gravity coupled to matter are more general than those considered in the past. One of the important differences is the presence of non-minimal scalar–gauge field couplings, leading to a breakdown of the “no hair” theorem (see the discussion in [57, 58]). Another new effect is the existence of certain scalars which, in the presence of an extremal charged black hole with regular horizon [59, 60, 61, 62, 63], acquire an effective potential [64, 65] which fixes their value at the horizon [66, 67, 68, 69]. These are the fixed scalars. The absorption of fixed scalars into $D = 4$ extremal black holes was considered in [70] and found to be suppressed compared to ordinary scalars: whereas the absorption cross-section of the latter approaches the horizon area A_h as $\omega \rightarrow 0$ [71], the fixed scalar cross-section was found to vanish as ω^2 .

This chapter is based on the paper [21]. Our main result is that the fixed scalar emission and absorption rates, as calculated using the methods of semi-classical gravity, are exactly reproduced by the effective string model of black holes based on intersecting D-branes. The D-brane description of the five-dimensional black holes involves n_1 1-branes and n_5 5-branes with some left-moving momentum along the intersection [5, 6]. The low-energy dynamics of the resulting bound state is believed to be well described by an effective string wound $n_1 n_5$ times around the compactification volume [72, 7, 8, 73, 74, 75]. This model has been successful in matching not only the extremal [5, 6] and near-extremal [54, 51, 72] entropies, but the rate of Hawking radiation of ordinary scalars as well [8, 73, 74, 75].

As part of our study, we have computed the semi-classical absorption cross-section of fixed scalars from both extremal and near-extremal $D = 5$ black holes. In general, we find cross-sections with a non-trivial energy dependence. In particular, for the extremal $D = 5$ black holes with two charges equal,

$$\sigma_{\text{abs}} = \frac{\pi^2}{2} R^2 r_K^3 \omega^2 \frac{\frac{\omega}{2T_L}}{1 - e^{-\frac{\omega}{2T_L}}} \left(1 + \frac{\omega^2}{16\pi^2 T_L^2} \right)$$

where r_K , $R \gg r_K$ and T_L are parameters related to the charges. At low energies the cross-section vanishes as ω^2 , just as in the $D = 4$ case studied in [70]. For non-extremal black holes, however, the cross-section no longer vanishes as $\omega \rightarrow 0$. For near-extremal $D = 5$ black holes, we find (for $\omega \sim T_H \ll T_L$)

$$\sigma_{\text{abs}}(\omega) = \frac{1}{4} A_h r_K^2 (\omega^2 + 4\pi^2 T_H^2) ,$$

where T_H is the Hawking temperature. A similar formula holds for the $D = 4$ case. Thus, even at low energies, the fixed scalar cross-section is sensitive to several features of the black hole geometry. By

comparison, the limiting value of the ordinary scalar cross-section is given by the horizon area alone. All of the complexities of the fixed scalar emission and absorption will be reproduced by, and find a simple explanation in, the effective string picture.

The absorption cross-section for ordinary scalars finds its explanation in the D-brane description in terms of the process $scalar \rightarrow L + R$ together with its time-reversal $L + R \rightarrow scalar$, where L and R represent left-moving and right-moving modes on the effective string [6, 7, 8, 73, 74, 75]. The absorption cross-section for fixed scalars is so interesting because, as we will show, it depends on the existence of eight kinematically permitted processes:

$$\begin{aligned} 1) & \quad scalar \rightarrow L + L + R + R \\ 2) & \quad scalar + L \rightarrow L + R + R \\ 3) & \quad scalar + R \rightarrow L + L + R \\ 4) & \quad scalar + L + R \rightarrow L + R \end{aligned} \tag{3.1}$$

and their time-reversals. One of the main results of this chapter is that competition among 1–4 and their time-reversals gives the following expression for the fixed scalar absorption cross-section,

$$\sigma_{\text{abs}}(\omega) = \frac{\pi r_1^2 r_5^2}{256 T_{\text{eff}}^2} \frac{\omega \left(e^{\frac{\omega}{T_H}} - 1 \right)}{\left(e^{\frac{\omega}{2T_L}} - 1 \right) \left(e^{\frac{\omega}{2T_R}} - 1 \right)} (\omega^2 + 16\pi^2 T_L^2)(\omega^2 + 16\pi^2 T_R^2), \tag{3.2}$$

where T_L and T_R are the left and right-moving temperatures, T_{eff} is the effective string tension [57, 58, 62, 72, 76, 77, 78, 79] and r_1^2 and r_5^2 are essentially the 1-brane and 5-brane charges. The only restriction on the validity of (3.2) is that $T_L, T_R, \omega \ll 1/r_1 \sim 1/r_5$ so that we stay in the dilute gas regime and keep the wavelength of the fixed scalar much larger than the longest length scale of the black hole. Remarkably, the very simple effective string result (3.2) is in complete agreement with the rather complicated calculations in semi-classical gravity! The semi-classical calculations involve no unknown parameters, so comparison with (3.2) allows us to infer T_{eff} . The result is in agreement with the fractional string tension necessary to explain the entropy of near-extremal 5-branes [76].

To set up the semi-classical calculations, we will develop in section 3.2 an effective action technique for deriving the equations of motion for fixed scalars. This technique shows how the fixed scalar equation couples with Einstein's equations when $r_1 \neq r_5$; therefore, we restrict ourselves to the regime $r_1 = r_5 = R$ where the fixed scalar equation is straightforward. We briefly digress to four dimensions, demonstrating how the same techniques lead to similar equations for fixed scalars. Clearly, comparisons analogous to the ones made in this chapter are possible for the four-dimensional case, where the effective string appears at the triple intersection of M-theory 5-branes [26]. In section 3.3 we use the Dirac-Born-Infeld (DBI) action to see how various scalars in $D = 5$ couple to the effective string. The main result of section 3.3 is that the leading coupling of the fixed scalar is to *four* fluctuation modes of the string. This highlights its difference from the moduli which couple to two fluctuation modes. In section 3.4 we return to five dimensions and exhibit approximate solutions to the fixed scalar equation, deriving the semi-classical emission and absorption rates. In section 3.5 we calculate the corresponding rates with D-brane methods, finding complete agreement with semi-classical gravity. We sum up in section 3.6.

3.2 Field theory effective action considerations

3.2.1 $D = 5$ case

First we shall concentrate on the case of a $D = 5$ black hole representing the bound state of n_1 RR strings and n_5 RR 5-branes compactified on a 5-torus [6]. This black hole may be viewed as a static

solution corresponding to the following truncation of type IIB superstring effective action compactified on 5-torus (cf. [80, 81])

$$S_5 = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} \left[R - \frac{4}{3}(\partial_\mu \phi_5)^2 - \frac{1}{4} G^{pl} G^{qn} (\partial_\mu G_{pq} \partial^\mu G_{ln} + e^{2\phi_5} \sqrt{G} \partial_\mu B_{pq} \partial^\mu B_{ln}) \right. \\ \left. - \frac{1}{4} e^{-\frac{4}{3}\phi_5} G_{pq} F_{\mu\nu}^{(K)p} F_{\mu\nu}^{(K)q} - \frac{1}{4} e^{\frac{2}{3}\phi_5} \sqrt{G} G^{pq} H_{\mu\nu p} H_{\mu\nu q} - \frac{1}{12} e^{\frac{4}{3}\phi_5} \sqrt{G} H_{\mu\nu\lambda}^2 \right], \quad (3.3)$$

where $\mu, \nu, \dots = 0, 1, \dots, 4; p, q, \dots = 5, \dots, 9$. ϕ_5 is the 5-d dilaton and G_{pq} is the metric of 5-torus,

$$\phi_5 \equiv \phi_{10} - \frac{1}{4} \ln G, \quad G = \det G_{pq},$$

and B_{pq} are the internal components of the RR 2-form field. $F_{\mu\nu}^{(K)p}$ is the Kaluza-Klein vector field strength, while $H_{\mu\nu p}$ and $H_{\mu\nu\lambda}$ are given explicitly by [80]

$$H_{\mu\nu p} = F_{\mu\nu p} - B_{pq} F_{\mu\nu}^{(K)q}, \quad F_p = dA_p, \quad F^{(K)p} = dA^{(K)p}, \quad (3.4)$$

$$H_{\mu\nu\lambda} = \partial_\mu B'_{\nu\lambda} - \frac{1}{2} A_\mu^{(K)p} F_{\nu\lambda p} - \frac{1}{2} A_{\mu p} F_{\nu\lambda}^{(K)p} + \text{cycles},$$

where $A_{\mu p} = B_{\mu p} + B_{pq} A_\mu^{(K)q}$ and $B'_{\mu\nu} = B_{\mu\nu} + A_{[\mu}^{(K)p} A_{\nu]p} - A_\mu^{(K)p} B_{pq} A_\nu^{(K)q}$ are related to the components of the $D = 10$ RR 2-form field B_{MN} .

The shifts in the field strengths in (3.4) will vanish for the black hole background considered below (for which the internal components of the 2-form B_{pq} will be zero and the two vector fields $A^{(K)p}$ and A_p will be electric), and, as it turns out, are also not relevant for the discussion of perturbations.

For comparison, a similar truncated $D = 5$ action with B_{pq} , $F_{\mu\nu p}$ and $H_{\mu\nu\lambda}$ from the NS-NS sector has the following antisymmetric tensor terms (the full action in general contains both RR and NS-NS antisymmetric tensor parts) [80]

$$- \frac{1}{4} G^{pl} G^{qn} \partial_\mu B_{pq} \partial^\mu B_{ln} - \frac{1}{4} e^{-\frac{4}{3}\phi_5} G^{pq} H_{\mu\nu p} H_{\mu\nu q} - \frac{1}{12} e^{-\frac{8}{3}\phi_5} H_{\mu\nu\lambda}^2.$$

We shall assume that there are non-trivial electric charges in only one of the five internal directions and that the metric corresponding to the internal 5-torus (over which the 5-brane will be wrapped) is

$$(ds_{10}^2)_{T^5} = e^{2\nu_5} dx_5^2 + e^{2\nu} (dx_6^2 + dx_7^2 + dx_8^2 + dx_9^2), \quad (3.5)$$

where x_5 is the string direction and ν is the scale of the four 5-brane directions transverse to the string. It is useful to introduce a different basis for the scalars, defining the six-dimensional dilaton, ϕ , and the scale λ of the x_5 (string) direction as measured in the $D = 6$ Einstein-frame metric:

$$\phi = \phi_{10} - 2\nu = \phi_5 + \frac{1}{2}\nu_5, \quad \lambda = \nu_5 - \frac{1}{2}\phi = \frac{3}{4}\nu_5 - \frac{1}{2}\phi_5. \quad (3.6)$$

The action (3.3) can be expressed either in terms of ϕ_5, ν_5, ν or ϕ, λ, ν (in both cases the kinetic term is diagonal). In the latter case (we set $B_{pq} = 0$)

$$S_5 = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} \left[R - (\partial_\mu \phi)^2 - \frac{4}{3} (\partial_\mu \lambda)^2 - 4 (\partial_\mu \nu)^2 \right. \\ \left. - \frac{1}{4} e^{\frac{8}{3}\lambda} F_{\mu\nu}^{(K)2} - \frac{1}{4} e^{-\frac{4}{3}\lambda + 4\nu} F_{\mu\nu}^2 - \frac{1}{12} e^{\frac{4}{3}\lambda + 4\nu} H_{\mu\nu\lambda}^2 \right]. \quad (3.7)$$

Here $F_{\mu\nu}^{(K)} \equiv F_{\mu\nu}^{(K)5}$ is the KK vector field strength corresponding to the string direction, while $F_{\mu\nu} \equiv F_{\mu\nu 5}$ and $H_{\mu\nu\lambda}$ correspond the “electric” (D1-brane) and “magnetic” (D5-brane) components of the field strength of the RR 2-form field. Evidently ϕ is an ordinary “decoupled” scalar while λ and ν are different: they interact with the gauge charges. We shall see that they are examples of the so-called “fixed scalars.”

To study spherically symmetric configurations corresponding to this action it is sufficient to choose the five-dimensional metric in the “2+3” form

$$ds_5^2 = g_{mn}dx^m dx^n + ds_3^2 = -e^{2a}dt^2 + e^{2b}dr^2 + e^{2c}d\Omega_3^2, \quad (3.8)$$

where a, b, c are functions of r and t . Solving first the equations for $H_{\mu\nu\lambda}$, $F_{\mu\nu}$ and $F_{\mu\nu}^{(K)}$ and assuming that the first two have, respectively, the magnetic and the electric components (with the charges P and Q corresponding to the D5-brane and the D1-brane), while the third has only the electric component with the Kaluza-Klein charge Q_K , we may eliminate them from the action (3.7). The result is an effective two-dimensional theory with coordinates $x^m = (t, r)$ and the action given (up to the constant prefactor) by¹

$$\begin{aligned} S_2 &= \int d^2x \sqrt{-g} e^{3c} \left[R + 6(\partial_m c)^2 - (\partial_m \phi)^2 - \frac{4}{3}(\partial_m \lambda)^2 - 4(\partial_m \nu)^2 + V(c, \nu, \lambda) \right] \\ &= \int dt dr \left[-e^{3c+b-a} (6\dot{c}\dot{b} + 6\dot{c}^2 - \dot{\phi}^2 - \frac{4}{3}\dot{\lambda}^2 - 4\dot{\nu}^2) \right. \\ &\quad \left. + e^{3c+a-b} (6c'a' + 6c'^2 - \phi'^2 - \frac{4}{3}\lambda'^2 - 4\nu'^2) \right. \\ &\quad \left. + 6e^{a+b+c} - 2e^{a+b-3c} f(\nu, \lambda) \right]. \end{aligned} \quad (3.9)$$

The first term in the potential originates from the curvature of the 3-sphere while the second is produced by the non-trivial charges,

$$f(\nu, \lambda) = Q_K^2 e^{-\frac{8}{3}\lambda} + e^{\frac{4}{3}\lambda} (P^2 e^{4\nu} + Q^2 e^{-4\nu}). \quad (3.10)$$

This is a special case of the more general expression following from (3.3): if the electric charges corresponding to the vector fields in (3.3) are Q_{Kp} and Q^p we get

$$f(\phi_5, G_{pq}) = e^{\frac{4}{3}\phi_5} Q_{Kp} Q_{Kq} G^{pq} + e^{-\frac{2}{3}\phi_5} \left(P^2 G^{1/2} + Q^p Q^q G_{pq} G^{-1/2} \right). \quad (3.11)$$

The potential f in (3.10) has the global minimum at $e^{4\nu} = QP^{-1}$, $e^{4\lambda} = Q_K^2 Q^{-1} P^{-1}$. These values of ν and λ are thus fixed points to which these fields are attracted on the horizon, which is why such fields can be called fixed scalars. By contrast, the decoupled scalar ϕ can be chosen to be equal to an arbitrary constant.

As an aside, we note that this structure of the potential (3.10) explains why one needs at least *three* different charges to get an extremal $D = 5$ black hole with a regular horizon (i.e. with scalar fields that do not blow up): it is necessary to have at least three exponential terms to “confine” the two fixed scalars. If the number of non-vanishing charges is smaller than three, then one or both scalars will blow up at the horizon.

Equivalent actions and potentials are found for theories that are obtained from the one above by U-duality. For example, in the case of the NS-NS truncation of type II action, which has a $D = 5$

¹The full set of equations and constraints is derived by first keeping the 2-d metric g_{mn} general and using its diagonal gauge-fixed form only after the variation. In addition to choosing g_{mn} diagonal as in (3.8), one can use the gauge freedom to impose one more relation between a and b .

black hole solution representing a bound state of NS-NS strings and solitonic 5-branes, we can put the action in the form (3.9), where λ is still the scale of the string direction as measured by the 6-d metric, while the roles of 2ν (the scale of the 4-torus) and $-\phi$ are interchanged.²

In order to find the static black hole solution to (3.9), we define $\rho = 2c + a$, $d\tau = -2e^{-3c-a+b}dr$. Now (3.9) reduces to a “particle” action (we choose $\phi = \text{const}$)

$$S_1 = \int d\tau \left[\frac{3}{2}(\partial_\tau \rho)^2 - \frac{3}{2}(\partial_\tau a)^2 - \frac{4}{3}(\partial_\tau \lambda)^2 - 4(\partial_\tau \nu)^2 + \frac{3}{2}e^{2\rho} - \frac{1}{2}e^{2a}f(\nu, \lambda) \right], \quad (3.12)$$

which should be supplemented by the “zero-energy” constraint,

$$\frac{3}{2}(\partial_\tau \rho)^2 - \frac{3}{2}(\partial_\tau a)^2 - \frac{4}{3}(\partial_\tau \lambda)^2 - 4(\partial_\tau \nu)^2 - \frac{3}{2}e^{2\rho} + \frac{1}{2}e^{2a}f(\nu, \lambda) = 0.$$

The special structure of f in (3.10) makes it possible to find a simple analytic solution of this Toda-type system. Introducing new variables $\alpha = a - \frac{4}{3}\lambda$, $\beta = a + \frac{2}{3}\lambda + 2\nu$, $\gamma = a + \frac{2}{3}\lambda - 2\nu$ and using the special form (3.10) of f , we can convert (3.12) to four non-interacting Liouville-like systems (related only through the constraint)

$$S_1 = \int d\tau \left[\frac{3}{2}(\partial_\tau \rho)^2 - \frac{1}{2}(\partial_\tau \alpha)^2 - \frac{1}{2}(\partial_\tau \beta)^2 - \frac{1}{2}(\partial_\tau \gamma)^2 + \frac{3}{2}e^{2\rho} - \frac{1}{2}Q_K^2 e^{2\alpha} - \frac{1}{2}P^2 e^{2\beta} - \frac{1}{2}Q^2 e^{2\gamma} \right]. \quad (3.13)$$

The general solution depends on the three gauge charges P, Q, Q_K and one parameter which we will call μ which governs the degree of non-extremality. In a convenient gauge, the solution reads [82, 6, 63, 51]

$$e^{2a} = h\mathcal{H}^{-2/3}, \quad e^{2b} = h^{-1}\mathcal{H}^{1/3}, \quad e^{2c} = r^2\mathcal{H}^{1/3}, \quad \mathcal{H} \equiv H_{\hat{P}}H_{\hat{Q}}H_{\hat{Q}_K}, \quad (3.14)$$

$$e^{2\lambda} = H_{\hat{Q}_K}(H_{\hat{Q}}H_{\hat{P}})^{-\frac{1}{2}}, \quad e^{4\nu} = H_{\hat{Q}}H_{\hat{P}}^{-1}, \quad e^{2\phi} = e^{2\phi_{10,\infty}}, \quad (3.15)$$

$$h = 1 - \frac{2\mu}{r^2}, \quad H_{\hat{q}} = 1 + \frac{\hat{q}}{r^2}, \quad \hat{q} \equiv \sqrt{q^2 + \mu^2} - \mu, \quad q = (P, Q, Q_K).$$

We have chosen the asymptotic values λ_∞ and ν_∞ to be zero. To compare with previous equations, we also note that $e^{2\rho} = r^2(r^2 - 2\mu)$.

In the extremal limit, $\mu = 0$, one finds

$$e^{-\alpha} = H_{Q_K}, \quad e^{-\beta} = H_P, \quad e^{-\gamma} = H_Q,$$

where $H_q = c_q + q\tau$ and $\tau = 1/r^2$. The constants c_{Q_K}, c_P, c_Q must satisfy $c_{Q_K}c_Pc_Q = 1$ in order for the 5-d metric to approach the Minkowski metric at infinity. The two remaining arbitrary constants correspond to the asymptotic values of λ and ν . As is clear from (3.9),(3.10), shifting λ and ν by constants is equivalent to a rescaling of Q_K, Q, P . The assumption that $\nu_\infty = 0$ and $\lambda_\infty = \nu_{5\infty} + \nu_\infty - \frac{1}{2}\phi_{10,\infty} = 0$ implies (setting $\alpha' = 1$):

$$V_4 = e^{4\nu_\infty} = 1, \quad \mathcal{R}^2 = e^{2\nu_{5\infty}} = g = e^{\phi_{10,\infty}}, \quad \kappa_5^2 = \frac{2\pi^2 g^2}{\mathcal{R}V_4}, \quad (3.16)$$

where $(2\pi)^4 V_4$ is the volume of T^4 in the (6789) directions, while $\mathcal{R}$ is the radius of the circle in direction 5. Then the charges Q_K, Q, P are related to the quantized charges n_1, n_5, n_K as follows:

$$n_1 = \frac{V_4 Q}{g} = \frac{Q}{g}, \quad n_5 = \frac{P}{g}, \quad n_K = \frac{\mathcal{R}^2 V_4 Q_K}{g^2} = \frac{Q_K}{g}. \quad (3.17)$$

²There exists an equivalent representation of this NS-NS action where the fixed scalars are the 5-d dilaton and the scale of the string direction, while the scale of the 4-torus is decoupled.

The somewhat unusual form of the last relation is due to our choice $\lambda_\infty = 0$ instead of more standard $\nu_{5\infty} = 0$.

In using the black hole solution (3.14), (3.15), we will often find it convenient to work in terms of characteristic radii rather than the charges, so we define

$$r_1^2 = \hat{Q} , \quad r_5^2 = \hat{P} , \quad r_K^2 = \hat{Q}_K , \quad r_0^2 = 2\mu . \quad (3.18)$$

From the classical GR point of view, these parameters can take on any values. Recent experience has shown, however, that when the radii satisfy [75]

$$r_0, r_K \ll r_1, r_5 \quad (3.19)$$

the black hole can be successfully matched to a bound state of D1-branes and D5-branes (with no anti-branes present) carrying a dilute gas of massless excitations propagating along the bound D1-branes. Evidence for this gas can be seen directly in the energy, entropy and temperature of the black hole solution. Introducing a new parameter σ through

$$r_K^2 = r_0^2 \sinh^2 \sigma$$

one finds the following expressions [51, 75] for the ADM mass, Hawking temperature and the entropy in the parameter region (3.19):

$$M = \frac{2\pi^2}{\kappa_5^2} \left(r_1^2 + r_5^2 + \frac{1}{2} r_0^2 \cosh 2\sigma \right) , \quad T_H^{-1} = \frac{2\pi r_1 r_5}{r_0} \cosh \sigma , \quad (3.20)$$

$$S = \frac{2\pi A_h}{\kappa_5^2} = \frac{4\pi^3}{\kappa_5^2} r_1 r_5 r_0 \cosh \sigma .$$

The entropy and energy are those of a gas of massless one-dimensional particles with the left-movers and right-movers each having its own temperature [75]:

$$T_L = \frac{r_0 e^\sigma}{2\pi r_1 r_5} , \quad T_R = \frac{r_0 e^{-\sigma}}{2\pi r_1 r_5} . \quad (3.21)$$

The Hawking temperature is related to these two temperatures by

$$\frac{2}{T_H} = \frac{1}{T_L} + \frac{1}{T_R} , \quad (3.22)$$

a fact which also has a natural thermodynamic interpretation. These results will be heavily used in later comparisons of classical GR results with D-brane calculations of corresponding quantities.

Let us now turn to the discussion of the propagation of perturbations on this black hole background. The goal will be to calculate the classical absorption cross-section of various scalar fields and eventually to compare them with comparable D-brane quantities. The behavior of “free” scalars, like ϕ , is quite different from that of “fixed” scalars, like λ and ν . The spherically symmetric fluctuations of ϕ obey the standard massless Klein-Gordon equation in this background. Namely, if $\delta\phi = e^{i\omega t} \tilde{\phi}(r)$, then

$$\left[r^{-3} \frac{d}{dr} \left(h r^3 \frac{d}{dr} \right) + \omega^2 h^{-1} H_{\hat{P}} H_{\hat{Q}} H_{\hat{Q}_K} \right] \tilde{\phi} = 0 , \quad (3.23)$$

This scattering problem, and its D-brane analog, have been analyzed at length recently and we will have no more to say about it. The spherically symmetric fluctuations of the metric functions a, b, c and the scalars λ, ν in general obey a complicated set of coupled differential equations.³ However,

³The spherically symmetric fluctuations of the gauge fields need not be considered explicitly: since the dependence on $H_{\mu\nu\lambda}$ and $F_{\mu\nu}$ is gaussian, they are automatically included when going from (3.7) to (3.9).

when the charges P and Q are set equal, a dramatic simplification occurs: the background value of ν in (3.15) (i.e. the scale of the transverse 4-torus) becomes constant and its small fluctuations $\delta\nu$ decouple from those of the other fields.⁴ The gaussian effective action for $\delta\nu$ extracted from (3.9) is

$$\delta S_2 = \int d^2x \sqrt{-g} e^{3c} \left[-4(\partial_m \delta\nu)^2 - 32P^2 e^{-6c + \frac{4}{3}\lambda} (\delta\nu)^2 + \dots \right] \quad (3.24)$$

and spherically symmetric fluctuations $\delta\nu = e^{i\omega t} \tilde{\nu}$ obey

$$\left[r^{-3} \frac{d}{dr} \left(h r^3 \frac{d}{dr} \right) + \omega^2 h^{-1} H_{\hat{P}}^2 H_{\hat{Q}_K} - 8P^2 r^{-6} H_{\hat{P}}^{-2} \right] \tilde{\nu} = 0 . \quad (3.25)$$

This is the standard Klein-Gordon equation (3.23) augmented by a space-dependent mass term originating from the expansion of the effective potential $f(\nu, \lambda)$ in (3.10). This mass term falls off as r^{-6} at large r , and, in the extremal case, blows up like $8/r^2$ near the horizon at $r = 0$. This is the $l(l+2)/r^2$ angular momentum barrier for an $l = 2$ partial wave in $D = 5$. This “transmutation” of angular momentum plays an important role in the behavior of the fixed scalar cross-section. For later analysis, it will be convenient to rewrite this equation using the coordinate $\tau = 1/r^2$:

$$\left[\left[(1 - 2\mu\tau) \frac{d}{d\tau} \right]^2 + \frac{1}{4} \omega^2 \tau^{-3} (1 + \hat{P}\tau)^2 (1 + \hat{Q}_K\tau) - 2 \frac{P^2 (1 - 2\mu\tau)}{(1 + \hat{P}\tau)^2} \right] \tilde{\nu} = 0 . \quad (3.26)$$

Note that, when all the three charges are equal, $P = Q = Q_K$, the background value of the other scalar, λ , is constant as well. Then the small fluctuations of this field decouple from gravitational perturbations and satisfy the same equation as ν , (3.25). If only two of the charges are equal, then there is only one fixed scalar which has a constant background value and decouples from gravitational perturbations. We would also like to know the fixed scalar scattering equations (and solutions) for the general $Q_K \neq Q \neq P$ black hole. This problem is surprisingly complicated due to mixing with gravitational perturbations. The disentanglement of the equations was achieved in [83].

To summarize, we have identified a set of scalars around the familiar type II string $D = 5$ black hole solution which merit the name of “fixed scalars” in that their horizon values are fixed by the background charges. Their fluctuations in the black hole background satisfy the Klein-Gordon equation, augmented by a position-dependent mass term. In section 3.4 we will solve the new equations to find the absorption cross-section by the black hole for these special scalars.

3.2.2 $D = 4$ case

Previous experience [84, 85, 26, 86, 74, 87] suggests that one may be able to extend the $D = 5$ successes in reproducing entropies and radiation rates with D-brane methods to $D = 4$ black holes carrying 4 charges. Although we will not pursue the p -brane approach to $D = 4$ black hole dynamics in this chapter, this is a natural place to discuss $D = 4$ fixed scalars and to record their scattering equations for later use.

A convenient representation of the $D = 4$ black hole with four different charges [60, 61, 62] is the $D = 11$ supergravity configuration $5 \perp 5 \perp 5$ of three 5-branes intersecting over a common string [26, 88]. The three magnetic charges are related to the numbers of 5-branes in three different hyperplanes, while the electric charge has Kaluza-Klein origin. The reduction to $D = 4$ of the relevant part of the $D = 11$ supergravity (or $D = 10$ type IIA) action has the form

$$S_4 = \frac{1}{2\kappa_4^2} \int d^4x \sqrt{-g} \left[R - 2(\partial_\mu \nu)^2 - \frac{3}{2}(\partial_\mu \zeta)^2 - \frac{4}{3}(\partial_\mu \xi)^2 - (\partial_\mu \eta)^2 \right] \quad (3.27)$$

⁴Similar simplification occurs when any two of the three charges are equal. For example, if $P = Q_K$ we may introduce $\lambda' = -\frac{1}{2}(\lambda - 3\nu)$, $\nu' = -\frac{1}{2}(\nu + \lambda)$ (in terms of which the kinetic part in the action (3.9) preserves its diagonal form) to discover that ν' has decoupled fluctuations. The resulting equation for $\delta\nu'$ has the same form as the equation for $\delta\nu$ in the case of $P = Q$.

$$-\frac{1}{4}e^{3\zeta}(F_{\mu\nu}^{(K)})^2 - \frac{1}{4}e^\zeta \left(e^{-\frac{8}{3}\xi}(F_{\mu\nu}^{(1)})^2 + e^{\frac{4}{3}\xi}[e^{2\eta}(F_{\mu\nu}^{(2)})^2 + e^{-2\eta}(F_{\mu\nu}^{(3)})^2] \right) \Big] .$$

The scalar fields are expressed in terms of components of the internal 7-torus part of the $D = 11$ metric. By the logic of the previous section, the scale ν of the 6-torus transverse to the intersection string is a decoupled scalar, while the fields ζ, ξ, η (related to the scale of the string direction and the ratios of sizes of 2-tori shared by pairs of 5-branes) are fixed scalars. If the internal part of the $D = 11$ metric is

$$ds_7^2 = e^{2\nu_4} dx_4^2 + e^{2\nu_1}(dx_5^2 + dx_6^2) + e^{2\nu_2}(dx_7^2 + dx_8^2) + e^{2\nu_3}(dx_9^2 + dx_{10}^2) ,$$

where x_4 is the direction of the common string, then

$$\nu = \nu_1 + \nu_2 + \nu_3, \quad \xi = \nu_1 - \frac{1}{2}\nu_2 - \frac{1}{2}\nu_3, \quad \eta = \nu_3 - \nu_2, \quad \zeta = \nu_4 + \frac{2}{3}(\nu_1 + \nu_2 + \nu_3).$$

Using an ansatz for the 4-d metric similar to (3.8), solving for the vector fields, and substituting the result back into the action, we get the following effective two-dimensional action (cf. (3.9))

$$S_2 = \int d^2x \sqrt{-g} e^{2c} \left[R + 2(\partial_m c)^2 - 2(\partial_\mu \nu)^2 - \frac{3}{2}(\partial_m \zeta)^2 - \frac{4}{3}(\partial_m \xi)^2 - (\partial_m \eta)^2 \right. \\ \left. + 2e^{-2c} - \frac{1}{2}e^{-4c} f(\zeta, \xi, \eta) \right] , \quad (3.28)$$

where

$$f(\zeta, \eta, \xi) = Q_K^2 e^{-3\zeta} + e^\zeta \left[P_1^2 e^{-\frac{8}{3}\xi} + e^{\frac{4}{3}\xi} (P_2^2 e^{2\eta} + P_3^2 e^{-2\eta}) \right] . \quad (3.29)$$

As in the $D = 5$ case, one finds that the special structure of f makes it possible to diagonalize the interaction term by a field redefinition and thus find the static solution in a simple factorized form [60, 61, 62, 88] (cf. (3.14))

$$ds_4^2 = -e^{2a} dt^2 + e^{2b} dr^2 + e^{2c} d\Omega_2^3 , \quad (3.30)$$

$$e^{2a} = h \mathcal{H}^{-1/2} , \quad e^{2b} = h^{-1} \mathcal{H}^{1/2} , \quad e^{2c} = r^2 \mathcal{H}^{1/2} , \quad \mathcal{H} \equiv H_{\hat{Q}_K} H_{\hat{P}_1} H_{\hat{P}_2} H_{\hat{P}_3} , \\ e^{2\eta} = H_{\hat{P}_3} H_{\hat{P}_2}^{-1} , \quad e^{2\xi} = H_{\hat{P}_1} (H_{\hat{P}_2} H_{\hat{P}_3})^{-1/2} , \quad e^{2\zeta} = H_{\hat{Q}_K} (H_{\hat{P}_1} H_{\hat{P}_2} H_{\hat{P}_3})^{-1/3} , \quad (3.31)$$

$$h = 1 - \frac{2\mu}{r} , \quad H_{\hat{q}} = 1 + \frac{\hat{q}}{r} , \quad \hat{q} \equiv \sqrt{q^2 + \mu^2} - \mu , \quad q = (Q_K, P_1, P_2, P_3) .$$

As in the $D = 5$ case, for the generic values of charges the spherically symmetric perturbations of this solution obey a complicated system of equations (for discussions of perturbations of single-charged dilatonic black holes see, e.g., [89, 43, 90, 91]). However, when the three magnetic charges are equal, η and ξ have constant background values, and so their small spherically-symmetric fluctuations decouple from the metric perturbations,

$$\delta S_2 = \int d^2x \sqrt{-g} e^{2c} \left[-(\partial_m \delta \eta)^2 - 2P^2 e^{-4c+\zeta+\frac{4}{3}\xi} (\delta \eta)^2 + \dots \right] , \quad (3.32)$$

leading to the following radial Klein-Gordon equation with an extra mass term:

$$\left[r^{-2} \frac{d}{dr} (h r^2 \frac{d}{dr}) + \omega^2 h^{-1} H_{\hat{P}}^3 H_{\hat{Q}_K} - 2P^2 r^{-4} H_{\hat{P}}^{-2} \right] \tilde{\eta} = 0 . \quad (3.33)$$

Here $\delta \eta(r, t) = e^{i\omega t} \tilde{\eta}(r)$; cf. (3.25). The same universal equation is found for $\delta \xi$. In terms of $\tau = 1/r$ this becomes

$$\left[[(1 - 2\mu\tau) \frac{d}{d\tau}]^2 + \omega^2 \tau^{-2} (1 + \hat{P}\tau)^3 (1 + \hat{Q}_K\tau) - 2 \frac{P^2 (1 - 2\mu\tau)}{(1 + \hat{P}\tau)^2} \right] \tilde{\eta} = 0 . \quad (3.34)$$

Represented in this form this is very similar to (3.26) found in the $D = 5$ case: the differential operator and mass terms are exactly the same, while the frequency terms are related by $\omega \rightarrow 2\omega$, $\tau^{-3}(1 + \hat{P}\tau)^2 \rightarrow \tau^{-2}(1 + \hat{P}\tau)^3$.

In the extremal case and with all four charges chosen to be equal, $Q_K = P$, (3.33) reduces to the equation studied in [70]. The characteristic coefficient 2 in the mass term gives the effective potential of the form $l(l+1)/r^2$ near the horizon, with $l = 1$. Away from the horizon, the fixed scalar equation differs from that of the $l = 1$ partial wave of the ordinary scalar.

Finally, let us note that there exist other representations of the 4-charge $D = 4$ black hole. For example in the case of the $2\perp 2\perp 5\perp 5$ representation [26], or, equivalently, the U-dual $D = 4$ configuration in the NS-NS sector with two (electric and magnetic) charges coming from the $D = 10$ antisymmetric tensor and two (electric and magnetic) charges being of Kaluza-Klein origin, we may parametrize the metric as

$$ds_{10}^2 = e^{2\nu_4} dx_4^2 + e^{2\nu_5} dx_5^2 + e^{2\nu} (dx_6^2 + dx_7^2 + dx_8^2 + dx_9^2) .$$

This leads to the effective Lagrangian related to the above one (3.28) by a linear field redefinition and re-interpretation of the charges. The potential is (cf. (3.29))

$$f(\phi, \nu_4, \nu_5) = e^{2\phi} (Q_K^2 e^{2\nu_4} + Q^2 e^{-2\nu_4}) + e^{-2\phi} (P_K^2 e^{2\nu_5} + P^2 e^{-2\nu_5}) ,$$

where ϕ is the 4-d dilaton. This shows that the scale of the remaining 4-torus, ν , decouples.

3.3 Effective string couplings

We now turn to a discussion of the effective action governing the absorption and emission of fixed scalars by the bound state of D1- and D5-branes. We use the same framework as the recent demonstrations of agreement between GR and D-brane treatments of the absorption of generic decoupled scalars [6, 7, 8, 73, 74]. We assume that: (i) the $D = 5$ black hole is equivalent to n_1 D1-branes bound to n_5 D5-branes, with some left-moving momentum; (ii) that the low-energy dynamics of this system is described by the DBI action for a string with an effective tension T_{eff} , and (iii) that the relevant bosonic oscillations of this effective string are only in the four 5-brane directions ($i = 6, 7, 8, 9$) transverse to the 1-brane. These assumptions serve to specify the detailed couplings of external closed string fields, in particular the fixed scalars, to the D-brane degrees of freedom. This is an essential input to any calculation of absorption and emission rates and, as we shall see, brings fairly subtle features of the effective action into play.

Specifically, we assume that the low-energy excitations of our system are described by the standard D-string action

$$I = -T_{\text{eff}} \int d^2\sigma e^{-\phi_{10}} \sqrt{-\det \gamma_{ab}} + \dots , \quad \gamma_{ab} = G_{\mu\nu}(X) \partial_a X^\mu \partial_b X^\nu , \quad (3.35)$$

where ϕ_{10} and $G_{\mu\nu}$ are the $D = 10$ dilaton and string-frame metric. The specific dependence on ϕ_{10} is motivated by the expected $1/g_{\text{str}}$ behavior of the D-string tension. The normalization constant of the tension, T_{eff} , is subtle and will be discussed later. Our goal is to read off the couplings between excitations of the effective string and the fluctuations of the metric and dilaton that correspond to the fixed scalars.

It should be noted that the essential structure of the effective string action we are interested in can be, at least qualitatively, understood using semi-classical effective field theory methods. A straightforward generalization of the extremal classical solution [82, 6] describing a BPS bound state of a string and 5-brane in which the string is localized on the 5-brane⁵ has the $D = 10$ string metric

⁵Instead of talking about a bound state of several single-charged D-strings and D5-branes with coinciding centers, it is sufficient, at the classical level, to consider just a single string and a single 5-brane having charges $Q \sim n_1$ and $P \sim n_5$.

($m = 1, 2, 3, 4; i = 6, 7, 8, 9$)

$$ds_{10}^2 = H_1^{-1/2} H_5^{-1/2} (-dt^2 + dx_5^2) + H_1^{1/2} H_5^{1/2} dx_m dx_m + H_1^{1/2} H_5^{-1/2} dx_i dx_i ,$$

where $H_5 = H_5(x_m) = 1 + P/x_m^2$ and $H_1 = H_1(x_m, x_i)$ is a solution of

$$[\partial^m \partial_m + H_5(x_n) \partial^i \partial_i] H_1 = 0$$

such that for $P \rightarrow 0$ it approaches the standard string harmonic function, $H_1 \rightarrow 1 + Q/(x_i^2 + x_m^2)^3$. If one averages over the x_i -dependence of H_1 one returns to the original delocalized case, $H_1 = 1 + Q/x_m^2$, which corresponds to the extremal limit of the $D = 5$ black hole (3.14), (3.15) with $Q_K = 0$ (here we consider the unboosted string). The presence of the 5-brane breaks the $O(1, 1) \times O(8)$ symmetry of the standard RR string solution down to $O(1, 1) \times O(4) \times O(4)$. Since the localized solution also breaks 4+4 translational invariances, the string soliton has 4+4 collective coordinates: $X^m(x_5, t)$, $X^i(x_5, t)$. The corresponding $O(4) \times O(4)$ invariant effective string action thus should have the following form in the static gauge,

$$I = \int d^2\sigma [T_0 + T_{\parallel} \partial^a X^i \partial_a X^i + T_{\perp} \partial^a X^m \partial_a X^m + \dots] .$$

The constants $T_0, T_{\parallel}, T_{\perp}$ can be determined using standard methods (see, e.g., [92]) by substituting the perturbed solution into the $D = 10$ effective field theory action, etc. T_0 is proportional to the ADM mass of the background, $T_0 \sim P + Q$. The same should be true also for $T_{\perp}$, $T_{\perp} \sim P + Q$, since X^m describe oscillations of the whole bound state in the common transverse 4-space. At the same time, $T_{\parallel}$ is the effective tension of the string within the 5-brane, so that $T_{\parallel} \sim Q$. In the special cases $P = 0$ and $Q = 0$ these expressions are in obvious agreement with the standard results for a free string and a free 5-brane. In the case when $P \gg Q$, i.e. $n_5 \gg n_1$, we learn that $T_{\perp} \gg T_{\parallel}$, so that oscillations of the string in the four directions X^m transverse to the 5-brane can be ignored.⁶ If we further assume, following [72, 94], that the string lying within the 5-brane has the effective length $L_{\text{eff}} \sim QP \sim n_1 n_5$, we finish with the following expression for the effective string tension in the relevant directions parallel to the 5-brane: $T_{\text{eff}} \sim T_{\parallel}/L_{\text{eff}} \sim 1/P \sim 1/n_5$. This picture is consistent with that suggested in [72, 76] and will pass a non-trivial test below.

In accord with the assumption (iii), we thus drop terms involving derivatives of $X^m = X^{1,2,3,4}$ (i.e. motions in the uncompactified directions). We also eliminate two more string coordinates by choosing the static gauge $X^0 = \sigma^0$, $X^5 = \sigma^1$ and write

$$\gamma_{ab} \equiv \eta_{ab} + \hat{\gamma}_{ab} = G_{ab}(x) + 2G_{i(a}(x) \partial_{b)} X^i + G_{ij}(x) \partial_a X^i \partial_b X^j . \quad (3.36)$$

We make the Kaluza-Klein assumption that the background fields ϕ_{10} and $G_{\mu\nu}$ depend only on the external coordinates $x^m \equiv X^m, m = 0, 1, \dots, 5$. Since we are interested in linear absorption and emission processes, we make a weak-field expansion in powers of ϕ_{10} and $h_{\mu\nu} \equiv G_{\mu\nu} - \eta_{\mu\nu}$, splitting h_{ij} into traceless and trace parts: $h_{ij} = \bar{h}_{ij} + \frac{1}{4} \delta_{ij} h$. Finally, we distinguish L and R string excitations by introducing $\partial_+ = \partial_0 + \partial_1$, $\partial_- = -\partial_0 + \partial_1$. We can then use the formula

$$\sqrt{-\det(\eta_{ab} + \hat{\gamma}_{ab})} = 1 + \frac{1}{2} \hat{\gamma}_{+-} - \frac{1}{8} \hat{\gamma}_{++} \hat{\gamma}_{--} + \frac{1}{16} \hat{\gamma}_{+-} \hat{\gamma}_{++} \hat{\gamma}_{--} + \dots$$

⁶One may also give the following argument in support of the claim that transverse oscillations of the string can be ignored when the string is light compared to the 5-brane. The classical action for a D-string probe moving in the above background produced by a bound state of R-R string and 5-brane has the following form:

$I_1 = T_0 \int d^2\sigma [e^{-\phi} \sqrt{-\det(G_{\mu\nu} + B_{\mu\nu}^{(NS)})} + B^{(R)} + \dots]$ (we shall set the world-sheet gauge field to zero and choose the static gauge). If the string is oriented parallel to the x^5 direction of the 5-brane (a BPS configuration), the non-trivial part of the potential term cancels out [93]. The same is true for the dependence on the H_1 -function in the second-derivative terms which have the form $I_1 = T_0 \int d^2\sigma [1 + \frac{1}{2} \partial^a X^i \partial_a X^i + \frac{1}{2} H_5(X) \partial^a X^m \partial_a X^m + \dots]$. The function $H_5 = 1 + \frac{P}{x_m^2}$ thus determines the metric of the transverse part of the moduli space (see also [11]), i.e. it plays the role of an effective $T_{\perp}$ which blows up when the string approaches the 5-brane. Thus the string probe can freely move within the 5-brane, but its transverse motions are suppressed.

to expand (3.35), finding the following action for X^i :

$$I_X = -T_{\text{eff}} \int d^2\sigma \left[1 + \frac{1}{2} \partial_+ X \partial_- X - \frac{1}{8} (\partial_+ X)^2 (\partial_- X)^2 + \dots \right. \quad (3.37)$$

$$\left. + L_0 + L_1 + L_2 + L_3 + L_4 + L'_4 + \dots \right] ,$$

$$L_0 = \frac{1}{2} (h_{55} - h_{00}) - \phi_{10} , \quad L_1 = \frac{1}{2} h_{5i} (\partial_+ + \partial_-) X^i , \quad (3.38)$$

$$L_2 = -\frac{1}{2} \phi \partial_+ X^i \partial_- X^i + \frac{1}{2} \bar{h}_{ij} \partial_+ X^i \partial_- X^j - \frac{1}{8} (h_{00} + h_{55}) [(\partial_+ X)^2 + (\partial_- X)^2] , \quad (3.39)$$

$$L_3 = -\frac{1}{4} h_{5i} [\partial_- X^i (\partial_+ X)^2 + \partial_+ X^i (\partial_- X)^2] , \quad (3.40)$$

$$L_4 = \frac{1}{8} (\phi_{10} - \frac{1}{2} h) (\partial_+ X)^2 (\partial_- X)^2 = \frac{1}{8} \phi (\partial_+ X)^2 (\partial_- X)^2 - \frac{1}{4} \nu (\partial_+ X)^2 (\partial_- X)^2 , \quad (3.41)$$

$$\begin{aligned} L'_4 &= \frac{1}{16} (h_{55} + h_{00}) \partial_+ X \partial_- X [(\partial_+ X)^2 + (\partial_- X)^2] + \frac{1}{16} (h_{55} - h_{00}) (\partial_+ X)^2 (\partial_- X)^2 \\ &= \frac{1}{8} \lambda [\partial_+ X \partial_- X ((\partial_+ X)^2 + (\partial_- X)^2) + (\partial_+ X)^2 (\partial_- X)^2] \\ &\quad + \frac{1}{16} \phi [\partial_+ X \partial_- X ((\partial_+ X)^2 + (\partial_- X)^2) + (\partial_+ X)^2 (\partial_- X)^2] + O(h_{00}) . \end{aligned} \quad (3.42)$$

The expansion has been organized in powers of derivatives of X^i and we have kept terms at most linear in the external fields (since we don't use them in what follows, we have dropped higher-order terms involving $\bar{h}_{ij}$). We have also reorganized those fields in a way appropriate for the compactification on a five-torus:

$$\nu \equiv \frac{1}{8} h , \quad \phi \equiv \phi_{10} - \frac{1}{4} h , \quad \lambda \equiv \frac{1}{2} h_{55} + \frac{1}{8} h - \frac{1}{2} \phi_{10} = \frac{1}{2} h_{55} - \frac{1}{2} \phi , \quad h \equiv h_{ii} , \quad (3.43)$$

where ν is the scale of the 4-torus part of the 5-brane (if $G_{ij} = e^{2\nu} \delta_{ij}$ then, in the linearized approximation, $h_{ii} = 8\nu$), ϕ is the corresponding six-dimensional dilaton and λ is the scale of the fifth (string) dimension measured in the six-dimensional Einstein metric. These are the same three scalar fields that appear in the GR effective action (3.7), (3.9).

Since the kinetic terms in the effective action (3.7), (3.9) are diagonal in ϕ , ν and λ , we can immediately read off some important conclusions from (3.39), (3.41). The expansion in powers of world-sheet derivatives is a low-energy expansion and, of the fields we have kept, only the dilaton ϕ is coupled at leading order. It is also easy to see that the off-diagonal components of the metric $\bar{h}_{ij}$ have the same coupling as ϕ to lowest order in energy. (These are the fields whose emission and absorption were considered in [7, 8, 73]). What is more interesting is that the scalar ν only couples at the next-to-leading order (fourth order in derivatives). Note that its interaction term can be written in terms of world-sheet energy-momentum tensors as $\nu T_{++}^X T_{--}^X$.⁷ The scalar λ likewise does not get emitted at the leading order and does couple at the same order as ϕ , but with a different vertex.⁸ The graviton components in the time and string directions h_{00} and h_{55} couple to the string in a way similar to λ , which reflects their mixing with λ in the effective action (3.9). Indeed, the vertex $(h_{00} + h_{55})(T_{++}^X + T_{--}^X)$ gives a vanishing contribution to the amplitude of production of a closed string state: it only couples a pair of left-movers or a pair of right-movers so that the production is forbidden kinematically. The important (and non-trivial) point is that the simplest DBI action

⁷There is a similar coupling for ϕ which produces a subleading correction to its emission rate.

⁸The different vertices for ν and λ probably reflect the different behavior of their fluctuations (the non-decoupling of $\delta\lambda$ from metric perturbations) in the case when Q_K is not equal to $Q = P$.

for the coupling of the external fields to the D-brane gives the fields ν and λ , previously identified as the “fixed scalars”, different (and weaker) couplings to the effective string than the fields like ϕ previously identified as “decoupled scalars.” The precise couplings will shortly be used to make precise calculations of absorption and emission rates.

The action (3.37) is at best the bosonic part of a supersymmetric action. In previous discussions of D-brane emission and absorption, it has been possible to ignore the coupling of external fields to the massless fermionic excitations of the D-brane. For the questions that interest us, that will no longer be possible and we make a specific proposal for the couplings of world-sheet fermions. In the successful D-brane description of the entropy of rotating black holes [38, 95], five-dimensional angular momentum is carried by the fermions alone. There are two world-sheet fermion doublets, one right-moving and one left-moving. The $SO(4)$ rotations of the uncompactified (1234) coordinates are decomposed as $SU(2)_L \times SU(2)_R$ and the obvious (and correct) choice is to take the left-moving fermions, S^a , to be a doublet under $SU(2)_L$ and the right-moving fermions, $S^{\dot{a}}$ to be a doublet under $SU(2)_R$. This set of fermions may be bosonized as two boson fields, φ^1 and φ^2 . As mentioned above, the next-to-leading terms in (3.40), (3.41) can be written in terms of the X -field stress-energy tensor,

$$T_{++}^X = (\partial_+ \vec{X})^2, \quad T_{--}^X = (\partial_- \vec{X})^2.$$

The obvious guess for the supersymmetric completion of these interaction terms is simply to add the bosonization fields $\vec{\varphi}$ to the world-sheet energy-momentum tensors:

$$T_{++}^X \rightarrow T_{++}^{\text{tot}} = (\partial_+ \vec{X})^2 + (\partial_+ \vec{\varphi})^2,$$

and similarly for T_{--}^X . This will have a crucial effect on the normalization of the fixed scalar absorption rate.

3.4 Semi-classical description of absorption

In this section we will mainly discuss the solution of the radial differential equation one obtains for s -wave perturbations in the fixed scalar ν related to the volume of the internal T^4 in string metric (3.5). Let us start by restating some results of section 3.2. From (3.8), (3.14) and (3.15) one can read off the five-dimensional Einstein metric:

$$ds_5^2 = -(H_{\hat{Q}} H_{\hat{P}} H_{\hat{Q}_K})^{-2/3} h dt^2 + (H_{\hat{Q}} H_{\hat{P}} H_{\hat{Q}_K})^{1/3} (h^{-1} dr^2 + r^2 d\Omega_3^2). \quad (3.44)$$

To avoid the mixing between gravitational perturbations and the fixed scalar, we will restrict ourselves to the case $P = Q$, i.e. $r_1 = r_5 = R$, $P^2 = R^2(R^2 + r_0^2)$ (see (3.18)). The small fluctuation equation, (3.25), may be written as

$$\left[(hr^3 \partial_r)^2 + (r^2 + R^2)^2 (r^2 + r_K^2) \omega^2 - \frac{8r^4 R^4}{(r^2 + R^2)^2} h \left(1 + \frac{r_0^2}{R^2} \right) \right] \tilde{\nu} = 0, \quad (3.45)$$

where $h = 1 - \frac{r_0^2}{r^2}$. Since we work in the regime $r_0 \ll R$, we will neglect the last factor in the last term.

Several different radial coordinates are useful in different regions. The ones we will use most often are u and y defined by the relations

$$1 - \frac{r_0^2}{r^2} = e^{-r_0^2/u^2}, \quad y = \frac{R^2 r_K \omega}{2u^2}. \quad (3.46)$$

Note that $u \rightarrow 0$ and $y \rightarrow \infty$ at the horizon.

The most efficient tool for obtaining the absorption cross-section is the ratio of fluxes method used in [75]. In all the cases we will treat, the solution to (3.45) whose near-horizon limit represents purely infalling matter has the limiting forms

$$\tilde{\nu} \approx e^{iy} \quad (3.47)$$

near the horizon and

$$\tilde{\nu} \approx \alpha \frac{J_1(\omega r)}{\omega r} = \frac{\alpha}{2} \frac{H_1^{(1)}(\omega r) + H_1^{(2)}(\omega r)}{\omega r} \quad (3.48)$$

far from the black hole, where J_1 is a Bessel function.⁹ The term in (3.48) containing $H_1^{(2)}(\omega r)$ is the incoming wave. Once the constant α is known, one can compute the flux for the incoming wave and compare it to the flux for the infalling wave (3.47) to find the absorption probability. In the present instance, fluxes are purely radial:

$$J = \frac{1}{2i}(\tilde{\nu}^* d\tilde{\nu} - \text{c.c.}) = J_r dr . \quad (3.49)$$

Observing that the number of particles passing through a sphere S_r^3 at radius r in a time interval $[0, t]$ is

$$\int_{S_r^3 \times [0, t]} *J = 2\pi^2 h r^3 J_r t , \quad (3.50)$$

one concludes that the flux per unit solid angle is

$$\mathcal{F} = h r^3 J_r = \frac{1}{2i}(\tilde{\nu}^* h r^3 \partial_r \tilde{\nu} - \text{c.c.}) . \quad (3.51)$$

The absorption probability is

$$1 - |S_0|^2 = \frac{\mathcal{F}_h}{\mathcal{F}_\infty^{\text{incoming}}} = \frac{2\pi}{|\alpha|^2} R^2 r_K \omega^3 . \quad (3.52)$$

We will always be interested in cases where this probability is small. By the Optical Theorem, the absorption cross-section is

$$\sigma_{\text{abs}} = \frac{4\pi}{\omega^3} (1 - |S_0|^2) = \frac{8\pi^2}{|\alpha|^2} R^2 r_K . \quad (3.53)$$

Readers unfamiliar with the solution matching technology may be helped by the analogy to tunneling through a square potential barrier in one dimension. If particles come from the left side of the barrier, the wave function is to a good approximation a standing wave on the left side of the barrier, a decreasing exponential inside the barrier, and a purely right moving exponential on the right side of the barrier.

To obtain the familiar result $\sigma_{\text{abs}} = A_h$ for low-energy, ordinary scalars falling into an extremal black hole, it suffices to match the limiting value of (3.47) for small y directly to the limiting value of (3.48) for small r [71]. Due to non-extremality and to the presence of the potential term in (3.45), this naive matching scheme is invalid. A more refined approximate solution must be used, and a more physically interesting low-energy cross-section will be obtained.

We will now present approximate solutions to (3.45) in two regimes most easily characterized in D-brane terms: we shall first consider $T_R = 0$ with ω/T_L arbitrary; then we shall consider T_R much less than T_L but not zero, and allow ω/T_R to vary arbitrarily. In [96] a more general case was considered: ω/T_L and ω/T_R both arbitrary.

When $T_R = 0$, the black hole is extremal: $r_0 = 0$ and $r = u$. As usual, one proceeds by joining a near horizon solution **I** to a far solution **III** using an exact solution **II** to the $\omega = 0$ equation [8, 73, 70].

⁹In fact there can be phase shifts in the arguments of the exponential in (3.47) and the Bessel functions in (3.48), but they are immaterial for computing fluxes at leading order.

The dominant terms of (3.45) and the approximate solutions in the three regions are

$$\begin{aligned}
\text{I.} \quad & \left[\partial_y^2 + 1 - \frac{2\eta}{y} - \frac{2}{y^2} \right] \tilde{\nu}_{\text{I}} = 0 & \tilde{\nu}_{\text{I}} &= G_1(y) + iF_1(y) \\
\text{II.} \quad & \left[(r^3 \partial_r)^2 - 8 \frac{R^4}{H^2} \right] \tilde{\nu}_{\text{II}} = 0 & \tilde{\nu}_{\text{II}} &= \frac{C}{H(r)} + DH^2(r) \\
\text{III.} \quad & [(r^3 \partial_r)^2 + r^6 \omega^2] \tilde{\nu}_{\text{III}} = 0 & \tilde{\nu}_{\text{III}} &= \alpha \frac{J_1(\omega r)}{\omega r} + \beta \frac{N_1(\omega r)}{\omega r} ,
\end{aligned} \tag{3.54}$$

where C, D, α and β are constants, $H = 1 + R^2/r^2$, and F_1 and G_1 are Coulomb functions [97] whose charge parameter η is given by

$$\eta = -\frac{1}{4} \left(2\omega r_K + \frac{\omega R^2}{r_K} \right) = -\frac{1}{4\pi} \frac{\omega}{T_L} \left(1 + 2 \frac{r_K^2}{R^2} \right) . \tag{3.55}$$

In the last equality we have used the definition of T_L in (3.21). The quantity r_K^2/R^2 is small in the dilute gas approximation, and we will neglect it when comparing the final semi-classical cross-section with the D-brane answer.

By design, $\tilde{\nu}_{\text{I}} \rightarrow e^{iy}$ as $y \rightarrow \infty$ up to a phase shift in y . An approximate solution can be patched together from $\tilde{\nu}_{\text{I,II,III}}$ if one sets

$$C = \frac{\alpha}{2} = \frac{2}{3C_1(\eta)\omega r_K} , \quad D = 0 , \quad \beta = 0 , \tag{3.56}$$

where $C_1(\eta) = \frac{1}{3} e^{-\pi\eta/2} |\Gamma(2 + i\eta)|$ [97]. A slightly better matching can be obtained by allowing D and β to be nonzero, but the changes in the final solution do not affect the fluxes $\mathcal{F}_h$ and $\mathcal{F}_\infty$ (these changes are however crucial in determining S_0 by the old methods of [37], and give phase information on the scattered wave which the flux method does not). Having only $C \neq 0$ in region **II** is analogous to the fact that for right-moving particles incident on a square potential barrier, the wave function under the barrier can be taken as a purely falling exponential with no admixture of the rising exponential.

From (3.56) and (3.53) the cross-section is immediate:

$$\sigma_{\text{abs}} = \frac{\pi^2}{2} r_K R^2 (\omega r_K)^2 |3C_1(\eta)|^2 = \frac{1}{4} A_h (\omega r_K)^2 (1 + \eta^2) \frac{2\pi\eta}{e^{2\pi\eta} - 1} , \tag{3.57}$$

where A_h is the area of the horizon (given in (3.20)). Note that the derivation of (3.57) does not require the assumption that $r_K \ll R$.

To make the comparison with the D-brane approach, we can write (3.57) in the following suggestive form

$$\sigma_{\text{abs}} = \frac{\pi^2}{2} r_K R^2 (\omega r_K)^2 \frac{\frac{\omega}{2T_L}}{1 - e^{-\frac{\omega}{2T_L}}} \left(1 + \frac{\omega^2}{16\pi^2 T_L^2} \right) [1 + O(r_K^2/R^2)] . \tag{3.58}$$

In section 3.5 we will compute the same quantity using effective D-string method and will find agreement to the indicated order of accuracy. To obtain $O(r_K^2/R^2)$ corrections on the D-brane side one would have to go beyond the dilute gas approximation. An interesting special case where these corrections vanish is when $T_L = 0$, corresponding to $\eta \rightarrow -\infty$. In the brane description, this corresponds to 1-branes and 5-branes only with no condensate of open strings running between them: a pure quantum state with no thermal averaging. The limiting forms of the GR result (3.57) and of the D-brane absorption cross-section (3.2) to be derived in section 3.5 then agree exactly:

$$\sigma_{\text{abs}} = \left(\frac{\pi}{4} \right)^3 R^8 \omega^5 . \tag{3.59}$$

Now let us continue on to the second regime in which an approximate solution to the radial equation (3.45) is fairly straightforward to obtain: $\omega, T_R \ll T_L$ with ω/T_R arbitrary. A quantity which enters more naturally into the differential equations than ω/T_R is

$$B = \frac{R^2 r_K \omega}{r_0^2} = \frac{\omega}{\kappa} \tanh \sigma \approx \frac{\omega}{\kappa}, \quad \kappa \equiv 2\pi T_H, \quad (3.60)$$

where κ is the surface gravity at the horizon, and in the last step we used the fact that $\sigma \gg 1$ when $T_R \ll T_L$. In dropping terms from the exact equation (3.45) to obtain soluble forms in the three matching regions, it is essential to retain B as a quantity of $O(1)$; however, r_0/r_K and $\omega R^2/r_K$ can be regarded as small because $T_R \ll T_L$ and $\omega \ll T_L$. In regions **II** and **III**, the approximate equations turn out to be precisely the same as in (3.54), but in **I** one obtains a more complicated differential equation:

$$\left[\partial_y^2 + 1 - \frac{8}{B^2} \frac{e^{-2y/B}}{(1 - e^{-2y/B})^2} \right] \tilde{\nu}_{\mathbf{I}} = 0. \quad (3.61)$$

This equation can be cast in the form of a supersymmetric quantum mechanics eigenfunction problem. Define a rescaled variable $z = y/B$ and supercharge operators

$$\mathcal{Q} = -\partial_z + \coth z, \quad \mathcal{Q}^\dagger = \partial_z + \coth z. \quad (3.62)$$

Then (3.61) can be rewritten in the form

$$\mathcal{Q}\mathcal{Q}^\dagger \tilde{\nu}_{\mathbf{I}} = [-\partial_z^2 + 2\text{csch}^2 z + 1] \tilde{\nu}_{\mathbf{I}} = (1 + B^2) \tilde{\nu}_{\mathbf{I}}. \quad (3.63)$$

The eigenfunctions of the related Hamiltonian $\mathcal{Q}^\dagger \mathcal{Q} = -\partial_z^2 + 1$ are just exponentials, and from them one can read off the solutions to (3.63): the infalling solution is

$$\tilde{\nu}_{\mathbf{I}} = \frac{\mathcal{Q} e^{iBz}}{1 - iB} = \frac{\coth z - iB}{1 - iB} e^{iBz} = \frac{\coth \frac{y}{B} - iB}{1 - iB} e^{iy}. \quad (3.64)$$

The factor in the denominator is chosen so that $\tilde{\nu}_{\mathbf{I}} \approx e^{iy}$ for large y .

Performing the matching as usual, one obtains

$$\sigma_{\text{abs}} = \frac{1}{4} A_h (\omega r_K)^2 (1 + B^{-2}) = \frac{1}{4} A_h (\omega r_K)^2 \left(1 + \frac{4\pi^2 T_H^2}{\omega^2} \right). \quad (3.65)$$

In section 3.5 we will show that the effective string calculation gives the same result when $\omega, T_R \ll T_L$.

3.5 D-brane absorption cross-sections and emission rates

In this section we give a detailed calculation of the emission and absorption of the fixed scalar ν , using the interaction vertices computed in section 3.3. We recall that ν is related (see (3.5)) to the volume (measured in the string metric) of the compactification 4-torus orthogonal to the string. To study the leading coupling of ν , it is sufficient to retain the following two terms in the string effective action (cf. (3.37)):

$$I = \int d^2\sigma \left\{ -\frac{1}{2} (\partial_+ \vec{X} \cdot \partial_- \vec{X} + \partial_+ \vec{\varphi} \cdot \partial_- \vec{\varphi}) + \frac{1}{4T_{\text{eff}}} \left[(\partial_+ \vec{X})^2 + (\partial_+ \vec{\varphi})^2 \right] \left[(\partial_- \vec{X})^2 + (\partial_- \vec{\varphi})^2 \right] \nu(x) \right\} \quad (3.66)$$

where we have absorbed $\sqrt{T_{\text{eff}}}$ into the fields to make them properly normalized. From (3.7) we see that the scalar field with the proper bulk kinetic term is $2\nu/\kappa_5$. Consider the invariant amplitude

for processes mediated by the quartic interaction in (3.66). If p_1 and p_2 are the left-moving energies, while q_1 and q_2 are the right-moving ones, the matrix element among properly normalized states is

$$\frac{\sqrt{2}\kappa_5}{T_{\text{eff}}} \sqrt{\frac{q_1 q_2 p_1 p_2}{\omega}}. \quad (3.67)$$

The basic assumption of the D-brane approach to black hole physics is that the left-movers and right-movers can be treated as thermal ensembles [6, 51]. Strictly speaking, they are microcanonical ensembles, but for our purposes the canonical ensemble is good enough and we proceed as if we are dealing with a massless one-dimensional gas of left-movers of temperature T_L and right-movers with temperature T_R . The motivation for this assumption has been explained at length in several recent papers [6, 51, 75]. To compute the rate for the process $scalar \rightarrow L + L + R + R$ we have to square the normalized matrix element (3.67) and integrate it over the possible energies of the final state particles. Because of the presence of the thermal sea of left-movers and right-movers, we must insert Bose enhancement factors: for example, each left-mover in the final state picks up a factor of $1 + \rho_L(p_i) = -\rho_L(-p_i)$, where

$$\rho_L(p_i) = \frac{1}{e^{\frac{p_i}{T_L}} - 1} \quad (3.68)$$

is the Bose-Einstein distribution. If there were a left-mover of energy p_i in the initial state, it would pick up an enhancement factor $\rho_L(p_i)$. Similar factors attach to right-movers.

Conservation of energy and of momentum parallel to the effective string introduces the factor

$$(2\pi)^2 \delta(p_1 + p_2 + q_1 + q_2 - \omega) \delta(p_1 + p_2 - q_1 - q_2) = \frac{1}{2} (2\pi)^2 \delta(p_1 + p_2 - \frac{\omega}{2}) \delta(q_1 + q_2 - \frac{\omega}{2}) \quad (3.69)$$

into the integrals over p_1 , p_2 , q_1 , and q_2 . Putting everything together, we find that the rate for $scalar \rightarrow L + L + R + R$ is given by

$$\begin{aligned} \Gamma(1) &= \Gamma(scalar \rightarrow L + L + R + R) \\ &= \frac{36}{4} \frac{\kappa_5^2 L_{\text{eff}}}{4\pi^2 T_{\text{eff}}^2 \omega} \int_0^\infty dp_1 dp_2 \delta\left(p_1 + p_2 - \frac{\omega}{2}\right) \frac{p_1}{1 - e^{-\frac{p_1}{T_L}}} \frac{p_2}{1 - e^{-\frac{p_2}{T_L}}} \\ &\quad \times \int_0^\infty dq_1 dq_2 \delta\left(q_1 + q_2 - \frac{\omega}{2}\right) \frac{q_1}{1 - e^{-\frac{q_1}{T_R}}} \frac{q_2}{1 - e^{-\frac{q_2}{T_R}}}, \end{aligned} \quad (3.70)$$

where L_{eff} is the length of the effective string. The factor of $36 = 6^2$ arises from the presence of six species of left-movers (four bosons and two bosonized fermions) and six species of right-movers. We divide by $4 = 2^2$ because of particle identity: because the two left-movers in the final state are identical particles, the integral over p_1, p_2 counts every left-moving final state twice (similarly for the right-movers).

To write down the rates for the three other absorptions processes (that is, processes 2, 3, and 4 in eq. (3.1)), it is convenient to define the integrals

$$\begin{aligned} I_L(s_1, s_2) &= \int_0^\infty dp_1 dp_2 \delta\left(s_1 p_1 + s_2 p_2 + \frac{\omega}{2}\right) s_1 p_1 \rho_L(s_1 p_1) \cdot s_2 p_2 \rho_L(s_2 p_2) \\ I_R(s_1, s_2) &= \int_0^\infty dq_1 dq_2 \delta\left(s_1 q_1 + s_2 q_2 + \frac{\omega}{2}\right) s_1 q_1 \rho_L(s_1 q_1) \cdot s_2 q_2 \rho_L(s_2 q_2). \end{aligned} \quad (3.71)$$

The choices $s_i = 1$ and $s_i = -1$ correspond, respectively, to putting a particle in the initial or final

state. Then the total absorption rate, including all four competing processes of (3.1), is

$$\begin{aligned}
\Gamma_{\text{abs}}(\omega) &= \Gamma(1) + \Gamma(2) + \Gamma(3) + \Gamma(4) \\
&= 36 \frac{\kappa_5^2 L_{\text{eff}}}{4\pi^2 T_{\text{eff}}^2 \omega} \left[\frac{1}{4} I_L(-1, -1) I_R(-1, -1) + \frac{1}{2} I_L(-1, 1) I_R(-1, -1) \right. \\
&\quad \left. + \frac{1}{2} I_L(-1, -1) I_R(-1, 1) + I_L(-1, 1) I_R(-1, 1) \right] \\
&= \frac{9\kappa_5^2 L_{\text{eff}}}{4\pi^2 T_{\text{eff}}^2 \omega} \int_{-\infty}^{\infty} dp_1 dp_2 \delta\left(p_1 + p_2 - \frac{\omega}{2}\right) \frac{p_1}{1 - e^{-\frac{p_1}{T_L}}} \frac{p_2}{1 - e^{-\frac{p_2}{T_L}}} \\
&\quad \times \int_{-\infty}^{\infty} dq_1 dq_2 \delta\left(q_1 + q_2 - \frac{\omega}{2}\right) \frac{q_1}{1 - e^{-\frac{q_1}{T_R}}} \frac{q_2}{1 - e^{-\frac{q_2}{T_R}}} \\
&= \frac{\kappa_5^2 L_{\text{eff}}}{(32\pi)^2 T_{\text{eff}}^2} \frac{\omega}{\left(1 - e^{-\frac{\omega}{2T_L}}\right) \left(1 - e^{-\frac{\omega}{2T_R}}\right)} (\omega^2 + 16\pi^2 T_L^2) (\omega^2 + 16\pi^2 T_R^2) .
\end{aligned} \tag{3.72}$$

The fractional coefficients inside the square brackets on the second line of (3.72) are symmetry factors for the final states (the initial states are always simple enough so that their symmetry factors are unity). It is remarkable that although the individual processes 1–4 have rates which cannot be expressed in closed form, their sum is expressible in terms of integrals which can be performed analytically [98] because they run over all p_1 , p_2 , q_1 , and q_2 .

A similar calculation may be performed for the four emission processes, with the result

$$\Gamma_{\text{emit}}(\omega) = e^{-\frac{\omega}{T_H}} \Gamma_{\text{abs}}(\omega) = -\Gamma_{\text{abs}}(-\omega) , \tag{3.73}$$

where the Hawking temperature characterizing the distribution of the emitted scalars is related to T_R and T_L by (3.22). Our convention has been to compute $\Gamma_{\text{abs}}(\omega)$ assuming that the flux of the incoming scalar is unity. We have also suppressed the phase space factor $d^4k/(2\pi)^4$ for the outgoing scalar in computing $\Gamma_{\text{emit}}(\omega)$, and we have assumed that the outgoing scalar is emitted into the vacuum state, so that $\Gamma_{\text{emit}}(\omega)$ includes no Bose enhancement factors. These conventions were chosen because they lead to simple expressions for $\Gamma_{\text{emit}}(\omega)$ (3.73) and σ_{abs} below, but they must be borne carefully in mind when considering questions of detailed balance. Suppose we put the black hole in a thermal bath of scalars at temperature T_H . Then $\Gamma_{\text{abs}}(\omega)$ and $\Gamma_{\text{emit}}(\omega)$ pick up Bose enhancement factors for the scalars: those factors are, respectively, $1/(e^{\omega/T_H} - 1)$ and $1/(1 - e^{-\omega/T_H})$. Once these factors are included, the emission and absorption rates become equal by virtue of the first equality in (3.73). The fact that calculating $\Gamma_{\text{emit}}(\omega)$ in the same way that we calculated $\Gamma_{\text{abs}}(\omega)$ leads to (3.73) is a nontrivial check on detailed balance. This check is analogous to verifying that QED reproduces the Einstein A and B coefficients for the decay of the first excited state of hydrogen.

Because $\Gamma_{\text{abs}}(\omega)$ was computed assuming unit flux, one would naively guess that the absorption cross-section to be compared with a semi-classical calculation is $\sigma_{\text{abs}} = \Gamma_{\text{abs}}(\omega)$. (Now we are back to the conventions where $\Gamma_{\text{abs}}(\omega)$ and $\Gamma_{\text{emit}}(\omega)$ do not include Bose enhancement factors for the scalars). This is not quite right; instead,

$$\sigma_{\text{abs}}(\omega) = \Gamma_{\text{abs}}(\omega) - \Gamma_{\text{emit}}(\omega) = \Gamma_{\text{abs}}(\omega) + \Gamma_{\text{abs}}(-\omega) . \tag{3.74}$$

To see why (3.74) is right, we have to remember what we are doing in a semi-classical computation. We send in a classical wave in the field whose quanta are the scalars of interest, and we watch to see what fraction of it is sucked up by the black hole and what fraction is re-emitted. The quantum field theory analog is to send in a coherent state of scalars with large average particle number, so that the flux is almost fixed at its classical expectation value $\mathcal{F}$. The dominant processes are then absorption and stimulated emission. The Bose enhancement factors collapse to $\mathcal{F}$ for both absorption and emission, up to errors which are insignificant in the semi-classical limit. The net rate at which

particles are absorbed is then $\Gamma_{\text{abs}}(\omega)\mathcal{F} - \Gamma_{\text{emit}}(\omega)\mathcal{F}$. But this rate is $\sigma_{\text{abs}}\mathcal{F}$ by definition, whence (3.74). Note that the last expression in (3.74) is manifestly invariant under time-reversal, which takes $\omega \rightarrow -\omega$.

In order to obtain definite results for the absorption cross-section, we must supply values for the effective length L_{eff} of the string, as well as its effective tension T_{eff} . It is a by-now-familiar story that multiple D-strings bound to multiple five-branes behave like a single D-string multiply wound about the compactification direction [72]. In the case at hand it is well-understood that the effective string length is [72, 75]

$$\kappa_5^2 L_{\text{eff}} = 4\pi^3 r_1^2 r_5^2 . \quad (3.75)$$

With this substitution, the fixed scalar ν absorption cross-section becomes

$$\sigma_{\text{abs}}(\omega) = \frac{\pi r_1^2 r_5^2}{256 T_{\text{eff}}^2} \frac{\omega \left(e^{\frac{\omega}{T_H}} - 1 \right)}{\left(e^{\frac{\omega}{2T_L}} - 1 \right) \left(e^{\frac{\omega}{2T_R}} - 1 \right)} (\omega^2 + 16\pi^2 T_L^2)(\omega^2 + 16\pi^2 T_R^2) . \quad (3.76)$$

This is similar to, but not quite the same as, the absorption cross-section for the ordinary “unfixed” scalar calculated in [75].

The object of our exercise is to offer further evidence that the D-brane configuration *is* the corresponding black hole by showing that (3.76) is identical to the corresponding quantity calculated by standard classical GR methods. For technical reasons, the GR calculation in a general black hole background is quite difficult and the results we have been able to obtain (presented in section 3.4) are only valid in certain simplifying limits. The most important simplification is to take equal brane charges $r_1 = r_5 = R$.

First we consider the extremal limit, $T_R = 0$. Here (3.76) reduces to

$$\sigma_{\text{abs}}(\omega) = \frac{\pi^3 r_1^2 r_5^2 T_L^3}{8 T_{\text{eff}}^2} \omega^2 \frac{\frac{\omega}{2T_L}}{1 - e^{-\frac{\omega}{2T_L}}} \left(1 + \frac{\omega^2}{16\pi^2 T_L^2} \right) . \quad (3.77)$$

This is to be compared with the classical fixed scalar absorption cross-section in the extremal background (eq. (3.58)):

$$\sigma_{\text{abs}}(\omega) = \frac{\pi^2}{2} R^2 r_K^3 \omega^2 \frac{\frac{\omega}{2T_L}}{1 - e^{-\frac{\omega}{2T_L}}} \left(1 + \frac{\omega^2}{16\pi^2 T_L^2} \right) . \quad (3.78)$$

Using that in the extremal limit

$$T_L = \frac{r_K}{\pi r_1 r_5} ,$$

and remembering that we were only able to do the classical calculation for $r_1 = r_5 = R$, we see that the D-branes and GR match if we take the effective string tension to be

$$T_{\text{eff}} = \frac{1}{2\pi R^2} = \frac{1}{2\pi\alpha' g n_5} , \quad (3.79)$$

where we have restored the dependence on α' that we have been suppressing since (3.16). This value for T_{eff} is precisely equal to the tension of the “fractionated” D-string moving inside n_5 5-branes [72, 76]. This is a highly non-trivial independent check on the applicability of the effective string model to fixed scalars, and also on the idea of D-string “fractionation!”

Another interesting comparison to be made is for near-extremal black holes. For $\omega, T_R \ll T_L$ but with ratio of ω to T_R otherwise arbitrary, (3.76) becomes, using (3.79) for the tension,

$$\sigma_{\text{abs}}(\omega) = \frac{\pi^2}{2} R^2 r_K^3 (\omega^2 + 4\pi^2 T_H^2) . \quad (3.80)$$

This is in exact agreement with the absorption cross-section on non-extremal black holes (3.65) computed using general relativity.

For the fixed scalar the coupling to $(\partial X)^2$ is absent from the D-brane action, and the cross-section we found is the leading effect. For an ordinary decoupled scalar, such as the 6-d dilaton, both terms are present. So, the cross-section computed above should be part of the correction to the leading effect determined in [75]. This is an interesting topic for future investigation.

3.6 Conclusions

Let us try to recapitulate in a few words what it has taken many equations to state. The main thrust of the chapter has been to explore the behavior of the type of fixed scalar studied earlier in [64, 65, 69, 66, 67, 68], and most recently in [70] – but now in the context of five-dimensional black holes that can be modeled by bound states of D1-branes and D5-branes [5, 6, 72, 7, 8, 73, 99, 75]. For the most part we have focused on the fixed scalar ν which corresponds to the volume of the internal four-torus as measured by the string metric. Through an interesting interplay between semi-classical computations (where the basic theory is well known but analytically intractable in general) and D-brane computations (where the theory is less well known but very tractable), we have arrived at a general formula (3.76) for the cross-section for low-energy fixed scalars to be absorbed into the black hole.

The absorption cross-section (3.76) has a much richer and more interesting functional form than the simple ω^2 dependence found in [70]. Even in the simple limit $\omega, T_R \ll T_L$ in which comparison calculations between GR and D-branes were initially performed [7, 8, 73], the fixed scalar cross-section goes not as ω^2 but as $\omega^2 + \kappa^2$, where $\kappa = 2\pi T_H$ is the surface gravity at the horizon. While we have derived the expression (3.76) in full generality only in the D-brane picture, we have demonstrated that it agrees with semi-classical calculations of the cross-section in the two regimes: one regime reproduces this novel $\omega^2 + \kappa^2$ behavior, while the other deals with absorption into extremal black holes. Because the equations for the gravitational perturbations and fixed scalar perturbations couple unless two of the three charges, e.g., the 1-brane charge and the 5-brane charge, are equal to each other, our semi-classical computations are limited to the equal charge case (similar equal-charge assumption was used in $D = 4$ case in [70]). Modulo this limitation, we have confidence that a full greybody factor computation along the lines of [75] would reproduce the general result (3.76).

One of the reasons why the extension of the semi-classical calculations to unequal 1-brane and 5-brane charges (but with both still greater than the third charge, $P, Q \gg Q_K$, to remain in the dilute gas region) would be interesting, is that the D-brane computations involve one free parameter, the tension T_{eff} of the effective string, which can be read off from a comparison with a semi-classical calculation. The expectation, based on the work of [72, 76] and on the arguments given at the beginning of section 3.3, is that $T_{\text{eff}} = \frac{1}{2\pi\alpha' g n_5}$. Our work confirms this relation when the 1-brane and 5-brane charges are equal. What a semi-classical calculation with unequal charges should confirm is that T_{eff} is independent of the number of 1-branes.

Although our ultimate goal has been to demonstrate a new agreement between semi-classical GR and a perturbative treatment of the effective string, we have along the way studied interesting facets of both formalisms separately. On the D-brane side, we have been forced to go beyond the leading quadratic terms in the expansion of the DBI action and examine terms quartic in the derivatives of the string collective coordinate fields X^i . As we argued in section 3.3, the generic form of the quadratic terms is practically inevitable given the invariances of the problem. But the decoupling of the fixed scalar from quadratic terms and the precise form of its coupling to quartic terms is a signature of the DBI action. The agreement between the D-brane and GR cross-sections for fixed scalars is thus a more stringent test of the DBI action than the agreements obtained previously [8, 73, 74, 75] for ordinary scalars.

From the open string theory point of view, the $(\partial X)^2$ term in the D-string action (3.35),(3.37) originates upon dimensional reduction from the $F_{\mu\nu}^2$ term in the $D = 10$ Born-Infeld action, while the $(\partial X)^4$ terms correspond to the $F_{\mu\nu}^4$ -terms. It is amusing to note that the fixed scalars, which are coupled to the Maxwell terms of the closed string vector fields in the spacetime effective action (3.7), thus do not couple to the Maxwell term of the open string vector field in the effective D-string action, while exactly the opposite is true for the decoupled scalars. It is thus the $F_{\mu\nu}^4$ -terms in the DBI action (which are important also in some other contexts) that are effectively responsible for the leading contribution to the cross-section of fixed scalars.

At the relevant $(\partial X)^4$ order, the processes involving fermionic excitations of the effective string contribute in the same way as purely bosonic processes. Fortunately, the coupling of bosonic excitations to the fixed scalar field predicted by the DBI action is of a particularly simple form, $T_{++}^X T_{--}^X \nu(x)$, which admits an obvious generalization to include fermions: $T_{++}^{\text{tot}} T_{--}^{\text{tot}} \nu(x)$. Obtaining precise agreement with GR using this coupling and the normalization of T_{eff} as in [72, 76] may be viewed as determining a partial supersymmetrization of the effective string action via D-brane spectroscopy.

On the GR side, we have to some extent systematized the study of spherical black hole configurations, including spherically symmetric perturbations around the basic $D = 5$ black hole with three charges, by reducing the problem to an effective two-dimensional one. For time-independent configurations, this gives a straightforward derivation of the basic black hole solution. We were disappointed to find, however, that, despite relative simplicity of the effective two-dimensional theory compared to the full supergravity equations, it still leads to complicated coupled differential equations for time-dependent fluctuations around the static solution. So far, we have been able to extract simple equations from the intractable general case only when some pair of charges are equal. Then the background value of one fixed scalar becomes constant and its fluctuations decouple from the other fields, leading to a non-extremal five-dimensional generalization of the equation studied in [70]. Similar two-dimensional effective theory techniques with similar equal charge limitations were applied to the basic four-dimensional black hole with four charges. In this chapter, we have taken the four-dimensional calculations only far enough to see that fixed scalars whose background values become constant when three of the four charges are equal have an absorption cross-section with the characteristic $\omega^2 + \kappa^2$ dependence.

One final comment is that we have focused almost exclusively on absorption rather than Hawking emission. This is not because Hawking emission is any more difficult, but rather because agreement between the semi-classical Hawking calculation and the D-brane result is inevitable once a successful comparison of absorption cross-sections is made. To see this, one must only note that detailed balance between emission and absorption is built into the Hawking calculation and that it can be checked explicitly in the D-brane description. Once detailed balance is established in both descriptions, it obviously suffices to check that the absorption cross-section agrees between the two in order to be sure that emission rates must agree as well.

Chapter 4

Partial wave absorption by effective strings

4.1 Introduction

The D1-brane D5-brane bound state toroidally compactified down to five dimensions has proven to be one of the most fruitful string theoretic models of black holes. Since the original paper of [6], which proposed the model as a way to study black hole dynamics in a manifestly unitary string theoretic framework, and the subsequent work in [72] clarifying the means by which a single multiply wound effective string arises in a description of the low-energy dynamics, there have been many exciting papers relating properties of five-dimensional and four-dimensional black holes to the effective string. An explanation of the near-extremal entropy was given in [6, 38, 95], following the ideas originally laid out in [5]. Absorption cross-sections and the corresponding Hawking emission rates were worked out in [7, 8, 73, 99, 75, 87, 21, 96, 100, 101] yielding impressive agreement at low energies with the effective string model. Suggestions that the effective string model may have some flaws or limitations have arisen in the work of [102, 103, 83].

In a recent paper by Strominger and Maldacena [100] it was found from a general analysis of thermal two-point functions that the effective string seems capable of explaining the semi-classical absorption cross-sections for arbitrary partial waves of ordinary scalars. Subsequent work by Mathur [104] exhibited more detailed agreement between the effective string model and General Relativity for these processes. Ordinary scalars are scalars whose equation of motion in five-dimensions is $\square \phi = 0$. The canonical examples of ordinary scalars are the off-diagonal gravitons h_{ij} with both i and j lying within the D5-brane but perpendicular to the D1-brane. These are the scalars for whose s -wave cross-section full agreement between General Relativity and the effective string was first achieved in [8].

The results of [100, 104] overlap substantially with [22], on which the present chapter is based. The organization is as follows. Section 4.2 covers the semi-classical analysis of partial wave absorption and includes a derivation of a form of the Optical Theorem for the absorption of scalar particles which was quoted without proof in [10]. Section 4.3 presents the effective string description of the same processes, exhibiting along the way a simple method for performing all the phase space integrals encountered in [104]. In section 4.4, the limitation on the number of partial waves the effective string can couple to arising from statistics and locality is compared with the limitation imposed semi-classically by cosmic censorship. Section 4.5 summarizes the results and indicates directions for further work.

4.2 The semi-classical computation

The quantity that can be conveniently computed using the matching technique is the absorption probability. To convert this to an absorption cross-section, it is necessary to use properties of the partial wave expansion in four spatial dimensions and to invoke the Optical Theorem. The details of this connection were worked out independently in [104], but because the derivation given below applies for arbitrary dimensions, it seems worthwhile to present it in full. In all of what follows, $n = d - 1$ will denote the number of spatial dimensions.

The Optical Theorem for scattering of a scalar field off a spherically symmetric potential states that if the scattering wave-function has for large r the asymptotic form

$$\phi(\vec{r}) \sim e^{ikx} + f(\theta) \frac{e^{ikr}}{r^{(n-1)/2}} \quad (4.1)$$

(here $x = r \cos \theta$), then the total cross-section is

$$\sigma_{\text{total}} = -2 \left(\frac{2\pi}{k} \right)^{\frac{n-1}{2}} \Re \left(i^{\frac{n-1}{2}} f(0) \right) . \quad (4.2)$$

To find the partial wave expansion of (4.1), it is first necessary to make a Neumann expansion of the exponential function, which can be done using Gegenbauer polynomials [105]:

$$\begin{aligned} e^{ir \cos \theta} &= 2^{n/2-1} \Gamma(n/2-1) \sum_{\ell=0}^{\infty} i^{\ell} P_{\ell}(\cos \theta) (\ell + n/2 - 1) \frac{J_{\ell+n/2-1}(r)}{r^{n/2-1}} \\ P_{\ell}(\cos \theta) &= \sum_{m=0}^{\lfloor \ell/2 \rfloor} (-1)^m 2^{\ell-2m} \frac{\Gamma(\ell + n/2 - 1 - m)}{\Gamma(n/2 - 1) m! (\ell - 2m)!} \cos^{\ell-2m} \theta . \end{aligned} \quad (4.3)$$

The $P_{\ell}(\cos \theta)$ are just the Legendre polynomials when $n = 3$. For arbitrary n , they can be defined by the expansion

$$(1 - 2a \cos \theta + a^2)^{1-n/2} = \sum_{\ell=0}^{\infty} P_{\ell}(\cos \theta) a^{\ell} . \quad (4.4)$$

An alternate normalization proves more convenient:

$$\tilde{P}_{\ell}(\cos \theta) = \sqrt{\frac{2}{\pi}} 2^{n/2-1} \Gamma(n/2-1) (\ell + n/2 - 1) P_{\ell}(\cos \theta) . \quad (4.5)$$

Using asymptotic properties of Bessel functions, one can now write down the partial wave expansion of (4.1) as

$$e^{ikx} + f(\theta) \frac{e^{ikr}}{r^{(n-1)/2}} \sim \sum_{\ell=0}^{\infty} \frac{1}{2} \tilde{P}_{\ell}(\cos \theta) \frac{S_{\ell} e^{ikr} + (-1)^{\ell} i^{n-1} e^{-ikr}}{(ikr)^{(n-1)/2}} . \quad (4.6)$$

When $f(\theta) = 0$ identically, $S_{\ell} = 1$ for all ℓ .

The absorption cross-section for the ℓ^{th} partial wave can now be computed as the difference between the total ℓ -wave cross-section computed via the Optical Theorem,

$$\sigma_{\text{total}}^{\ell} = - \left(\frac{\sqrt{2\pi}}{k} \right)^{n-1} \tilde{P}_{\ell}(1) \Re(S_{\ell} - 1) , \quad (4.7)$$

and the ℓ -wave scattering cross-section,

$$\sigma_{\text{scattered}}^{\ell} = (\text{Vol } S^{n-2}) \frac{|S_{\ell} - 1|^2}{4k^{n-1}} \int_0^{\pi} d\theta \sin^{n-2} \theta \tilde{P}_{\ell}(\cos \theta)^2 . \quad (4.8)$$

The final result,

$$\sigma_{\text{abs}}^\ell = \frac{2^{n-2}\pi^{n/2-1}}{k^{n-1}} \Gamma(n/2-1)(\ell+n/2-1) \binom{\ell+n-3}{\ell} (1-|S_\ell|^2) , \quad (4.9)$$

relates the absorption cross-section σ_{abs}^ℓ to the absorption probability $1-|S_\ell|^2$. The results for $n=3$ and $n=4$ are

$$\begin{aligned} \sigma_{\text{abs}}^\ell &= \frac{\pi}{k^2} (2\ell+1) (1-|S_\ell|^2) && \text{in three spatial dimensions} \\ \sigma_{\text{abs}}^\ell &= \frac{4\pi}{k^3} (\ell+1)^2 (1-|S_\ell|^2) && \text{in four spatial dimensions.} \end{aligned} \quad (4.10)$$

With these results in hand, let us proceed to the semi-classical computation of the cross-section for an ordinary scalar ϕ in the ℓ^{th} partial wave to be absorbed into a black hole. It is hoped that this piece of “spectroscopic data” will be illuminating of the form of the effective string action.

An ordinary scalar is one whose equation of motion is just the Laplace equation following from the black hole metric, which in five dimensions is

$$\begin{aligned} ds^2 &= -F^{-2/3} h dt^2 + F^{1/3} (h^{-1} dr^2 + r^2 d\Omega_{S^3}^2) \\ F &= f_1 f_5 f_K = \left(1 + \frac{r_1^2}{r^2}\right) \left(1 + \frac{r_5^2}{r^2}\right) \left(1 + \frac{r_K^2}{r^2}\right) \\ h &= 1 - \frac{r_0^2}{r^2} . \end{aligned} \quad (4.11)$$

The mass, entropy, Hawking temperature, $U(1)$ charges, and characteristic radii are conveniently parametrized as

$$\begin{aligned} M &= \frac{\pi}{8} r_0^2 \sum_{i=1,5,K} \cosh 2\sigma_i & S &= \frac{\pi^2}{2} r_0^3 \prod_{i=1,5,K} \cosh \sigma_i & \beta_H &= 2\pi r_0 \prod_{i=1,5,K} \cosh \sigma_i \\ Q_i &= \frac{r_0^2}{2} \sinh 2\sigma_i & r_i &= r_0 \sinh \sigma_i \end{aligned} \quad (4.12)$$

in five-dimensional Planck units. Using a separation of variables,

$$\phi = e^{-i\omega t} P_\ell(\cos \theta) R(r) ,$$

one can extract from the Laplace equation $\square \phi = 0$ the radial equation

$$[(hr^3 \partial_r)^2 + r^6 F \omega^2 - r^4 h \ell(\ell+2)] R = 0 . \quad (4.13)$$

Because of the left-right symmetry of the effective string description for five-dimensional black holes, the absorption cross-section for the near-extremal case provides essentially no more information about the effective string than the extremal case does. In the interest of a simple presentation, we will therefore restrict the calculations in both this section and the next to the extremal case. The near-extremal generalizations of the results are summarized at the end of each section.

In the near horizon region (denoted **I** for consistency with the literature [37, 87, 21]), (4.13) for an extremal black hole can be approximated by a Coulomb equation [97] in the variable $y = (r_1 r_5 r_K \omega)/(2r^2)$, while in the far horizon region **III** it can be approximated as a Bessel equation:

$$\begin{aligned} \text{I.} \quad & \left[\partial_y^2 + 1 - \frac{2\eta}{y} - \frac{\ell(\ell+2)/4}{y^2} \right] R_{\text{I}} = 0 & R_{\text{I}} &= G_{\ell/2}(y) + i F_{\ell/2}(y) \\ \text{III.} \quad & [(r^3 \partial_r)^2 + r^6 \omega^2 - r^4 \ell(\ell+2)] R_{\text{III}} = 0 & R_{\text{III}} &= \alpha \frac{J_{\ell+1}(\omega r)}{\omega r} + \beta \frac{N_{\ell+1}(\omega r)}{\omega r} , \end{aligned} \quad (4.14)$$

where α and β are constants to be determined in the matching and

$$\eta = -\frac{1}{4} \sum_{i=1,5,K} \frac{r_1 r_5 r_K \omega}{r_i^2} \equiv -\frac{\omega}{4\pi T_L} \quad (4.15)$$

is the charge parameter of the Coulomb functions. The infalling solution $R_{\mathbf{I}}$ can be matched directly onto $R_{\mathbf{III}}$ without the aid of an intermediate region $\mathbf{II}$, with the result

$$\alpha = \frac{\ell!}{C_{\ell/2}(\eta)} \frac{2^{2\ell+1} \pi^\ell}{(A_h \omega^3)^{\ell/2}}, \quad \beta = 0. \quad (4.16)$$

The quantity

$$C_{\ell/2}(\eta) = \frac{2^{\ell/2} e^{-\pi\eta/2} \left| \Gamma\left(\frac{\ell}{2} + 1 + i\eta\right) \right|}{\Gamma(\ell + 2)} \quad (4.17)$$

enters into the series expansion of Coulomb functions.

A more accurate matching can be obtained with $\beta \neq 0$, but the level of accuracy embodied in (4.16) is sufficient for the flux ratio method [75, 87, 21]. In this method, the absorption probability is computed as the ratio of the infalling flux at the horizon to the flux in the incoming wave at infinity. The result is

$$1 - |S_\ell|^2 = \frac{1}{\pi} \frac{A_h \omega^3}{|\alpha|^2} = 4\pi \left(\frac{A_h \omega^3}{16\pi^2} \right)^{\ell+1} \frac{C_{\ell/2}^2(\eta)}{\ell!^2}. \quad (4.18)$$

Now the formula (4.10) comes into play to give the final result:

$$\begin{aligned} \sigma_{\text{abs}}^\ell &= A_h (\ell + 1)^2 \left(\frac{A_h \omega^3}{16\pi^2} \right)^\ell \frac{C_{\ell/2}^2(\eta)}{\ell!^2} \\ &= \frac{A_h}{\ell!^4} \left(\frac{A_h \omega^3}{8\pi^2} \right)^\ell e^{\frac{\omega}{4T_L}} \left| \Gamma\left(\frac{\ell}{2} + 1 - i\frac{\omega}{4\pi T_L}\right) \right|^2 \end{aligned} \quad (4.19)$$

The right hand side of (4.19) depends on r_1 , r_5 , and r_K only through A_h and T_L . Both quantities are symmetric in the three radii, and in fact admit U-duality invariant generalizations [106]. The near-extremal generalization of (4.19),

$$\sigma_{\text{abs}}^\ell = A_h \frac{(\omega r_0/2)^{2\ell}}{\ell!^4} \left| \frac{\Gamma\left(1 + \frac{\ell}{2} - \frac{i\omega}{4\pi T_L}\right) \Gamma\left(1 + \frac{\ell}{2} - \frac{i\omega}{4\pi T_R}\right)}{\Gamma\left(1 - i\frac{\omega}{2\pi T_H}\right)} \right|^2, \quad (4.20)$$

also treats the three charges on an equal footing. The temperatures T_L and T_R are given by [106]

$$\beta_{L,R} = 2\pi r_0 \left(\prod_i \cosh \sigma_i \mp \prod_i \sinh \sigma_i \right) \quad (4.21)$$

in the general non-extremal case.

4.3 The effective string analysis

Despite recent progress in formulating superspace actions for branes [33, 34, 107] and in generalizing the DBI action to nonabelian gauge theory (see [108] and references therein), a first-principles derivation of a complete action for the effective string, including all couplings to fields in the bulk of spacetime, has yet to be achieved. The goal of this section is to write down a reasonable form for the

part of the action responsible for coupling the effective string to higher partial waves of an ordinary scalar and see how the cross-sections it predicts compare with the semi-classical result (4.19).

Consider the off-diagonal graviton h_{ij} with i and j parallel to the D5-brane but perpendicular to the D1-brane. The lowest-order interaction of this field with excitations on the effective string can be read off from the DBI action [8]: in static gauge where $t = \tau$ and $x^5 = \sigma$,

$$V_{\text{int}} = -t_{\text{eff}} \int_0^{L_{\text{eff}}} d\sigma \, 2h_{ij}(\tau, \sigma, \vec{x}=0) \partial_+ X^i \partial_- X^j . \quad (4.22)$$

The convention in (4.22) and elsewhere is to sum over all $i \neq j$. The fields h_{ii} couple somewhat differently; an exploration of those couplings and their physical consequences was initiated in [21].

In [76], an analysis of the entropy and temperature of near-extremal 5-branes led to an effective string with $c_{\text{eff}} = 6$ and $T_{\text{eff}} = 1/(2\pi r_5^2)$. An extension of the methods used in [76] to the case $r_1 \sim r_5$ leads to [109]

$$T_{\text{eff}} = \frac{1}{2\pi(r_1^2 + r_5^2)} . \quad (4.23)$$

The natural assumption is that t_{eff} , by definition the tension that appears in front of the DBI action, is precisely T_{eff} . Strangely enough, all previous scattering calculations except the fixed scalar computation of [83] (whose implications regarding the effective string tension are unclear as yet) either do not depend on t_{eff} or require $r_1 = r_5$. Thus, purely from the point of view of scattering computations, t_{eff} seems ambiguous by a factor of the form $f(r_1/r_5)$ where $f(1) = 1$. One of the motivations for studying higher partial waves is to resolve this ambiguity. The result we will obtain is

$$t_{\text{eff}} = \frac{1}{2\pi r_1 r_5} . \quad (4.24)$$

Because of the evaluation of h_{ij} at $\vec{x} = 0$, (4.22) is a coupling to the s -wave of h_{ij} only. How might it be generalized to include the dominant couplings to arbitrary partial waves? To begin with, a coupling to the ℓ^{th} partial wave should include ℓ derivatives of h_{ij} since the wave-function vanishes like $|\vec{x}|^\ell$. It is the fermions on the effective string which carry the angular momentum [38, 95]: the left-moving and right-moving fermions transform in a fundamental of $SU(2)_L$ and $SU(2)_R$ respectively, where the $SO(4)$ of rotations in the four non-compact spatial dimensions is written as $SO(4) = SU(2)_L \times SU(2)_R$. Purely on group theory grounds, one thus expects the ℓ^{th} partial wave to couple to ℓ left-moving and ℓ right-moving fermions. The order of the absorption process in the string coupling can be read off from (4.19) as $g^{\ell+1}$ where g is the closed string coupling. Exactly two more open string vertex operators should be included in the interaction to make the disk diagram come out with this power of g . The natural candidate for the interaction is

$$\begin{aligned} V_{\text{int}} &= -t_{\text{eff}} \int_0^{L_{\text{eff}}} d\sigma \, 2h_{ij}(\tau, \sigma, x^m = \bar{\Psi} \gamma^m \Psi) \partial_+ X^i \partial_- X^j \\ &= -t_{\text{eff}} \int_0^{L_{\text{eff}}} d\sigma \, 2 \sum_{\ell=0}^{\infty} \frac{1}{\ell!} \left(\prod_{k=1}^{\ell} \bar{\Psi} \gamma^{m_k} \Psi \right) \partial_{m_1} \cdots \partial_{m_\ell} h_{ij}(\tau, \sigma, x^m = 0) \partial_+ X^i \partial_- X^j . \end{aligned} \quad (4.25)$$

Extra derivatives on the fermion fields are possible *a priori*, but power counting in ω for $\omega/T_L \ll 1$ shows that they must be absent if (4.19) is to be reproduced. The same general form of the coupling was deduced independently in [100] through a greybody factor analysis.

The outstanding fallacy of (4.25) is that the sum terminates at $\ell = 4$ because there are only four types of left-moving fermions and the same number of right-moving fermions. The situation is even worse when one factors in the restrictions from $SO(4)$ group theory. As we shall see after (4.29), only the $\ell = 0$ and $\ell = 1$ partial waves can be absorbed.

In (4.25), the γ^m are gamma matrices of $SO(4, 1)$:

$$\gamma^0 = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \quad \gamma^m = \begin{pmatrix} 0 & \tau_{\alpha\beta}^m \\ \tau^{m\dot{\alpha}\beta} & 0 \end{pmatrix} \quad (4.26)$$

where

$$\tau_{\alpha\dot{\alpha}}^m = (\mathbf{1}, i\sigma_1, i\sigma_2, i\sigma_3) \quad \tau^{m\dot{\alpha}\alpha} = \epsilon^{\dot{\alpha}\beta} \epsilon^{\alpha\beta} \tau_{\beta\dot{\beta}}^m = (\mathbf{1}, -i\sigma_1, -i\sigma_2, -i\sigma_3) , \quad (4.27)$$

σ_i being the usual Pauli matrices. We follow northwest contraction conventions for raising and lowering spinor indices, and we set $\epsilon_{01} = \epsilon^{01} = \epsilon_{\dot{0}\dot{1}} = \epsilon^{\dot{0}\dot{1}} = 1$. The four-component spinor Ψ decomposes into $SU(2)_L$ and $SU(2)_R$ fundamentals, which are left-movers and right-movers on the effective string, respectively. These complex fermions decompose further into real components of the 10-dimensional Majorana-Weyl spinor that one would expect to emerge most simply from a full string theory analysis:

$$\Psi = \begin{pmatrix} \Psi_{+\alpha} \\ \bar{\Psi}_{-\dot{\alpha}} \end{pmatrix} \quad (4.28)$$

$$\Psi_{\pm}^1 = \frac{\psi_{\pm}^1 + i\psi_{\pm}^2}{\sqrt{2}} \quad \Psi_{\pm}^2 = \frac{\psi_{\pm}^3 + i\psi_{\pm}^4}{\sqrt{2}} .$$

Note that complex conjugation raises or lowers a spinor index, rather than dotting or undotting it as in the case of $SO(3, 1)$.

For $\ell \geq 2$ the ℓ^{th} term in the interaction (4.25) makes subleading contributions to the absorption of lower partial waves because the expression $\partial_{m_1} \cdots \partial_{m_\ell} h_{ij}$ does not pick out a pure ℓ^{th} partial wave from a plane wave. The cure for this is to symmetrize $SU(2)_L$ and $SU(2)_R$ spinor indices:

$$V_{\text{int}} = -2t_{\text{eff}} \int_0^{L_{\text{eff}}} d\sigma \sum_{\ell=0}^4 \frac{i^\ell}{\ell!} \prod_{k=0}^{\ell} \left(\Psi_+^{\alpha_k} \Psi_{-\dot{\beta}_k} + \bar{\Psi}_+^{\alpha_k} \bar{\Psi}_{-\dot{\beta}_k} \right) \left(\tau_{(\alpha_1}^{\dot{\beta}_1} \cdots \tau_{\alpha_\ell)}^{\dot{\beta}_\ell} \right) \cdot \partial_{m_1} \cdots \partial_{m_\ell} h_{ij} \partial_+ X^i \partial_- X^j + \dots \quad (4.29)$$

Terms have been omitted in (4.29) which make subleading contributions. The product of fermion fields is antisymmetric in $\alpha_1 \dots \alpha_\ell$ and in $\dot{\beta}_1 \dots \dot{\beta}_\ell$. Hence all terms in (4.29) vanish except $\ell = 0$ and $\ell = 1$. The conclusion is that only the first two partial waves can be absorbed. This limitation is not merely a failing of the specific form (4.25); it is intrinsic to the approach of coupling partial wave to a product of fermion fields evaluated at a single point on the effective string without derivatives. To reiterate, the addition of derivatives introduces extra powers of the energy in the final cross-section which would cause disagreement with (4.19). Let us proceed with the analysis of $\ell = 1$ and consider possible extensions to $\ell \geq 2$ later.

There are many steps involved in passing from the interaction (4.29) to the cross-section for $\ell = 1$. To avoid losing factors it pays to be as explicit as possible. Let us begin with mode expansions of the fields appearing in (4.29). The forms of mode expansions are dictated by the kinetic terms in the action. In the present case, the kinetic terms are

$$S_{\text{bulk}} = \frac{1}{2\kappa_6^2} \int d^6x \frac{1}{4} (\partial_\mu h_{ij}) (\partial^\mu h^{ij}) + \dots \quad (4.30)$$

$$S_{\text{string}} = -2t_{\text{eff}} \int d^2\sigma \left[\partial_+ X_i \partial_- X^i + \psi_+^\Delta i \partial_- \psi_+^\Delta + \psi_-^\Delta i \partial_+ \psi_-^\Delta + \dots \right] ,$$

resulting in the mode expansions

$$\begin{aligned}
\psi_+^\Delta(\tau + \sigma) &= \sum_{k^5 \in \frac{2\pi}{L_{\text{eff}}}(\mathbf{Z}^- - 1/2)} \frac{1}{\sqrt{2L_{\text{eff}}t_{\text{eff}}}} (b_k^\Delta e^{ik \cdot \sigma} + \text{h.c.}) \\
\psi_-^\Delta(\tau - \sigma) &= \sum_{k^5 \in \frac{2\pi}{L_{\text{eff}}}(\mathbf{Z}^+ + 1/2)} \frac{1}{\sqrt{2L_{\text{eff}}t_{\text{eff}}}} (b_k^\Delta e^{ik \cdot \sigma} + \text{h.c.}) \\
X^i(\sigma, \tau) &= \sum_{k^5 \in \frac{2\pi}{L_{\text{eff}}}\mathbf{Z}} \frac{1}{\sqrt{2L_{\text{eff}}t_{\text{eff}}k^0}} (a_k^i e^{ik \cdot \sigma} + \text{h.c.}) \\
h^{ij}(x^\mu) &= \sum_{k^5, \vec{k}} \sqrt{\frac{2\kappa_6^2}{2VL_5k^0}} (g_k^{ij} e^{ik \cdot x} + \text{h.c.}) .
\end{aligned} \tag{4.31}$$

The sum over $k^5, \vec{k}$ has $k^5 \in \frac{2\pi}{L_5}\mathbf{Z}$ and $k^m \in \frac{2\pi}{\sqrt[4]{V}}\mathbf{Z}$ for $m = 1, 2, 3, 4$. V is the volume of a large box in which we imagine enclosing the four uncompactified spatial dimensions. The indices Δ and $\dot{\Delta}$ run from 1 to 4, and since ψ_+^Δ and ψ_-^Δ are real, conjugation does not change the position of the indices. Typographical convenience will dictate the position of Δ and $\dot{\Delta}$.

It is important that h^{ij} is moded differently in the x^5 direction from the effective string excitations: the minimal quantum of Kaluza-Klein charge for an excitation on the effective string is $1/(n_1 n_5)$ of the minimal quantum for a particle in the bulk [94, 72]. In (4.31) we have not been careful about zero modes because it is the oscillator states which are important for the absorption processes. The factors in (4.31) were chosen to make the commutation relations simple:

$$\begin{aligned}
\{b_k^\Delta, b_q^{\Gamma^\dagger}\} &= \delta_{k^5 - q^5} \delta^{\Delta\Gamma} \\
\{b_k^{\dot{\Delta}}, b_q^{\dot{\Gamma}^\dagger}\} &= \delta_{k^5 - q^5} \delta^{\dot{\Delta}\dot{\Gamma}} \\
[a_k^i, a_q^{j^\dagger}] &= \delta_{k^5 - q^5} \delta^{ij} \\
[g_k^{ij}, g_q^{fh^\dagger}] &= \delta_{\vec{k} - \vec{q}} \delta^{if} \delta^{jh} \quad \text{if } i < j \text{ and } f < h .
\end{aligned} \tag{4.32}$$

The goal now is to compute the amplitude $\langle \tilde{f} | V_{\text{int}} | \tilde{i} \rangle$ for an absorption process where a scalar in the $\ell = 1$ partial wave turns into two bosons and two fermions on the effective string. The tildes on $|\tilde{i}\rangle$ and $|\tilde{f}\rangle$ are meant to indicate that these state vectors are not the real initial and final states: they include only the particles that participate in the interaction and not the whole thermal sea of left-movers that give the effective string its Kaluza-Klein charge. Restoring the thermal sea is an easy exercise which will be postponed until (4.39).

Two other slight simplifications will be made to ease the notational burden. First, indices can be dropped on all the X^i fields, but then one must include an extra factor of 2 in the rate, as shown in (4.35). The 2 accounts for the fact that h_{ij} can turn into a left-moving X^i and a right-moving X^j or a left-moving X^j and a right-moving X^i . The second simplification is to consider only

$$\begin{aligned}
|\tilde{i}\rangle &= g_k^\dagger |0\rangle \\
|\tilde{f}\rangle &= a_{p_b}^\dagger a_{q_b}^\dagger b_{p_f}^{\Delta^\dagger} b_{q_f}^{\dot{\Delta}^\dagger} |0\rangle ,
\end{aligned} \tag{4.33}$$

which is to say we put all the particles on the effective string into the final state and none into the initial state. A simple way to account for all the crossed processes which also contribute to absorption will be discussed after equation (4.42). In (4.33) and below, k refers to the momentum of the bulk scalar, p refers to the momentum of a left-mover on the effective string, and q refers to the momentum of a right-mover.

The desired matrix element can now be read off from (4.29) and (4.32) as

$$\langle \tilde{f} | V_{\text{int}} | \tilde{i} \rangle = C_{\Delta}^{\dot{\Delta}} \frac{k_1}{2L_{\text{eff}} t_{\text{eff}}} \kappa_5 \sqrt{\frac{p_b^0 q_b^0}{V k^0}} \delta_{k^5 - p_b^5 - p_f^5 - q_b^5 - q_f^5} , \quad (4.34)$$

where $C_{\Delta}^{\dot{\Delta}}$ is the 4×4 matrix $\text{diag}\{1, -1, 1, -1\}$. To extract the rate is necessary to use a generalization of Fermi's Golden Rule that includes Bose enhancement factors and Fermi suppression factors:

$$\begin{aligned} \Gamma &= 2 \sum_{|\tilde{f}\rangle} (\rho_L^X(p_b^0) + 1)(\rho_R^X(q_b^0) + 1)(1 - \rho_L^\psi(p_f^0))(1 - \rho_L^\psi(q_f^0)) \\ &\quad \cdot \left| \langle \tilde{f} | V_{\text{int}} | \tilde{i} \rangle \right|^2 2\pi \delta(k^0 - p_b^0 - p_f^0 - q_b^0 - q_f^0) \\ &= 2 \sum_{\Delta, \dot{\Delta}} |C_{\Delta}^{\dot{\Delta}}|^2 \cdot \sum_{\text{modes}} (\rho_L^X(p_b^0) + 1)(\rho_R^X(q_b^0) + 1)(1 - \rho_L^\psi(p_f^0))(1 - \rho_L^\psi(q_f^0)) \\ &\quad \cdot \frac{k_1^2 \kappa_5^2}{V k^0 (2L_{\text{eff}} t_{\text{eff}})^2} p_b^0 q_b^0 \delta_{k^5 - p_b^5 - p_f^5 - q_b^5 - q_f^5} 2\pi \delta(k^0 - p_b^0 - p_f^0 - q_b^0 - q_f^0) \end{aligned} \quad (4.35)$$

where

$$\rho_L^X(p_b^0) = \frac{1}{e^{p_b^0/T_L} - 1} , \quad \rho_L^\psi(p_i^0) = \frac{1}{e^{p_i^0/T_L} + 1} \quad (4.36)$$

and similarly for the right-moving thermal occupation factors. Again, the explicit factor of 2 in the first line of (4.35) is present to account for the two distinct choices, $i+ j-$ or $i- j+$, for polarizing the bosonic fields.

The vanishing of all cross-sections beyond $\ell = 1$ is an egregious failing of the most naive effective string model. The simplest fix would be to allow an incoming scalar to couple to a product of fermion fields evaluated at a single point on the spatial S^1 which the effective string wraps, but not necessarily at a single point in the effective string coordinates σ . In terms of the (4,4) SCFT from which the effective string emerges as a particular twisted sector, this more general coupling seems very natural because it still involves only a local operator constructed from a product of the $4n_1 n_5$ species of fermions in the SCFT.

The present treatment extends easily to cover this more general interaction. Let D_k and $\dot{D}_k$ be $4n_1 n_5$ -valued indices for the left- and right-moving fermion fields, respectively. Consider the final state

$$|\tilde{f}\rangle = a_{p_b}^\dagger a_{q_b}^\dagger \left(b_{p_1}^{D_1^\dagger} b_{q_1}^{\dot{D}_1^\dagger} \dots b_{p_\ell}^{D_\ell^\dagger} b_{q_\ell}^{\dot{D}_\ell^\dagger} \right) |0\rangle . \quad (4.37)$$

The matrix element is now of the form

$$\langle \tilde{f} | V_{\text{int}} | \tilde{i} \rangle = C_{D_1 \dots D_\ell}^{\dot{D}_1 \dots \dot{D}_\ell} \frac{k_1^\ell}{(2L_{\text{eff}} t_{\text{eff}})^\ell} \kappa_5 \sqrt{\frac{p_b^0 q_b^0}{V k^0}} \delta_{k^5 - p_b^5 - \Sigma p_i^5 - q_b^5 - \Sigma q_i^5} . \quad (4.38)$$

The coefficient tensor $C_{D_1 \dots D_\ell}^{\dot{D}_1 \dots \dot{D}_\ell}$ is antisymmetric in $D_1 \dots D_\ell$ and in $\dot{D}_1 \dots \dot{D}_\ell$. It encodes the $SO(4)$ group theory factors isolating the ℓ^{th} partial wave as well as restrictions on the possible final states

arising from D1-brane and D5-brane Chan-Paton factors. The rate is

$$\begin{aligned}
\Gamma &= 2 \sum_{|\tilde{f}\rangle} (\rho_L^X(p_b^0) + 1)(\rho_R^X(q_b^0) + 1) \prod_{i=1}^{\ell} \left[(1 - \rho_L^{\psi}(p_i^0))(1 - \rho_L^{\psi}(q_i^0)) \right] \\
&\quad \cdot \left| \langle \tilde{f} | V_{\text{int}} | \tilde{i} \rangle \right|^2 2\pi \delta(k^0 - p_b^0 - \sum p_i^0 - q_b^0 - \sum q_i^0) \\
&= \frac{2}{\ell!^2} \sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_{\ell}}^{\dot{D}_1 \dots \dot{D}_{\ell}}|^2 \cdot \sum_{\text{modes}} (\rho_L^X(p_b^0) + 1)(\rho_R^X(q_b^0) + 1) \prod_{i=1}^{\ell} \left[(1 - \rho_L^{\psi}(p_i^0))(1 - \rho_L^{\psi}(q_i^0)) \right] \\
&\quad \cdot \frac{k_1^{2\ell} \kappa_5^2}{V k^0 (2L_{\text{eff}} t_{\text{eff}})^{2\ell}} p_b^0 q_b^0 \delta_{k^5 - p_b^5 - \sum p_i^5 - q_b^5 - \sum q_i^5} 2\pi \delta(k^0 - p_b^0 - \sum p_i^0 - q_b^0 - \sum q_i^0) .
\end{aligned} \tag{4.39}$$

The $1/\ell!^2$ in the last expression arises because the sums over D_k , $\dot{D}_k$, p_k , and q_k are unrestricted, and there are $\ell!^2$ different permutations of a given set of values for these quantities which yield the same final state $|\tilde{f}\rangle$.

When the energy of the incoming scalar is much greater than the gap, a continuum approximation can be made in (4.39):

$$\sum_p \rightarrow \int dp \frac{L_{\text{eff}}}{2\pi} , \quad \delta_p \rightarrow \frac{2\pi}{L_{\text{eff}}} \delta(p) . \tag{4.40}$$

For the sake of simplicity, only massless particle absorption will be considered. In that case the flux associated with the state $g_k^{\dagger}|0\rangle$ is $\mathcal{F} = 1/V$. The absorption cross-section is

$$\begin{aligned}
\sigma_{\text{abs}} &= V \Gamma(\text{scalar} \rightarrow b_L + b_R + \ell f_L + \ell f_R) + \text{crossed processes} \\
&= \frac{\sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_{\ell}}^{\dot{D}_1 \dots \dot{D}_{\ell}}|^2}{\ell!^2 4^{\ell}} \frac{\kappa_5^2 L_{\text{eff}}}{(2\pi t_{\text{eff}})^{2\ell}} \omega^{2\ell-1} I_L I_R
\end{aligned} \tag{4.41}$$

where

$$I_L = \int_{-\infty}^{\infty} dp_b^0 \prod_{i=1}^{\ell} dp_i^0 \delta\left(\frac{\omega}{2} - p_b^0 - \sum_{i=1}^{\ell} p_i^0\right) \frac{p_b^0}{1 - e^{-p_b^0/T_L}} \prod_{i=1}^{\ell} \frac{1}{1 + e^{-p_i^0/T_L}} \tag{4.42}$$

and similarly for I_R . $\Gamma(\text{scalar} \rightarrow b_L + b_R + \ell f_L + \ell f_R)$ is what was computed in (4.39). Arbitrary crossings of the basic process $\text{scalar} \rightarrow b_L + b_R + \ell f_L + \ell f_R$ and their time-reversals also contribute to the net absorption rate from which σ_{abs} is computed. However, the simple trick of extending the integrals in (4.42) over the entire real line can be used to keep track of all of them. A demonstration of this with careful attention paid to symmetry factors can be found in chapter 3 for the special case where only two left-moving bosons and two right-moving bosons are involved. For $\ell = 0$ the dependence on t_{eff} disappears in (4.41), as was noted in chapter 3.

The integral (4.42) is a convolution of $\ell + 1$ simple functions and so can be done most directly by transforming to Fourier space, where convolutions become products. Three integrals which are useful for doing the Fourier transforms are

$$\begin{aligned}
\int_{-\infty}^{\infty} dp e^{ixp} \frac{p}{2 \sinh \frac{p}{2T_L}} &= (\pi T_L)^2 \text{sech}^2(\pi T_L x) \\
\int_{-\infty}^{\infty} dp e^{ixp} \frac{1}{2 \cosh \frac{p}{2T_L}} &= (\pi T_L) \text{sech}(\pi T_L x) \\
\int_{-\infty}^{\infty} dx e^{-ixp} (\pi T_L)^{\ell+2} \text{sech}^{\ell+2}(\pi T_L x) &= (2\pi T_L)^{\ell+1} \frac{\left| \Gamma\left(\frac{\ell}{2} + 1 - i \frac{p}{2\pi T_L}\right) \right|^2}{(\ell+1)!} .
\end{aligned} \tag{4.43}$$

The first two integrals are Fourier inversions of the third in the special cases $\ell = 0$ and -1 . Now the computation is straightforward:

$$\begin{aligned}
I_L &= e^{\frac{\omega}{4T_L}} \int_{-\infty}^{\infty} dp_b^0 \prod_{i=1}^{\ell} dp_i^0 \delta \left(\frac{\omega}{2} - p_b^0 - \sum_{i=1}^{\ell} p_i^0 \right) \frac{p_b^0}{2 \sinh \frac{p_b^0}{2T_L}} \prod_{i=1}^{\ell} \frac{1}{2 \cosh \frac{p_i^0}{2T_L}} \\
&= \frac{e^{\frac{\omega}{4T_L}}}{2\pi} \int_{-\infty}^{\infty} dx e^{-ix\omega/2} (\pi T_L)^{\ell+2} \operatorname{sech}^{\ell+2}(\pi T_L x) \\
&= \frac{(\ell+1)!}{\pi} (\pi T_L)^{\ell+1} C_{\ell/2}^2(\eta) .
\end{aligned} \tag{4.44}$$

The last step uses (4.15) and (4.17). I_R can be computed similarly, but since $T_R = 0$ by assumption, the result is much simpler:

$$I_R = \frac{(\omega/2)^{\ell+1}}{(\ell+1)!} . \tag{4.45}$$

Now the absorption cross-section can be given in closed form:

$$\begin{aligned}
\sigma_{\text{abs}}^{\ell} &= \frac{\sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_{\ell}}^{\dot{D}_1 \dots \dot{D}_{\ell}}|^2}{\ell!^2 4^{\ell}} \frac{\kappa_5^2 L_{\text{eff}}}{(2\pi t_{\text{eff}})^{2\ell}} \frac{\omega^{3\ell}}{\pi} \left(\frac{\pi T_L}{2} \right)^{\ell+1} C_{\ell/2}^2(\eta) \\
&= A_h \sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_{\ell}}^{\dot{D}_1 \dots \dot{D}_{\ell}}|^2 \left(\frac{A_h \omega^3}{16\pi^2} \right)^{\ell} \frac{C_{\ell/2}^2(\eta)}{\ell!^2} .
\end{aligned} \tag{4.46}$$

In the second equality two key relations have been used:

$$T_L = \frac{r_K}{\pi r_1 r_5} \quad \frac{\kappa_5^2 L_{\text{eff}} T_L}{2} = A_h . \tag{4.47}$$

Both are valid when $r_K \ll r_1, r_5$. The first can be derived by setting the effective string entropy equal to the Bekenstein-Hawking entropy. The second is a limiting case of (4.21). In addition, the tension has at last been fixed:

$$t_{\text{eff}} = \frac{1}{2\pi r_1 r_5} . \tag{4.48}$$

Because the cross-section (4.19) depends on n_1 and n_5 only through the product $n_1 n_5$ in the dilute gas regime, and because the same is true of the quantities L_{eff} , T_L , and A_h when $r_0 = 0$ and $r_K \ll r_1, r_5$, the choice $t_{\text{eff}} \sim 1/\sqrt{n_1 n_5}$ seems inevitable.

Precise agreement between General Relativity and the effective string now depends only on the relation

$$\sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_{\ell}}^{\dot{D}_1 \dots \dot{D}_{\ell}}|^2 = (\ell+1)^2 . \tag{4.49}$$

Agreement in the case $\ell = 0$ is trivial. For $\ell = 1$, the original treatment in terms of free fermions on the effective string is adequate: one can easily trace through the computations and verify that D_1 , $\dot{D}_1$, and $C_{D_1}^{\dot{D}_1}$ can be replaced in every equation by Δ , $\dot{\Delta}$, and $C_{\Delta}^{\dot{\Delta}}$. Any numerical discrepancy could have been fixed by introducing a multiplicative constant in the relation $x^m = \bar{\Psi} \gamma^m \Psi$; however to see perfect agreement without such artifice is pleasing and also rather suggestive of the form one expects for a gauge-fixed kappa symmetric action.

The real test is $\ell \geq 2$. Here it seems essential to depart from the simplistic effective string picture and return to a more fundamental description of the D1-D5 bound state in order to compute the coefficients $C_{D_1 \dots D_{\ell}}^{\dot{D}_1 \dots \dot{D}_{\ell}}$.

Finally, it is worth noting that if agreement can be established for extremal absorption, agreement for the near-extremal case follows automatically. In the effective string computation for near-extremal absorption one must subtract off the stimulated emission contribution as described in chapter 3 in order to respect detailed balance and time reversal invariance. Modulo this subtraction, the result can be read off from (4.41) and (4.44):

$$\begin{aligned}\sigma_{\text{abs}}^\ell &= \frac{\sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_\ell}^{\dot{D}_1 \dots \dot{D}_\ell}|^2}{\ell!^2 (\ell+1)!^2} \frac{\kappa_5^2 L_{\text{eff}} T_L T_R}{T_H} \left(\frac{\omega \sqrt{T_L T_R}}{2t_{\text{eff}}} \right)^{2\ell} \\ &\quad \cdot \left| \frac{\Gamma\left(1 + \frac{\ell}{2} - \frac{i\omega}{4\pi T_L}\right) \Gamma\left(1 + \frac{\ell}{2} - \frac{i\omega}{4\pi T_R}\right)}{\Gamma\left(1 - i\frac{\omega}{2\pi T_H}\right)} \right|^2 \\ &= \frac{\sum_{D_k, \dot{D}_k} |C_{D_1 \dots D_\ell}^{\dot{D}_1 \dots \dot{D}_\ell}|^2}{\ell!^2 (\ell+1)!^2} A_h \left(\frac{\omega r_0}{2} \right)^{2\ell} \left| \frac{\Gamma\left(1 + \frac{\ell}{2} - \frac{i\omega}{4\pi T_L}\right) \Gamma\left(1 + \frac{\ell}{2} - \frac{i\omega}{4\pi T_R}\right)}{\Gamma\left(1 - i\frac{\omega}{2\pi T_H}\right)} \right|^2.\end{aligned}\tag{4.50}$$

The second line relies on a modified version of (4.47) applicable to the near-extremal case with $r_0, r_K \ll r_1, r_5$:

$$\sqrt{T_L T_R} = \frac{r_0}{2\pi r_1 r_5} \quad \frac{\kappa_5^2 L_{\text{eff}} T_L T_R}{T_H} = A_h.\tag{4.51}$$

The same tension (4.48) and the same relation (4.49) establish agreement between (4.50) and (4.20).

It has been suggested [106] that the effective string picture can be used to describe in a U-duality invariant fashion black holes with arbitrary charges, possibly even far from extremality. The expression on the first line of (4.50) depends only on T_L, T_R, T_H , and L_{eff} —all quantities that have meaning to the effective string considered in the abstract, independent of the microscopic D1-D5-brane model. It cries out to be reconciled with (4.20) for arbitrary values of r_0, Q_1, Q_5 , and Q_K . But the treatment of absorption given in this section relies on the dilute gas approximation and thus is not general enough to be matched in any meaningful way to General Relativity when $Q_K \not\ll Q_1, Q_5$.

4.4 Limitations on partial wave absorption

Consider the more general couplings described in the paragraph preceding (4.37): local on spatial S^1 wrapped by the effective string but not on the effective string itself.¹ Although the details of the $SO(4)$ group theory and Chan-Paton factors have yet to be worked out fully in the context of the (4,4) SCFT description of the D1-D5-brane bound state, it seems clear that a coupling of the ℓ^{th} partial wave to an operator built out of any combination of the $4n_1 n_5$ fermionic fields raises the maximum value of ℓ which the effective string can absorb from 1 to some number on the order $n_1 n_5$. One might suppose that by putting derivatives on some of the fermion fields, the problem can be avoided altogether. But such derivatives raise the dimension of the operator and hence suppress the cross-section by more powers of ω than are present in the semi-classical result. To sum up, the assumption of locality prevents the effective string from coupling to partial waves above a certain maximum ℓ with the strength required to match General Relativity. The situation does not seem as satisfactory as for the D3-brane, where couplings to all partial waves exist with appropriate dimensions to reproduce semi-classical cross-sections [9] (normalizations however are problematic, as we shall see in chapter 6).

There is a reason, however, why one might expect not to observe agreement between the effective string and General Relativity at high values of ℓ . If the black hole absorbs some very high partial wave, it winds up with a large angular momentum, so the geometry before and after is appreciably different. Back reaction is not included in the General Relativity calculations of section 4.2. In fact

¹The ideas in this section originate largely in discussions with C. Callan, L. Thorlacius, and J. Maldacena.

the only back reaction calculation to date for the black hole under consideration [110] is restricted to $\ell = 0$. But on the grounds of cosmic censorship one would expect that absorption processes which drive the black hole past extremality are forbidden even semi-classically. From an adaptation of the work of [38, 95] one can read off the corresponding bound on ℓ as $\ell \lesssim \sqrt{n_K n_1 n_5}$.

We have two different bounds on ℓ indicating the maximum partial wave that the effective string should be capable of absorbing:

$$\begin{aligned} \ell &\lesssim \ell_{\max}^L \equiv n_1 n_5 && \text{from locality and statistics} \\ \ell &\lesssim \ell_{\max}^C \equiv \sqrt{n_K n_1 n_5} && \text{from cosmic censorship.} \end{aligned} \quad (4.52)$$

Now we would like to inquire which is the more restrictive. Using the standard relations (see for example [99])

$$n_1 n_5 = \frac{4\pi^3 r_1^2 r_5^2}{\kappa_5^2 L_5} \quad n_K = \frac{\pi L_5 r_K^2}{\kappa_5^2} \quad (4.53)$$

and the formula (4.47) for T_L , one can show that $\ell_{\max}^C / \ell_{\max}^L = L_5 T_L / 2$.

The validity of any comparison between General Relativity and the effective string model as treated in section 4.3 relies on being in the dilute gas regime [6] and at low energies [111]:

$$r_K \ll r_1, r_5 \ll 1/\omega. \quad (4.54)$$

These inequalities still do not determine whether $\ell_{\max}^L$ is larger or smaller than $\ell_{\max}^C$. But if it is agreed to examine only fat black holes [72], which is to say if one assumes

$$L_5 \ll \sqrt{r_1 r_5}, \quad (4.55)$$

then it is easy to obtain the inequality

$$\ell_{\max}^L \gg \ell_{\max}^C \quad (4.56)$$

by combining the dilute gas inequality in (4.54) with (4.55). Now, (4.56) is a hopeful state of affairs for the effective string model, because it indicates that making couplings local only on S^1 in principle enables the effective string to absorb all the partial waves for which reasonable comparisons can be made with General Relativity. It is perhaps not the ideal state of affairs: one might have hoped that the bounds $\ell_{\max}^L$ and $\ell_{\max}^C$ would coincide, indicating that the effective string knew about cosmic censorship. The results of [100] suggest that a more careful treatment of these issues using techniques of conformal field theory would result in a translation of cosmic censorship into unitarity of the effective string description. Such a treatment would need to address the problem that if one moves deep into the black string region of parameter space by making L_5 large, one can obtain $\ell_{\max}^L \ll \ell_{\max}^C$. The perturbative D-brane region is in fact closer to the black string region than the fat black hole region, so it would be surprising to find such a disaster for comparisons in the black string region when agreement seems possible for fat black holes.

4.5 Conclusion

In this chapter we have shown that the leading order coupling of the effective string to an ordinary scalar correctly predicts the cross-sections for the $\ell = 1$ partial wave. Due to the Grassmannian character of the fermionic fields which carry the angular momentum, it is impossible for the simplest effective string model (a single long string with $c_{\text{eff}} = 6$) to couple properly to $\ell > 1$ partial waves through an operator local on the string. Generalizing the model to include what one might intuitively regard as multi-strand interactions of the effective string postpones this difficulty to $\ell \gtrsim n_1 n_5$, a higher bound on ℓ for fat black holes than the one arising from cosmic censorship. An analysis of the unique form which such interactions must have in order to make a leading order contribution to the

absorption of the ℓ^{th} partial wave demonstrates that the correct energy dependence arises from the finite temperature kinematics. This demonstration, together with the general proof of the Optical Theorem for absorption of scalars given in section 4.2, can be viewed as a full investigation of the kinematics involved in higher partial waves. What is left is to calculate the coefficients $C_{D_1 \dots D_\ell}^{\dot{D}_1 \dots \dot{D}_\ell}$ and thereby verify or falsify (4.49), on which agreement between General Relativity and the effective string model relies.

The means to achieve a clear description of the dynamics and hopefully a derivation of the $C_{D_1 \dots D_\ell}^{\dot{D}_1 \dots \dot{D}_\ell}$ is a more precise treatment of the low-energy SCFT dictating the dynamics of the D1-D5-brane bound state. The effective string might continue to be a useful picture, perhaps supplemented by rules governing how different strands of the effective string interact.

In a way the finding that the effective string tension needs to scale as $1/\sqrt{Q_1 Q_5}$ is a more serious difficulty than the vanishing of $\ell > 1$ cross sections. Indeed, the necessity of choosing this peculiar value for the tension appears already at $\ell = 1$, where the naive effective string model in other ways seems completely adequate. A study of T-duality in [104] led to the conclusion that a tension scaling as $1/\sqrt{Q_1 Q_5}$ is more natural than $1/Q_1$ or $1/Q_5$, and it was further described how a simple modification in the calculation of disk diagrams would lead to such a scaling. However, in the absence of a first-principles derivation of the tension, a $1/(Q_1 + Q_5)$ scaling seems equally natural. This scaling is the one favored by entropy and temperature arguments. Thus it appears that a single energy scale does not fully characterize effective strings in the way that α' does fundamental strings.

Although effective string models of black holes have recently enjoyed a number of remarkable successes, a unifying picture has been slow in emerging. The General Relativity calculations for near-extremal black holes, on which most of the evidence for effective strings is based, are conceptually straightforward. The difficulty of studying bound states of solitons has caused the link between fundamental string theory and effective strings to remain imprecise in certain respects—most importantly in the interaction between the effective string and fields in the bulk of spacetime.

The puzzles presented by higher partial waves may push effective string theory in the directions it needs to go in order to become a fully viable model of near-extremal black holes. Discrepant results for the tension may be a clue to nature of effective string's interactions with bulk fields. Rescuing the $\ell > 1$ cross-sections must surely lead to a consideration of the multi-strand interactions which are as yet virgin territory in the theory of effective strings. Already, the absorption of higher partial waves into five-dimensional black holes is the cleanest dynamical test of the role of fermions on the effective string. Achieving a full understanding of these processes would constitute a major advance, not only in establishing the viability of effective strings in black hole physics, but also in comprehending the D1-D5-brane bound state.

Chapter 5

Photons and fermions falling into four-dimensional black holes

5.1 Introduction

Microscopic models of near-extremal black holes in terms of effective strings have recently been employed with great success to explain the Bekenstein-Hawking entropy [5, 6, 54, 51, 72]. These models have also proven their value by correctly predicting certain Hawking emission rates and absorption cross-sections at low energies [7, 8, 73, 99, 87, 21, 103, 102, 110, 112, 101, 75, 100, 113, 114].

The many successful predictions have all been for scalar particles. Most in fact have been for minimally coupled scalars. Usually it is said that minimally coupled scalars are those whose equation of motion is $\square \phi = 0$. This is a slight misnomer: a more precise way to say it is that when the equations of motion are linearized in small fluctuations around the black hole solution under consideration, one of them is simply $\square \delta\phi = 0$. Fortunately, it is usually easy to see which scalars in the theory are minimally coupled: for example, when one obtains a 5-dimensional black hole by toroidally compactifying the D1-brane D5-brane bound state, the off-diagonal gravitons with both indices lying within the D5-brane but perpendicular to the D1-brane are clearly minimally coupled scalars in the 5-dimensional theory.

This chapter, based on the paper [23], presents a study of particles with nonzero spin falling into effective string black holes. As a starting point, it is easiest to consider minimally coupled photons or fermions. For photons, minimally coupled means that the relevant linearized equation of motion is

$$\nabla_\mu \delta F^{\mu\nu} = 0 , \quad (5.1)$$

and for chiral fermions it means

$$\sigma_\mu^{\alpha\beta} \nabla^\mu \delta\psi_\alpha = 0 . \quad (5.2)$$

Minimally coupled fermions in arbitrary dimensions were studied in [71]. The authors of [71] correctly point out that the fermions in supergravity theories do not in general obey minimally coupled equations of motion in the presence of charged black holes. Likewise, it is not usually the case that photons will be minimally coupled in the presence of a supergravity black hole solution. Indeed, it seems that the gauge fields which carry the charges of the black hole are never minimally coupled: there is mixing between them and the graviton.

However, for the case we shall study in this chapter, namely the equal charge black hole [59] of $\mathcal{N} = 4$ supergravity [115, 116, 117, 118], two of the four Weyl fermions are in fact minimally coupled, as are four of the six gauge fields. The same black hole provided the simplest framework in which

to study fixed scalars [70]. Its metric is that of an extreme Reissner-Nordstrom black hole. In [100] it was shown that an effective string model is capable of reproducing the minimally coupled scalar cross-section of the Kerr-Newman metric, in the near-extremal limit. The effective string picture carries over naturally to the equal charge extreme black hole in $\mathcal{N} = 4$ supergravity and its near-extremal generalization. The recent work [114] presents evidence that the effective string can model a much broader class of black holes which have arbitrary $U(1)$ charges and are far from extremality. The essential features of the effective string, however, seem much the same in all its four-dimensional applications. We will show that minimally coupled fermions can be incorporated naturally into the effective string picture through a coupling to the supercurrent. Minimally coupled photons fit in in a somewhat unexpected way: the coupling of the gauge field to the string seems to occur via the field strength rather than the gauge potential.

The organization of this chapter is as follows. In section 5.2, the black hole solutions are exhibited and the minimally coupled photons are identified. In section 5.3, separable equations are derived for these photons. In section 5.4, these equations are solved to yield absorption cross-sections. The parallel analysis of minimally coupled fermions is postponed to section 5.5, in which also the axion cross-section is computed. The axion turns out to have the same cross-section as the dilaton, not because it is a fixed scalar in the usual sense of attractors [66, 67], but because of a dynamical version of the Witten effect [119]. Section 5.6 discusses the effective string interpretation of these cross-sections. Although the overall normalizations of the cross-sections are not computed, it is shown that the effective string correctly reproduces the relative normalization of the dilaton, axion, and minimal fermion cross-sections. Some concluding remarks are made in section 5.7. The appendix presents some results of the dyadic index formalism needed for the rest of the chapter.

5.2 Minimally coupled photons in $\mathcal{N} = 4$ supergravity

The fields of the $SU(4)$ version of $\mathcal{N} = 4$, $d = 4$ supergravity [118] are the graviton e_μ^a , four Majorana gravitinos ψ_μ^i , three vector fields A_μ^n , three axial vectors B_μ^n , four Majorana fermions χ^i , the dilaton ϕ , and the axion B . The doubly extreme black hole of [59] is electrically charged under A_3^μ , magnetically charged under B_3^μ , and neutral with respect to the other four gauge fields. These four extra gauge fields are minimally coupled photons, as we shall see shortly.

The full bosonic lagrangian of $\mathcal{N} = 4$ supergravity in the $SU(4)$ picture is

$$\begin{aligned} \mathcal{L} = \sqrt{-g} \Big[& -R + 2(\partial_\mu \phi)^2 + 2e^{4\phi}(\partial_\mu B)^2 - e^{-2\phi} \sum_n (F_n^2 + G_n^2) \\ & - 2iB \sum_n (F_n * F_n + G_n * G_n) \Big] \end{aligned} \quad (5.3)$$

where $F_n = dA_n$ and $G_n = dB_n$. The conventions used here are those of [59]. In particular, Hodge duals are defined by

$$*F_{\mu\nu} = \frac{\sqrt{-g}}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma} \quad (5.4)$$

where $\epsilon_{0123} = \epsilon_{tr\theta\phi} = -i$. The equations of motion following from (5.3) are

$$\begin{aligned} \nabla_\mu (e^{-2\phi} F_n^{\mu\nu} + 2iB * F_n^{\mu\nu}) &= 0 \\ \nabla_\mu (e^{-2\phi} G_n^{\mu\nu} + 2iB * G_n^{\mu\nu}) &= 0 \\ \square \phi - \frac{1}{2} e^{-2\phi} \sum_n (F_n^2 + G_n^2) - 2e^{4\phi} (\partial_\mu B)^2 &= 0 \\ \square B + 4\partial^\mu \phi \partial_\mu B + \frac{i}{2} e^{-4\phi} \sum_n (F_n * F_n + G_n * G_n) &= 0 \\ R_{\mu\nu} + 2\partial_\mu \phi \partial_\nu \phi + 2e^{4\phi} \partial_\mu B \partial_\nu B - e^{-2\phi} \sum_n (2F_{n\mu\lambda} F_{n\nu}{}^\lambda - \frac{1}{2} g_{\mu\nu} F_n^2 \\ &+ 2G_{n\mu\lambda} G_{n\nu}{}^\lambda - \frac{1}{2} g_{\mu\nu} G_n^2) = 0. \end{aligned} \quad (5.5)$$

Including the fermions introduces extra terms into these equations involving fermion bilinears. These terms affect neither the black hole solution nor the linearized bosonic equations around that solution since the fermions' background values are zero. Duals of the field strengths F_n and G_n are defined by

$$\tilde{F} = ie^{-2\phi} * F - 2BF . \quad (5.6)$$

$\tilde{F}_n$ and $\tilde{G}_n$ are closed forms by the equations of motion (5.5). In the $SO(4)$ version of $\mathcal{N} = 4$ supergravity, one writes $\tilde{G}_n = d\tilde{B}_n$.

The equal charge, axion free, extreme black hole solution is

$$\begin{aligned} ds^2 &= \frac{1}{(1 + M/r)^2} dt^2 - (1 + M/r)^2 (dr^2 + r^2 d\Omega^2) \\ \tilde{F}_3 &= Q \text{vol}_{S^2} \quad G_3 = P \text{vol}_{S^2} \\ e^{2\phi} &= 1 \quad B = 0 . \end{aligned} \quad (5.7)$$

The electric charge Q , the magnetic charge P , and the mass M are related by $Q = P = M/\sqrt{2}$. By definition, $\text{vol}_{S^2} = \sin \theta d\theta \wedge d\phi$. Gauss' law for the electric and magnetic charges reads

$$\begin{aligned} \int_{S^2} F_3 &= 0 & \int_{S^2} \tilde{F}_3 &= 4\pi Q \\ \int_{S^2} G_3 &= 4\pi P & \int_{S^2} \tilde{G}_3 &= 0 . \end{aligned} \quad (5.8)$$

Keeping the charges Q and P fixed but increasing the mass, one obtains the non-extremal generalization of (5.7):

$$\begin{aligned} ds^2 &= \frac{h}{f^2} dt^2 - f^2 (h^{-1} dr^2 + r^2 d\Omega^2) \\ \tilde{F}_3 &= Q \text{vol}_{S^2} \quad G_3 = P \text{vol}_{S^2} \\ e^{2\phi} &= 1 \quad B = 0 \end{aligned} \quad (5.9)$$

where

$$h = 1 - \frac{r_0}{r} \quad f = 1 + \frac{r_0 \sinh^2 \alpha}{r} . \quad (5.10)$$

The mass, charges, area, and temperature of this black hole are given by

$$\begin{aligned} M &= \frac{r_0}{2} \cosh 2\alpha & Q = P &= \frac{r_0}{2\sqrt{2}} \sinh 2\alpha \\ A &= 4\pi r_0^2 \cosh^4 \alpha & T &= \frac{1}{4\pi r_0 \cosh^4 \alpha} . \end{aligned} \quad (5.11)$$

Taking $\alpha \rightarrow \infty$ with Q and P held fixed, one recovers the extremal solution (5.7).

A crucial property of (5.7) and (5.9), without which there will be no minimally coupled photons, is the vanishing of the dilaton. This happens only when $Q = P$. The situation is similar to the case of fixed scalars, where the $Q = P$ case [70] was much easier to deal with than the $Q \neq P$ case [83].

When the equations (5.5) are linearized around the solution (5.9), the variations $\delta F_3^{\mu\nu}$ and $\delta G_3^{\mu\nu}$ appear in all the equations of motion because the background values of F_3 and G_3 are nonzero. But because F_1 , F_2 , G_1 , and G_2 do have vanishing background values (as does the axion B) the variations of these gauge fields appear in the linearized equations only as

$$\nabla_\mu (e^{-2\phi} F^{\mu\nu}) = 0 , \quad (5.12)$$

where $F = \delta F_1$, δF_2 , δG_1 , or δG_2 . Including $e^{-2\phi}$ in (5.12) was unnecessary since it is identically 1 when the charges are equal; but (5.12) is still the right linearized equation of motion for these fields when the charges are unequal (provided the background value of the axion remains zero). Surprisingly enough, a non-constant dilaton background makes it much more difficult to decouple the equations for different components of the gauge field.

5.3 Separable equations for gauge fields

Having shown that minimally coupled gauge fields do indeed exist, let us now show how to solve their equations of motion. The dilaton background will be kept arbitrary just long enough to observe why it makes the job much harder. The goal of this section is to convert Maxwell's equations

$$dF = 0 \quad d * e^{-2\phi} F = 0 \quad (5.13)$$

into decoupled separable differential equations. The equation of motion for the vector potential A ,

$$d * e^{-2\phi} dA = 0, \quad (5.14)$$

does not lend itself to this task. It turns out to be easier to dispense with A altogether and analyze Maxwell's equations, (5.13), directly. Even this is quite challenging if one sticks to the traditional tools of tensor analysis. Fortunately, several authors [120, 121, 122] in the 60's and 70's worked out an elegant approach to this sort of problem using Penrose's dyadic index formalism [123]. Appendix A provides a summary of some of the standard notation.

The field strength $F_{\mu\nu}$ (six real quantities) is replaced by a symmetric matrix $\Phi_{\Delta\Gamma}$ (three complex quantities) using the equation

$$F_{\mu\nu} \sigma_{\Delta\dot{\Delta}}^{\mu} \sigma_{\Gamma\dot{\Gamma}}^{\nu} = \Phi_{\Delta\Gamma} \epsilon_{\Delta\dot{\Gamma}} + \bar{\Phi}_{\dot{\Delta}\dot{\Gamma}} \epsilon_{\Delta\Gamma}. \quad (5.15)$$

Now define ϕ_1 , ϕ_0 , and ϕ_{-1} as follows:

$$\begin{aligned} \phi_1 &= \Phi_{00} = F_{\mu\nu} \ell^{\mu} m^{\nu} \\ \phi_0 &= \Phi_{10} = \Phi_{01} = \frac{1}{2} F_{\mu\nu} (\ell^{\mu} n^{\nu} + \bar{m}^{\mu} m^{\nu}) \\ \phi_{-1} &= \Phi_{11} = F_{\mu\nu} \bar{m}^{\mu} n^{\nu} \end{aligned} \quad (5.16)$$

where ℓ^{μ} , n^{μ} , m^{μ} , and $\bar{m}^{\mu}$ form the complex null tetrad (see the appendix). In the literature, it is more common to write ϕ_0 , ϕ_1 , and ϕ_2 instead of ϕ_1 , ϕ_0 , and ϕ_{-1} . The present convention has the advantage that the subscript is essentially the helicity.

The Bianchi identity $dF = 0$ can be rewritten as $D^{\Gamma\dot{\Delta}} \Phi_{\Delta\Gamma} = D^{\Delta\dot{\Gamma}} \bar{\Phi}_{\dot{\Gamma}}^{\dot{\Delta}}$. Using this identity one can rewrite the equation of motion $d * e^{-2\phi} F = 0$ as

$$D^{\Gamma\dot{\Delta}} \Phi_{\Delta\Gamma} = \frac{1}{2} (\partial_{\Gamma\dot{\Gamma}} e^{-2\phi}) (\Phi^{\Delta\Gamma} \epsilon^{\dot{\Delta}\dot{\Gamma}} + \bar{\Phi}^{\dot{\Delta}\dot{\Gamma}} \epsilon^{\Delta\Gamma}). \quad (5.17)$$

One immediately sees that the equations simplify greatly if the coupling $e^{-2\phi} = 1$. If this is not the case, then because the right hand side involves $\bar{\Phi}_{\dot{\Delta}\dot{\Gamma}}$ as well as $\Phi_{\Delta\Gamma}$, the advantage of compressing the real field strength components into complex components of $\Phi_{\Delta\Gamma}$ is lost. In this case, it seems difficult to decouple the equations. It is striking that the condition for Maxwell's equations to be simple is the same as the condition found in [70] and in chapter 3 for the fixed scalar equation to decouple from Einstein's equations.

Let us proceed with the case where $e^{-2\phi} = 1$, so that Maxwell's equations can be succinctly written as $D^{\Delta\dot{\Delta}} \Phi_{\Delta\Gamma} = 0$. The spin coefficients for the general spherically symmetric metric,

$$ds^2 = e^{2A(r)} dt^2 - e^{2B(r)} dr^2 - e^{2C(r)} (d\theta^2 + \sin^2 \theta d\phi^2) \quad (5.18)$$

are presented in (5.114). Six of them vanish, and the remaining six can be expressed in terms of γ , ρ , and α , which are real. Maxwell's equations written out in components therefore take on a particularly simple form:

$$\begin{aligned} (\Delta - 2\gamma + \rho)\phi_1 &= \delta\phi_0 \\ (D - 2\rho)\phi_0 &= (\bar{\delta} - 2\alpha)\phi_1 \\ (\Delta + 2\rho)\phi_0 &= (\delta - 2\alpha)\phi_{-1} \\ (D + 2\gamma - \rho)\phi_{-1} &= \bar{\delta}\phi_0. \end{aligned} \quad (5.19)$$

The form of Maxwell's equations in a more general metric can be found in [123].

A straightforward generalization of the preceding treatment can be given for fields of arbitrary nonzero spin. The simplest Lorentz covariant wave equation for a massless field of spin $n/2$ is

$$D^{\Delta_1 \dot{\Delta}} \Psi_{\Delta_1 \dots \Delta_n} = 0 \quad (5.20)$$

where $\Psi_{\Delta_1 \dots \Delta_n}$ is symmetric in all its indices. The case $n = 1$ gives the Weyl fermion equation. The case $n = 2$ is, as we have seen, Maxwell's equations in vacuum. The case $n = 4$ can be obtained by linearizing pure gravity around Minkowski space, as discussed in section 5.7 of [124]. The case $n = 3$ can be obtained in Minkowski space from the massless Rarita-Schwinger equation, as follows. The constraint $\gamma^\mu \psi_\mu = 0$ is imposed on the Rarita-Schwinger field

$$\psi^\mu = \begin{pmatrix} \sigma^{\mu \dot{\Delta} \Delta} \psi_{\Delta \Gamma \dot{\Delta}} \\ \sigma^{\mu \dot{\Delta} \Delta} \bar{\psi}_{\dot{\Delta}}^{\dot{\Gamma}}{}_{\Delta} \end{pmatrix} \quad (5.21)$$

to project out the spin-1/2 components. This constraint is equivalent to making $\psi_{\Delta \Gamma \dot{\Delta}}$ symmetric in its two undotted indices. In the supergravity literature, the equation of motion is usually written as $\epsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\nu \Psi_{\rho\sigma} = 0$ where $\Psi_{\rho\sigma} = \partial_\rho \psi_\sigma - \partial_\sigma \psi_\rho$. The original paper by Rarita and Schwinger [125] (see also p. 323 of [126]) proposes $\not{\partial} \psi_\mu = 0$ as the equation of motion. Using the constraint one can show that both are equivalent to $\partial^{\dot{\Delta} \Delta} \psi_{\Delta \Gamma \dot{\Sigma}} = 0$. As a result, the field strength

$$\Psi_{\Delta \Sigma \Gamma} = \partial_{\Sigma \dot{\Delta}} \psi_{\Delta}^{\dot{\Delta}}{}_{\Gamma} \quad (5.22)$$

is symmetric in all its indices and obeys the $n = 3$ case of (5.20).

Although the cases $n = 3$ and $n = 4$ of (5.20) are not in general the correct curved-space equations of motion for the gravitino and graviton, and although for $n > 2$ there are problems defining local, gauge-invariant number currents and stress-energy tensors, still a brief investigation of (5.20) serves to illustrate some of the general features one expects for fields of higher spin. Furthermore, the near-Minkowskian limit of the equations we will derive should be close in form to the actual graviton and gravitino equations far from a black hole.

Define helicity components ψ_s according to

$$\Psi_{\Delta_1 \dots \Delta_n} = \psi_{\frac{n}{2} - \sum_i \Delta_i} \quad (5.23)$$

Then (5.20) can be written out in components. There are $2n$ equations:

$$\begin{aligned} (\Delta - n\gamma + \rho) \psi_{\frac{n}{2}} &= (\delta + (n-2)\alpha) \psi_{\frac{n}{2}-1} \\ (D - (n-2)\gamma - n\rho) \psi_{\frac{n}{2}-1} &= (\bar{\delta} - n\alpha) \psi_{\frac{n}{2}} \\ &\vdots \\ (\Delta - 2s\gamma + (\frac{n}{2} + 1 - s)\rho) \psi_s &= (\delta + (2s-2)\alpha) \psi_{s-1} \\ (D - (2s-2)\gamma - (\frac{n}{2} + s)\rho) \psi_{s-1} &= (\bar{\delta} - 2s\alpha) \psi_s \\ &\vdots \\ (\Delta + (n-2)\gamma + n\rho) \psi_{-\frac{n}{2}+1} &= (\delta - n\alpha) \psi_{-\frac{n}{2}} \\ (D + n\gamma - \rho) \psi_{-\frac{n}{2}} &= (\bar{\delta} + (n-2)\alpha) \psi_{-\frac{n}{2}+1} \end{aligned} \quad (5.24)$$

The equations (5.24) are invariant under PT, which sends $\psi_s \rightarrow \psi_{-s}$, $\gamma \rightarrow -\gamma$, $\rho \rightarrow -\rho$, $D \leftrightarrow \Delta$, $\delta \leftrightarrow \bar{\delta}$.

The commutation relations

$$[D, \delta] = \rho\delta \quad [D, \alpha] = \rho\alpha \quad [\Delta, \bar{\delta}] = -\rho\bar{\delta} \quad [\Delta, \alpha] = -\rho\alpha \quad (5.25)$$

are easily established by direct computation. They can be used to convert the pair of equations in (5.24) relating ψ_s and ψ_{s-1} into decoupled second order equations for ψ_s and ψ_{s-1} separately. In this way one obtains

$$\begin{aligned} & \left[(D - (2s - 2)\gamma - (\frac{n}{2} + 1 + s)\rho)(\Delta - 2s\gamma + (\frac{n}{2} + 1 - s)\rho) \right. \\ & \quad \left. - (\delta + (2s - 2)\alpha)(\bar{\delta} - 2s\alpha) \right] \psi_s = 0 \\ & \left[(\Delta - (2s + 2)\gamma + (\frac{n}{2} + 1 - s)\rho)(D - 2s\gamma - (\frac{n}{2} + 1 + s)\rho) \right. \\ & \quad \left. - (\bar{\delta} - (2s + 2)\alpha)(\delta + 2s\alpha) \right] \psi_s = 0 . \end{aligned} \quad (5.26)$$

The first of these can be derived for $s > -n/2$, while the second can be derived for $s < n/2$. In fact they are different forms of the same equation, which can be written out more simply in terms of the fields

$$\tilde{\psi}_s = e^{|s|A + (\frac{n}{2} - |s| + 1)C} \psi_s \quad (5.27)$$

as

$$\begin{aligned} & \left[(D + (2 - 4s)\gamma - 2s\rho)\Delta - (\delta + (2s - 2)\alpha)(\bar{\delta} - 2s\alpha) \right] \tilde{\psi}_s = 0 \quad \text{for } s \geq 0 \\ & \left[(\Delta - (2 + 4s)\gamma - 2s\rho)D - (\bar{\delta} - (2s + 2)\alpha)(\delta + 2s\alpha) \right] \tilde{\psi}_s = 0 \quad \text{for } s \leq 0. \end{aligned} \quad (5.28)$$

More explicitly,

$$\begin{aligned} & \left[\partial_r^2 + ((1 - 2|s|)A' - B' + 2|s|C')\partial_r - e^{-2A+2B}\partial_t^2 + 2s e^{-A+B}(A' - C')\partial_t \right. \\ & \quad \left. + e^{2B-2C}(\partial_\theta^2 + \cot\theta\partial_\theta + \csc^2\theta\partial_\phi^2 + 2is\cot\theta\csc\theta\partial_\phi - s^2\cot^2\theta - |s|) \right] \tilde{\psi}_s = 0 \end{aligned} \quad (5.29)$$

for all values of s , positive and negative. It is interesting to note that there is no explicit dependence on the spin $n/2$ of the particle in (5.29), only on its helicity s . In practice we will mainly be interested in the equations for $\tilde{\psi}_{\pm\frac{n}{2}}$ since these are the only components that can be radiative. The fact that the equations for these radiative fields are identical to equations obeyed by non-radiative components of fields of higher spin suggests that mixing of different spins is possible. Such mixing between photons and gravitons was observed by Chandrasekhar in his analysis of perturbations of the Reissner-Nordstrom black hole [127].

Equations similar to (5.29) were worked out for the Kerr metric by Teukolsky [120, 121]. In that case, only the equations for the radiative fields turned out to be separable. But in the present context, spherical symmetry makes the separability of (5.29) trivial: the general solution is

$$\tilde{\psi}_s(t, r, \theta, \phi) = e^{-i\omega t} R_{s\ell}(r) Y_{s\ell m}(\theta, \phi) \quad (5.30)$$

where $Y_{s\ell m}$ is a spin-weighted spherical harmonic [128] and $R_{s\ell}(r)$ satisfies the ODE

$$\begin{aligned} & \left[\partial_r^2 + ((1 - 2|s|)A' - B' + 2|s|C')\partial_r + \omega^2 e^{-2A+2B} - 2si\omega e^{-A+B}(A' - C') \right. \\ & \quad \left. - e^{2B-2C}(\ell + |s|)(\ell - |s| + 1) \right] R_{s\ell} = 0 . \end{aligned} \quad (5.31)$$

The minimal value of ℓ is $|s|$. In the case of photons, $\ell \geq 1$ indicates that the fields of lowest moment that can be radiated are dipole fields. For fermions, ℓ , s , and m are all half-integer.

5.4 Semi-classical absorption probabilities

Returning now to the case of photons, let us investigate how an absorption cross-section can be extracted from a solution to (5.31). Section 5.4.1 derives a formula for the absorption probability. In section 5.4.2 matching solutions are exhibited and absorption probabilities calculated for the black holes (5.7) and (5.9).

5.4.1 Probabilities from energy fluxes

For the photon as for other fields of spin greater than $1/2$, there is no gauge-invariant number current analogous to $J_\mu = \frac{1}{2i} \bar{\phi} \overleftrightarrow{\partial}_\mu \phi$ for spin 0 and $J_\mu = \bar{\psi} \gamma_\mu \psi$ for spin $1/2$. In order to count the photons falling into the black hole, it is therefore necessary to examine the energy flux through the horizon and adjust for the gravitational blueshift that the infalling photons experience. The stress-energy tensor can be written in terms of ϕ_1 , ϕ_0 , and ϕ_{-1} :

$$\begin{aligned} T^{\mu\nu} &= \frac{1}{4} g^{\mu\nu} F^2 + F^{\mu\rho} F_\rho{}^\nu = 2\sigma^\mu \Delta \dot{\Delta} \sigma^\nu \Gamma \dot{\Gamma} \phi_{\Delta\Gamma} \bar{\phi}_{\dot{\Delta}\dot{\Gamma}} \\ &= \left[|\phi_1|^2 n_\mu n_\nu + 2|\phi_0|^2 (\ell_{(\mu} n_{\nu)} + m_{(\mu} \bar{m}_{\nu)}) + |\phi_{-1}|^2 \ell_\mu \ell_\nu \right. \\ &\quad \left. - 4\bar{\phi}_1 \phi_0 n_{(\mu} m_{\nu)} - 4\bar{\phi}_0 \phi_{-1} \ell_{(\mu} m_{\nu)} + 2\phi_{-1} \bar{\phi}_1 m_\mu m_\nu \right] + \text{c.c.} \end{aligned} \quad (5.32)$$

Consider a sphere S^2 located anywhere outside the horizon. Taking into account the blueshift factor as described, the number of photons passing through S^2 in a time interval $[0, t]$ is

$$N = \frac{1}{\omega} \int_{S^2 \times [0, t]} *(T_{tr} dr) = \frac{t}{\omega} \int_{S^2} \mathcal{F} \text{vol}_{S^2} \quad (5.33)$$

where $\text{vol}_{S^2} = \sin\theta d\theta \wedge d\phi$, and

$$\mathcal{F} = e^{A-B+2C} T_{tr} = e^{2A+2C} (|\phi_{-1}|^2 - |\phi_1|^2) = |\tilde{\phi}_{-1}|^2 - |\tilde{\phi}_1|^2 \quad (5.34)$$

is essentially the radial photon number flux.

The goal now is to find an approximate solution to (5.29) for photons whose wavelength is much longer than the size of the black hole, and to extract from it an absorption probability and cross-section. The dominant contribution to this absorption comes from dipole fields.

Far from the black hole, (5.31) for dipole fields simplifies to

$$\left[\partial_\rho^2 + \frac{2}{\rho} \partial_\rho + 1 + \frac{2si}{\rho} - \frac{2}{\rho^2} \right] R = 0 \quad (5.35)$$

where $\rho = \omega r$. The general solution to (5.35) with $s = 1$ is

$$R = 2ae^{-i\rho} \left(1 - \frac{i}{\rho} - \frac{1}{2\rho^2} \right) + b \frac{e^{i\rho}}{\rho^2}. \quad (5.36)$$

The general solution with $s = -1$ is just the conjugate of (5.36). By using (5.19) ϕ_0 can be calculated as well. Let us choose the spatial orientation by setting $m = 0$ in (5.30). Then the final result is

$$\begin{aligned} \tilde{\phi}_1 &= e^{-i\omega t} \sin\theta \left[2ae^{-i\rho} \left(1 - \frac{i}{\rho} - \frac{1}{2\rho^2} \right) + b \frac{e^{i\rho}}{\rho^2} \right] \\ \tilde{\phi}_0 &= e^{-i\omega t} \cos\theta \frac{\sqrt{2}i}{\omega} \left[ae^{-i\rho} \left(1 - \frac{i}{\rho} \right) + be^{-i\rho} \left(1 + \frac{i}{\rho} \right) \right] \\ \tilde{\phi}_{-1} &= e^{-i\omega t} \sin\theta \left[a \frac{e^{-i\rho}}{\rho^2} + 2be^{i\rho} \left(1 + \frac{i}{\rho} - \frac{1}{2\rho^2} \right) \right]. \end{aligned} \quad (5.37)$$

It is clear from (5.37) that ϕ_{-1} is the radiative component of the field for outgoing waves, while ϕ_1 is the radiative component for ingoing waves.

The boundary conditions at the horizon [121] require the radial group velocity to point inward. It can be shown that $\tilde{\phi}_1$ remains finite at the horizon while $\tilde{\phi}_{-1}$ vanishes. Normalizations are fixed by

requiring $|\tilde{\phi}_1|^2 \rightarrow \sin^2 \theta$ at the horizon. The net flux of photons into the black hole can be computed in two ways:

$$\begin{aligned} \text{at the horizon: } \mathcal{F}_h &= -|\tilde{\phi}_1|^2 = -\sin^2 \theta \\ \text{at infinity: } \mathcal{F}_\infty &= \mathcal{F}_\infty^{\text{out}} + \mathcal{F}_\infty^{\text{in}} = |\tilde{\phi}_{-1}|^2 - |\tilde{\phi}_1|^2 = 4(|b|^2 - |a|^2) \sin^2 \theta . \end{aligned} \quad (5.38)$$

The two must agree, $\mathcal{F}_h = \mathcal{F}_\infty$, and so $|a|^2 = |b|^2 + 1/4$. One can thus easily perceive the equivalence of the two common methods for computing the absorption probability. The first examines the deficit in the outgoing flux compared to the ingoing flux:

$$1 - P = \frac{\mathcal{F}_\infty^{\text{out}}}{\mathcal{F}_\infty^{\text{in}}} = \frac{|b|^2}{|a|^2} , \quad (5.39)$$

while the second simply compares the flux on the horizon to the ingoing flux at infinity:

$$P = \frac{\mathcal{F}_h}{\mathcal{F}_\infty^{\text{in}}} = \frac{1}{4|a|^2} . \quad (5.40)$$

For low-energy photons, the absorption probability is small and a and b are large and nearly equal, so from a calculational point of view the second method is to be preferred over the first. Indeed, it will be standard practice in the matching calculations of later sections to ignore the small difference between a and b and simply set them equal. This approximation suffices when (5.40) is used.

Finally, to obtain the absorption cross-section from the probability, the Optical Theorem is needed. Averaging over polarizations is unnecessary in view of the spherical symmetry of the background. The result for photons in a dipole wave is

$$\sigma_{\text{abs}} = \frac{3\pi}{\omega^2} P . \quad (5.41)$$

5.4.2 Matching solutions

The work of previous sections can be boiled down to a simple prescription for computing the absorption probability and cross-section to leading order in the energy for minimally coupled photons falling into a spherically symmetric black hole. The probability can be obtained by solving (5.31) with $\ell = 1$ and $s = 1$, subject to the boundary condition $R(r) \sim e^{if(r)}$ as r approaches the horizon, $f(r)$ being some real decreasing function of r . Far from the black hole, one will find $R(r) \sim 2ae^{-i\omega r}$, and the probability is then given by

$$P = \frac{1}{4|a|^2} = \frac{|R(r)|^2 \Big|_{\text{horizon}}}{|R(r)|^2 \Big|_{\infty}} . \quad (5.42)$$

First consider the extreme black hole (5.7). The radial equation (5.31) with $\ell = s = 1$ is

$$\left[\partial_r^2 + \frac{2}{r+M} \partial_r + \omega^2 \left(1 + \frac{M}{r} \right)^4 + \frac{2i\omega}{r} \left(1 - \frac{M^2}{r^2} \right) - \frac{2}{r^2} \right] R = 0 . \quad (5.43)$$

A different radial variable, $y = \omega M^2/r$, is more natural near the horizon. In terms of y , (5.43) can be rewritten as

$$\left[(y^2 \partial_y)^2 - \frac{2\omega M}{y + \omega M} y^3 \partial_y + (y + \omega M)^4 - 2iy(y^2 - \omega^2 M^2) - 2y^2 \right] R = 0 . \quad (5.44)$$

A matching solution can be pieced together as usual from a near region (**I**), an intermediate region (**II**), and a far region (**III**). In the near region, (5.44) is simplified by setting to zero all terms containing

explicit factors of ωM . The intermediate region solution is obtained from (5.43) with $\omega = 0$. In the far region, we simply make the flat space approximation, obtaining (5.35) and (5.36). The solutions in the three regions are

$$\begin{aligned} R_{\text{I}} &= e^{iy} \left(1 + \frac{i}{y} - \frac{1}{2y^2} \right) \\ R_{\text{II}} &= \frac{C_1 r}{1 + M/r} + \frac{C_2}{r^2(1 + M/r)} \\ R_{\text{III}} &= 2ae^{-i\rho} \left(1 - \frac{i}{\rho} - \frac{1}{2\rho^2} \right) + b \frac{e^{i\rho}}{\rho^2} . \end{aligned} \quad (5.45)$$

A match is obtained by setting

$$\begin{aligned} C_1 &= -\frac{1}{2\omega^2 M^3} & C_2 &= 0 \\ a &= b = -\frac{3i}{8(\omega M)^3} . \end{aligned} \quad (5.46)$$

The absorption probability and cross-section are

$$P = \frac{16}{9}(\omega M)^6 \quad \sigma_{\text{abs}} = \frac{16\pi}{3}\omega^4 M^6 . \quad (5.47)$$

Now consider the non-extremal generalization, (5.9). The radial equation (5.31) with $\ell = 1$ and $s = 1$ is

$$\left[\partial_r^2 + \frac{2}{fr} \partial_r + \omega^2 \frac{f^4}{h^2} - 2i\omega \left(\frac{1}{2} + \frac{1}{2h} - \frac{2}{f} \right) - \frac{2}{hr^2} \right] R = 0 . \quad (5.48)$$

As before, the far region is treated in the flat space approximation, and a solution in the intermediate region is obtained by solving (5.48) with $\omega = 0$. In the near region, the useful radial variable is h itself. Having the black hole near extremality is useful since one can approximate $f \approx (1 - h) \cosh^2 \alpha$ and drop terms in (5.48) which are small in the limit where $\alpha \rightarrow \infty$ and $\omega r_0 \rightarrow 0$ with

$$\lambda = \omega r_0 \cosh^4 \alpha = \frac{\omega}{4\pi T} \quad (5.49)$$

held fixed. The result is that (5.48) simplifies to

$$\left[h(1 - h) \partial_h^2 - 2h \partial_h + \lambda^2 \frac{1 - h}{h} - i\lambda \frac{1 + h}{h} - \frac{2}{1 - h} \right] R = 0 , \quad (5.50)$$

which is representative of the general form of differential equation which is solved by a hypergeometric function of h times powers of h and $1 - h$.

The solutions in the three regions are

$$\begin{aligned} R_{\text{I}} &= \frac{h^{-i\lambda}}{(1 - h)^2} F(-2, -1 - 2i\lambda, -2i\lambda; h) \\ &= \frac{h^{-i\lambda}}{(1 - h)^2} \left(1 + \frac{ih(1 + 2i\lambda)}{\lambda} - \frac{h^2(1 + 2i\lambda)}{1 - 2i\lambda} \right) \\ R_{\text{II}} &= C_1 \frac{h}{f(1 - h)} + C_2 \frac{h}{f} \left(1 + \frac{1}{h} + \frac{2 \log h}{1 - h} \right) \\ R_{\text{III}} &= 2ae^{-i\rho} \left(1 - \frac{i}{\rho} - \frac{1}{2\rho^2} \right) + b \frac{e^{i\rho}}{\rho^2} , \end{aligned} \quad (5.51)$$

and a match is obtained by setting

$$\begin{aligned} C_1 &= \frac{i \cosh^2 \alpha}{\lambda(1 - 2i\lambda)} & C_2 &= 0 \\ a = b &= -\frac{3 \cosh^2 \alpha}{4\omega r_0 \lambda(1 - 2i\lambda)} . \end{aligned} \tag{5.52}$$

The absorption probability and cross-section are

$$P = \frac{4}{9}(\omega r_0)^3 \lambda(1 + 4\lambda^2) \quad \sigma_{\text{abs}} = \frac{4\pi}{3} \omega r_0^3 \lambda(1 + 4\lambda^2) . \tag{5.53}$$

5.5 The axion and minimally coupled fermions

Because the solution (5.7) preserves a quarter of the supersymmetry, it is clearly of interest to compare cross-sections of particles with different spins related by the unbroken supersymmetry. The other particles in $\mathcal{N} = 4$ supergravity whose cross-sections are straightforward to compute are the dilaton, the axion, and those fermions which obey the Weyl equation. The dilaton has been dealt with at length in the fixed scalar literature [70, 96, 83]. The axion in fact is also a fixed scalar, as section 5.5.1 will show. Of the four massless fermions, two are minimally coupled. The purpose of section 5.5.2 is to demonstrate this fact and to compute the minimal fermion cross-section. For comparison with (5.53) let us quote here the final results:

$$\begin{aligned} \text{axion, dilaton:} \quad & P = (\omega r_0)^2(1 + 4\lambda^2) & \sigma_{\text{abs}} &= \pi r_0^2(1 + 4\lambda^2) \\ \text{minimal fermions:} \quad & P = \frac{(\omega r_0)^2}{4}(1 + 16\lambda^2) & \sigma_{\text{abs}} &= \frac{\pi r_0^2}{2}(1 + 16\lambda^2) \end{aligned} \tag{5.54}$$

where $\lambda = \omega/(4\pi T)$, as in (5.53). Note that in the extremal limit the bosonic and fermionic absorption probabilities quoted in (5.54) coincide.¹

5.5.1 The axion

The strategy for deriving the linearized equation of motion for the axion is the same as the one used in [70] for the dilaton: only spherical perturbations of the solution (5.9) are considered, and a gauge is chosen where the only components of the metric that fluctuate are g_{rr} and g_{tt} . The minimally coupled gauge fields F_1 , F_2 , G_1 , and G_2 do not affect the linearized axion equation because they have no background value and enter into the axion equation quadratically. Spherical symmetry dictates that only tr and $\theta\phi$ components of these field strengths can fluctuate, corresponding respectively to radial electric and radial magnetic field fluctuations. These fluctuations are constrained further by Gauss' law (5.8):

$$\begin{aligned} F_{\theta\phi}^3 &= 0 & F_3^{tr} &= \frac{Q \sin \theta}{\sqrt{-g}} e^{2\phi} \\ G_{\theta\phi}^3 &= P \sin \theta & G_3^{tr} &= \frac{2B\epsilon^{2\phi}}{\sqrt{-g}} G_{\theta\phi}^3 . \end{aligned} \tag{5.55}$$

The asymmetry between F_3 and G_3 arises because the black hole is electrically charged under F_3 and magnetically charged under G_3 . The axion B is a dynamical theta-angle for the gauge fields, so the last relation in (5.55) should be viewed as a dynamical version of the Witten effect [119]: when the

¹Similar results on the agreement of absorption probabilities due to residual supersymmetry have appeared elsewhere in the literature [129, 130] for the case of $N = 2$ supergravity. See also [131]. Thanks to G. Horowitz and A. Peet for bringing these papers to my attention.

axion fluctuates, an object that was magnetically charged picks up what seems like an electric charge in that there are radial electric fields.

Using (5.55) and ignoring the dilaton terms in the axion equation of (5.5) (which is valid for the purpose of deriving the linearized axion equation because the dilaton has zero background value), one obtains

$$\square B + \frac{i}{2}(F_3 * F_3 + G_3 * G_3) = \square B + i(G_3^{tr} * G_{tr}^3 + G_3^{\theta\phi} * G_{\theta\phi}^3) = \left[\square + \frac{4P^2}{f^4 r^4} \right] B = 0 . \quad (5.56)$$

This is indeed identical to the linearized equation for the dilaton, although the “mass” term for the dilaton receives equal contributions $2Q^2 + 2P^2$ from the electric and magnetic charges in place of the $4P^2$ we see in (5.56).

The radial equation for B and ϕ is

$$\left[(hr^2 \partial_r)^2 + \omega^2 r^4 f^4 - \frac{hr_0^2 \sinh^2 2\alpha}{2f^2} \right] R = 0 . \quad (5.57)$$

By an analysis sufficiently analogous to the treatments in [70, 96] that it seems superfluous to present the details, one obtains the result already quoted in (5.54):

$$P = (\omega r_0)^2 (1 + 4\lambda^2) \quad \sigma_{\text{abs}} = \pi r_0^2 (1 + 4\lambda^2) . \quad (5.58)$$

5.5.2 Minimally coupled fermions

It was shown in [118] that the complete fermionic equations of motion for simple $\mathcal{N} = 4$, $d = 4$ supergravity take on a simple form when written in terms of the supercovariant derivatives introduced in [117]. The relevant one of these equations for spin-1/2 fermions is

$$i\hat{D}\Lambda_I - \frac{3}{2}e^{2\phi}(\hat{D}B)\Lambda_I = 0 , \quad (5.59)$$

where $\hat{D}_\mu$ denotes a supercovariant derivative.²

Supercovariant derivatives in general can be read off from the supersymmetry variations of a field: if $\delta f = F_I \epsilon^I$, then $\hat{D}_\mu f = D_\mu f - \frac{1}{4}F_I \Psi_\mu^I$. Thus the supercovariant derivative of the axion is $\hat{D}_\mu B = \partial_\mu B + (\text{two fermion terms})$. Again, terms in the equations of motion which are quadratic in fields with zero background value do not contribute to the linearized first-varied equations of motion. Because the background values of the axion as well as all fermions are zero for the solution (5.9), the second term in (5.59) can be discarded.

A further simplification of (5.59) can be made by dropping terms from $\hat{D}\Lambda_I$ which are quadratic in fields with zero background value. The supersymmetry variation of Λ_I is

$$\delta\Lambda_I = \sqrt{2}\sigma^{\rho\sigma}(F_{\rho\sigma}^3\alpha_{IJ}^3 - \tilde{G}_{\rho\sigma}^3\beta_{IJ}^3)\epsilon^J \quad (5.60)$$

plus terms which vanish for the solution (5.9). So

$$\hat{D}_\mu\Lambda_I = D_\mu\Lambda_I - \frac{1}{2\sqrt{2}}\sigma^{\rho\sigma}(F_{\rho\sigma}^3\alpha_{IJ}^3 - \tilde{G}_{\rho\sigma}^3\beta_{IJ}^3)\Psi_\mu^J \quad (5.61)$$

²The spinor and gamma matrix conventions used here are those described in Appendix A of [59]. Λ_I is a chiral spinor with $\gamma_5\Lambda_I = \Lambda_I$ which replaces the Majorana spinor χ^i of [118]. The gravitinos are also written in terms of chiral spinors Ψ_μ^I with $\gamma_5\Psi_\mu^I = \Psi_\mu^I$. I runs from 1 to 4. Conversion to these conventions from those of [118] is discussed in [59]. It is worth adding that in the current conventions, each field is identified with K times its counterpart in [118], and for notational simplicity K is then set equal to 1/2. With this choice of the gravitational constant, ϕ and Λ_I are not canonically normalized; rather, they are twice the canonically normalized fields. So for example the kinetic term of ϕ in (5.3) is $2(\partial_\mu\phi)^2$ rather than the canonical $\frac{1}{2}(\partial_\mu\phi)^2$.

plus terms quadratic in fields with zero background value. In (5.60) and (5.61), we have introduced $\sigma^{\rho\sigma} = \frac{1}{4}[\gamma^\rho, \gamma^\sigma]$ and the matrices

$$\alpha_{IJ}^3 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \quad \beta_{IJ}^3 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \quad (5.62)$$

The matrices α_{IJ}^n and β_{IJ}^n were introduced in [132] and used in [118] to establish the $SU(4)$ invariance of $\mathcal{N} = 4$ supergravity.

Since $F_{\rho\sigma}^3 = \tilde{G}_{\rho\sigma}^3$, the first variation of the equation of motion (5.59) is

$$i\hat{D}\delta\Lambda_I = i\hat{D}\delta\Lambda_I - \frac{i}{2\sqrt{2}}\sigma^{\rho\sigma}F_{\rho\sigma}^3(\alpha_{IJ}^3 - \beta_{IJ}^3)\delta\Psi_\mu^J = 0. \quad (5.63)$$

For $I = 3$ and 4 , the gravitino part gets killed and (5.59) is nothing but the Weyl equation (5.2). For $I = 1$ and 2 , the gravitino mixes in.³

The key point in this analysis is that the supersymmetry transformation (5.60) leaves two of the Λ_I invariant no matter what ϵ^J is chosen to be. In view of the form of (5.59) and the prescription for reading off supercovariant derivatives from the supersymmetry transformation laws, this makes it inevitable that two of the Λ_I are minimally coupled fermions. The situation is related to unbroken supersymmetries, but only loosely: the non-extremal black hole has the same minimally coupled fermions that the extremal one does, and the extremal black hole preserves only one supersymmetry but admits two minimally coupled fermions. The condition for an unbroken supersymmetry is that the supersymmetry variations of all the fermions fields must vanish. So one would expect that there are at least as many minimally coupled fermions as unbroken supersymmetries. More precisely, suppose that there are n *broken* supersymmetries and m fermions in the theory whose equation of motion is of the form $i\hat{D}\Lambda = 0$, up to terms quadratic in fields with zero background value. Then there must be at least $m - n$ minimally coupled fermions. For axion-free solutions to pure $\mathcal{N} = 4$ supergravity, $m = 4$.

The existence of minimally coupled fermions having been established, the computation of their absorption cross-section now proceeds in parallel to the case of minimally coupled photons. If $R(r)$ is a solution of the radial equation (5.31) with $\ell = s = 1/2$, then the absorption probability is once again

$$P = \frac{|R(r)|^2|_{\text{horizon}}}{|R(r)|^2|_{\infty}}. \quad (5.64)$$

The justification for (5.64) is a little different than for its analog (5.42) for photons because for fermions there is a conserved number current,

$$J_\mu = -\sqrt{2}\sigma_\mu^{\Delta\Delta}\bar{\Psi}_\Delta\Psi_\Delta. \quad (5.65)$$

Here Ψ_Δ is the dyadic version of $\delta\Lambda_3$ or $\delta\Lambda_4$. The number of fermions passing through a sphere in a time t is

$$N = \int_{S^2 \times [0, t]} *(J_r dr) = t \int_{S^2} \mathcal{F} \text{vol}_{S^2} \quad (5.66)$$

³The gravitino equation of motion has the form of the Rarita-Schwinger equation plus interactions. This equation also has an $SO(4)$ index structure which is block diagonal, and the spin-1/2 particles decouple from the $I = 3, 4$ equations. The resulting gravitino equation, $\epsilon^{\mu\nu\rho\sigma}\gamma_\nu\hat{\Psi}_{\rho\sigma}^I = 0$, is less simple than the equations studied previously because the supercovariant field strength $\hat{\Psi}_{\rho\sigma}^I$ involves a non-vanishing combination of the field strengths F_3 and $\tilde{G}_3$. The problem of extracting from it a separable PDE like (5.29) is under investigation.

where

$$\mathcal{F} = e^{A-B+2C} J_r = e^{A+2C} (|\psi_{-1/2}|^2 - |\psi_{1/2}|^2) = |\tilde{\psi}_{-1/2}|^2 - |\tilde{\psi}_{1/2}|^2 . \quad (5.67)$$

The component $\psi_{1/2}$, like ϕ_1 in the case of photons, is both the radiative component at infinity for infalling solutions and the nonzero component at the black hole horizon. The formula (5.64) thus follows from (5.67) by the same analysis that gave (5.42) from (5.34).

The radial equation (5.31) with $\ell = s = 1/2$ is

$$\left[\partial_r^2 + \frac{1}{r} \left(\frac{1}{2} + \frac{1}{2h} \right) + \omega^2 \frac{f^4}{h^2} - i\omega \frac{f^2}{h} \frac{1}{r} \left(\frac{1}{2} + \frac{1}{2h} - \frac{2}{f} \right) - \frac{1}{hr^2} \right] R = 0 . \quad (5.68)$$

A matching solution can be obtained in the usual fashion:

$$\begin{aligned} R_{\text{I}} &= \frac{h^{-i\lambda}}{1-h} F(-1, -\frac{1}{2} - 2i\lambda, \frac{1}{2} - 2i\lambda; h) \\ &= \frac{h^{-i\lambda}}{1-h} \left(1 + \frac{1+4i\lambda}{1-4i\lambda} h \right) \\ R_{\text{II}} &= C_1 \frac{1+h}{1-h} + C_2 \frac{\sqrt{h}}{1-h} \\ R_{\text{III}} &= 2a e^{-i\rho} \left(1 - \frac{i}{2\rho} \right) + ib \frac{e^{i\rho}}{\rho} \end{aligned} \quad (5.69)$$

with

$$\begin{aligned} C_1 &= \frac{1}{1-4i\lambda} & C_2 &= 0 \\ a = b &= \frac{i}{\omega r_0} \frac{1}{1-4i\lambda} \end{aligned} \quad (5.70)$$

where as before $\lambda = \omega/(4\pi T)$. The absorption probability and cross-section are

$$P = \frac{(\omega r_0)^2}{4} (1 + 16\lambda^2) \quad \sigma_{\text{abs}} = \frac{2\pi}{\omega^2} P = \frac{\pi r_0^2}{2} (1 + 16\lambda^2) . \quad (5.71)$$

The $\lambda \rightarrow 0$ limit of (5.71) agrees with the general result of [71].

5.6 The effective string model

The usual approach to effective string calculations (see for example [8, 21, 99, 101]) has been to derive couplings between bulk fields and the effective string by expanding the Dirac-Born-Infeld (DBI) action to some appropriate order. The leading terms in the expansion specify a conformal field theory (CFT) which would describe the effective string in the absence of interactions with bulk fields. The interactions are dictated at leading order by terms in the expansion which are linear in the bulk fields: for a scalar field ϕ , a typical coupling would be

$$S_{\text{int}} = \int d^2x \phi(t, x, \vec{x}=0) \mathcal{O}(t, x) \quad (5.72)$$

where $\mathcal{O}(t, x)$ is some local conformal operator in the CFT. The integration is over the effective string world-volume, and $\vec{x}$ is set to zero because this is the location of the effective string in transverse space. One can then consider tree-level processes mediated by S_{int} where a bulk particle is converted into excitations on the effective string. Although the applicability of the DBI action to D-brane bound states can be called into question, the prescription described here for computing absorption or emission rates appears to be very robust. In the spirit of [100], where effective string calculations were

used to account for properties of Reissner-Nordstrom black holes without reference to any underlying microscopic picture, let us examine the consequences of couplings of the general form (5.72).

Consider the absorption of a quanta of ϕ with energy ω , momentum p along the effective string, and transverse momentum $\vec{p}$. We shall continue to use $+- - -$ signature, so for example $p \cdot x = \omega t - px - \vec{p} \cdot \vec{x}$. The absorption cross-section can be calculated by setting $\phi(t, x, \vec{x}=0) = e^{-ip \cdot x}$ and then treating (5.72) as a time-dependent perturbation to the CFT which describes the effective string in isolation. Stimulated emission would be calculated by choosing $e^{ip \cdot x}$ rather than $e^{-ip \cdot x}$. The t and x dependence of $\mathcal{O}(t, x)$ is fixed by the free theory:

$$\mathcal{O}(t, x) = e^{i\hat{p} \cdot x} \mathcal{O}(0, 0) e^{-i\hat{p} \cdot x} \quad (5.73)$$

where $\hat{p} \cdot x = Ht - Px$, H and P being the Hamiltonian and momentum operators of the CFT. If one considers the perturbation (5.72) to act for a time t , then Fermi's Golden Rule gives the thermally averaged transition probability as

$$\mathcal{P} = \sum_{i,f} \frac{e^{-\beta \cdot p_i}}{Z} P_{i \rightarrow f} = Lt \sum_{i,f} \frac{e^{-\beta \cdot p_i}}{Z} (2\pi)^2 \delta^2(p + p_i - p_f) |\langle f | \mathcal{O}(0, 0) | i \rangle|^2. \quad (5.74)$$

This formula is valid for when the length L of the effective string is much larger than the Compton wavelength of the incoming scalar. In (5.74), β has two components, $\beta^+ = \beta_L$ and $\beta^- = \beta_R$. The partition function splits into left and right sectors:

$$Z = \text{tr} e^{-\beta \cdot \hat{p}} = (\text{tr}_L e^{-\beta_L \hat{p}^+}) (\text{tr}_R e^{-\beta_R \hat{p}^-}). \quad (5.75)$$

For simplicity we take the momentum $p = (\omega, 0, \vec{p})$ of the incoming particle perpendicular to the brane, but clearly (5.74) remains valid for the case of particles with Kaluza-Klein charge.

The summation over final states can become tedious when the scalar turns into more than two excitations on the effective string. Already in the case of fixed scalars (chapter 3), which split into two right-movers and two left-movers, the evaluation of this summation was a nontrivial exercise. It therefore seems worthwhile to develop further a method employed in [100] in which the absorption probability is read off from the two-point function of the operator $\mathcal{O}$ in the effective string CFT.

Allow t to take on complex values, defining $\mathcal{O}(t, x)$ by (5.73) for arbitrary complex t . According to usual notational conventions [133], $\mathcal{O}^\dagger(t, x)$ is no longer the adjoint of $\mathcal{O}(t, x)$ except when t is real; instead, $\mathcal{O}^\dagger(t, x)$ is evolved from $\mathcal{O}^\dagger(0, 0)$ using (5.73). The conventional thermal Green's function takes $t = -i\tau$ where τ is the Euclidean time:

$$\mathcal{G}(-i\tau, x) = \langle \mathcal{O}^\dagger(-i\tau, x) \mathcal{O}(0, 0) \rangle = \text{tr} (\rho T_\tau \{ \mathcal{O}^\dagger(-i\tau, x) \mathcal{O}(0, 0) \}) \quad (5.76)$$

where $\rho = e^{-\beta \cdot \hat{p}}/Z$. One can continue to arbitrary complex t , defining T_τ to time-order with respect to $-\Im(t)$. The convenience of doing this is that the integral

$$\int d^2x e^{ip \cdot x} \mathcal{G}(t - i\epsilon, x) = \sum_{i,f} \frac{e^{-\beta \cdot p_i}}{Z} (2\pi)^2 \delta^2(p + p_i - p_f) |\langle f | \mathcal{O}(0, 0) | i \rangle|^2 \quad (5.77)$$

reproduces the right-hand side of (5.74). The proof of (5.77) proceeds by inserting $\sum_i |i\rangle\langle i|$ and $\sum_f |f\rangle\langle f|$ into (5.76) before $\mathcal{O}^\dagger$ and $\mathcal{O}$ respectively.

Now let us turn to the evaluation of $\mathcal{G}(t, x)$. Assume that $\mathcal{O}(t, x)$ has the form

$$\mathcal{O}(t, x) = \mathcal{O}_+(x^+) \mathcal{O}_-(x^-) \quad (5.78)$$

where $x^\pm = t \pm x$ and $\mathcal{O}_+$ and $\mathcal{O}_-$ are primary fields of dimensions h_L and h_R , respectively. Set $z = ix^-$ so that, for x real and $t = -i\tau$ imaginary, $\bar{z} = ix^+$. The singularities in $\mathcal{G}(t, x)$ are determined by the

OPE's of $\mathcal{O}_+$ and $\mathcal{O}_-$ with themselves:

$$\begin{aligned}\mathcal{O}_+(\bar{z})\mathcal{O}_+^\dagger(\bar{w}) &= \frac{C_{\mathcal{O}_+}}{(\bar{z}-\bar{w})^{2h_L}} + \text{less singular} \\ \mathcal{O}_-(z)\mathcal{O}_-^\dagger(w) &= \frac{C_{\mathcal{O}_-}}{(z-w)^{2h_R}} + \text{less singular}.\end{aligned}\tag{5.79}$$

$\mathcal{G}(t, x)$ factors into a left-moving and right-moving piece. The imaginary time periodicity properties of each piece, together with their singularities, suffice to fix the form of $\mathcal{G}(t, x)$ completely:⁴

$$\mathcal{G}(t, x) = \frac{C_{\mathcal{O}}}{i^{2h_L+2h_R}} \left(\frac{\pi T_L}{\sinh \pi T_L x^+} \right)^{2h_L} \left(\frac{\pi T_R}{\sinh \pi T_R x^-} \right)^{2h_R}\tag{5.80}$$

where $C_{\mathcal{O}} = C_{\mathcal{O}_+} C_{\mathcal{O}_-}$.

At nonzero temperature, the absorption cross-section cannot be calculated straight from (5.74): for bosons, the stimulated emission probability must be subtracted off in order to obtain a result consistent with detailed balance, as described in chapter 3. The net result is to set $\sigma_{\text{abs}} \mathcal{F} t = \mathcal{P}(1 - e^{-\beta \cdot p})$ where $\mathcal{F}$ is the flux and $\mathcal{P}$ is read off from (5.74). For fermions, the presence of an incoming wave inhibits by the Exclusion Principle emission processes leading to another fermion in the same state as the incoming wave. The absorption cross-section must therefore be calculated using $\sigma_{\text{abs}} \mathcal{F} t = \mathcal{P}(1 + e^{-\beta \cdot p})$. Again, this result is in accord with detailed balance.

The considerations of the previous paragraph can be restated compactly in terms of the Green's function:

$$\begin{aligned}\sigma_{\text{abs}} &= \frac{L}{\mathcal{F}} \int d^2 x e^{ip \cdot x} (\mathcal{G}(t - i\epsilon, x) - \mathcal{G}(t + i\epsilon, x)) \\ &= \frac{LC_{\mathcal{O}}}{\mathcal{F}} \frac{(2\pi T_L)^{2h_L-1} (2\pi T_R)^{2h_R-1}}{\Gamma(2h_L)\Gamma(2h_R)} \frac{e^{\beta \cdot p/2} - (-1)^{2h_L+2h_R} e^{-\beta \cdot p/2}}{2} \\ &\quad \cdot \left| \Gamma\left(h_L + i\frac{p_+}{2\pi T_L}\right) \Gamma\left(h_R + i\frac{p_-}{2\pi T_R}\right) \right|^2.\end{aligned}\tag{5.81}$$

One way to perform the integral in the first line is first to separate into x^+ and x^- factors and then to deform the contours in the separate factors by setting $\epsilon = \beta_L/2$ or $\beta_R/2$. Assuming that bosons and fermions couple, respectively, to conformal fields with $h_L + h_R$ an integer or half an odd integer, one indeed obtains the factor $1 \mp e^{-\beta \cdot p}$ required by detailed balance.

The formula (5.81) represents almost the most general functional form for an absorption cross-section that the effective string model is capable of predicting. One possible generalization is for the bulk field to couple to a sum of different operators $\mathcal{O}(t, x)$, in which case a sum of terms like (5.81) would be expected. Another generalization can arise from a coupling of a bulk field ϕ to the effective string not through its value $\phi(t, x, \vec{x}=0)$ on the string, as shown in (5.72), but rather through its derivatives: for instance $\partial_i \phi(t, x, \vec{x}=0)$ where i labels a transverse dimension. In case of fields without Kaluza-Klein charge, the effect of n such derivatives is simply to introduce an extra factor ω^{2n} on the right hand side of (5.81). Since the flux is $\mathcal{F} = \omega$ for a canonically normalized scalar, the ω dependence of the cross-section is

$$\sigma_{\text{abs}} \sim \omega^{2n-1} \sinh\left(\frac{\omega}{2T_H}\right) \left| \Gamma\left(h_L + i\frac{\omega}{4\pi T_L}\right) \Gamma\left(h_R + i\frac{\omega}{4\pi T_R}\right) \right|^2.\tag{5.82}$$

As we shall see in a specific example below, the flux factor for massless fermions cancels out a similar factor in $C_{\mathcal{O}}$. So for a fermionic field which couples to the effective string through a term in the

⁴Actually, there is a subtlety here: the information from periodicity and singularities must be supplemented by a sum rule [133] on the spectral density to squeeze out an ambiguity in the analytic continuation.

lagrangian of the form $\partial^n \psi \mathcal{O}$, the energy dependence of the cross-section is

$$\sigma_{\text{abs}} \sim \omega^{2n} \cosh\left(\frac{\omega}{2T_H}\right) \left| \Gamma\left(h_L + i\frac{\omega}{4\pi T_L}\right) \Gamma\left(h_R + i\frac{\omega}{4\pi T_R}\right) \right|^2. \quad (5.83)$$

The remarkable fact is that numerous classical absorption calculations that have appeared in the literature [7, 8, 73, 99, 87, 21, 103, 102, 110, 112, 101, 75, 100, 113, 114, 71] all give results consistent with (5.81) or (5.82) in the near-extremal limit. As an example, consider massless minimally coupled scalars falling into the four-dimensional black hole considered in [87], whose effective string model is derived from the picture of three intersecting sets of M5-branes. The absorption cross-section for the ℓ^{th} partial wave is

$$\sigma_{\text{abs}}^\ell = \frac{2}{\omega^2} \frac{(\omega T_H A_h)^{2\ell+1}}{(2\ell)!^2 (2\ell+1)!!} \sinh\left(\frac{\omega}{2T_H}\right) \left| \Gamma\left(\ell+1 + i\frac{\omega}{4\pi T_L}\right) \Gamma\left(\ell+1 + i\frac{\omega}{4\pi T_R}\right) \right|^2, \quad (5.84)$$

consistent with a coupling to the effective string of the form $(\partial^\ell \phi) \mathcal{O}$ where $\mathcal{O}$ has dimensions $h_L = h_R = \ell + 1$. Another interesting example is the fixed scalar [21, 96]. The cross-section for the s -wave in the four-dimensional case [96], with three charges equal and much greater than the fourth ($R = r_1 = r_2 = r_3 \gg r_K \sim r_0$) is

$$\sigma_{\text{abs}} = \frac{r_0^3}{2\omega R^2} \sinh\left(\frac{\omega}{2T_H}\right) \left| \Gamma\left(2 + i\frac{\omega}{4\pi T_L}\right) \Gamma\left(2 + i\frac{\omega}{4\pi T_R}\right) \right|^2, \quad (5.85)$$

consistent with a coupling of the form $\phi T_{++} T_{--}$. It is not hard to convince oneself that the analysis of the two-point function works the same when $\mathcal{O}(t, x) = T_{++}(x^+) T_{--}(x^-)$ as it did when $\mathcal{O}(t, x)$ was the product of left and right moving primary fields.

One naturally expects that an equal charge black hole whose metric is extreme Reissner-Nordstrom, like the $\mathcal{N} = 4$ example we focused on in sections 5.2, 5.4.2, and 5.5, can be obtained in the effective string picture by taking $T_L \gg T_R, \omega$. Indeed, it was found in [100] that ordinary scalar cross-sections in this metric, and even in the Kerr-Newman metric, have precisely the form one would expect from an effective string with $T_L \gg T_R, \omega$. The cross-sections have no dependence on T_L in this limit, and the authors of [100] suggested a model which made no reference to the left-moving sector. Note however that left-movers seem the natural explanation for the finite entropy of extremal black holes—a subject not addressed in [100]. In [25] it was argued in the context of $\mathcal{N} = 8$ compactifications that the CFT on the effective string is a $(0, 4)$ theory with central charges $c_L = c_R = 6$. The (local) $SU(2)$ R -symmetry of the right-moving sector was identified with the group $SO(3)$ of spatial rotations of the black hole. The same identification of a local $SU(2)$ on the effective string with $SO(3)$ was used in [100]. We will assume that an effective string description with 4 supersymmetries and $c = 6$ in the right-moving sector also applies to the equal charge black hole of $\mathcal{N} = 4$ supergravity.

Without committing to specific assumptions about the nature or existence of left-movers, one can conclude that the general form for an effective string absorption cross-section of massless particles is

$$\begin{aligned} \text{bosons:} \quad \sigma_{\text{abs}} &\sim \omega^{2n} \left| \frac{\Gamma(h_R + 2i\lambda)}{\Gamma(1 + 2i\lambda)} \right|^2 = \omega^{2n} \prod_{r=1}^{h_R-1} (r^2 + 4\lambda^2) \quad \text{if } h_R \in \mathbf{Z} \\ \text{fermions:} \quad \sigma_{\text{abs}} &\sim \omega^{2n} \left| \frac{\Gamma(h_R + 2i\lambda)}{\Gamma(\frac{1}{2} + 2i\lambda)} \right|^2 = \omega^{2n} \prod_{r=1/2}^{h_R-1/2} (r^2 + 4\lambda^2) \quad \text{if } h_R \in \mathbf{Z} + \frac{1}{2} \end{aligned} \quad (5.86)$$

where r runs over integers in the case of bosons and integers plus $1/2$ in the case of fermions, and

$$\lambda = \frac{\omega}{4\pi T} = \frac{\omega}{8\pi T_R}. \quad (5.87)$$

The cross-section (5.53) for minimally coupled photons fits the form (5.86) with $h_R = 2$ and $n = 1$. The form of the coupling of the minimal photon to the effective string is further constrained by rotation invariance. If we represent the right-moving sector using 4 free chiral bosons (which are neutral under $SU(2)$) and an $SU(2)$ doublet of fermions, then simplest coupling to the minimal photon with the right group theoretic properties is

$$\mathcal{L}_{\text{int}} = \phi_{\alpha\beta} \Psi_-^\alpha \partial_- \Psi_-^\beta F_+ + \text{h.c.} \quad (5.88)$$

where F_+ is a left-moving field which, as noted previously, does not affect the form of the absorption cross-section. Recall that $\phi_{\alpha\beta}$ is a field strength: one derivative is hidden inside it, so indeed $n = 1$. The indices on Ψ_-^α are for the group $SO(3)$ of spatial rotations, while the indices on $\phi_{\alpha\beta}$ are for the $SU(2)_L$ half of the Lorentz group $SO(3,1)$. But they can be contracted as shown in a static gauge description since the generators of $SO(3)$ are just sums of the generators of $SU(2)_L$ and $SU(2)_R$. With the current spinor index conventions, an upper dotted index is equivalent to a lower undotted index if only spatial $SO(3)$ rotations are considered.

The dilaton and axion cross-sections (5.54) fit the form (5.86) with $h_R = 2$, and so does the minimal fermion cross-section with $h_R = 3/2$. The natural guess is a coupling of these fields to the stress-energy tensor and the supercurrents: to linear order in all the fields,

$$\mathcal{L}_{\text{int}} = [(\phi + iB)T_{--} + \Lambda_{3\alpha} T_{F-}^\alpha - i\Lambda_{4\alpha} T_{F-}^{\dagger\alpha}] F_+ + \text{h.c.} \quad (5.89)$$

The form of (5.89) is dictated by the quarter of the $\mathcal{N} = 4$ supersymmetry which is preserved by the extreme black hole. The terms in the supersymmetry variation of $\mathcal{L}_{\text{int}}$ with no derivatives can be shown to cancel using

$$\begin{aligned} \delta(\phi + iB) &= \epsilon^\alpha \Lambda_{3\alpha} + \bar{\epsilon}_{\dot{\alpha}} i\sqrt{2}\sigma^{0\dot{\alpha}\beta} \Lambda_{4\beta} \\ \delta T_{F-}^\alpha &= \epsilon^\alpha T_{--} . \end{aligned} \quad (5.90)$$

Here ϵ_α parametrizes what was referred to in [59] as the ϵ_+^{34} supersymmetry. Note that in the conventions outlined in the appendix, both $\sqrt{2}\sigma^{0\dot{\alpha}\beta}$ and $\sqrt{2}\sigma_{\alpha\dot{\beta}}^0$ are numerically the identity matrix. The left-moving sector of the CFT is assumed to be neutral under supersymmetry.

The equivalence (in static gauge) of upper undotted and lower dotted indices has been used to simplify (5.89). The matrices $\sqrt{2}\sigma^{0\dot{\alpha}\beta}$ and $\sqrt{2}\sigma_{\alpha\dot{\beta}}^0$ can be used to convert between them. The index on $T_{F-}^\dagger$ has been raised in (5.89) using $\epsilon^{\alpha\beta}$. A consequence of the second line in (5.90) is thus $\delta T_{F-}^{\dagger\alpha} = -\bar{\epsilon}_{\dot{\alpha}} T_{--}$.

The normalization of T_{F-}^α used here differs from [134] by a factor of $\sqrt{2}$: if G_n^α and L_n are the supercurrent and Virasoro generators of [134], then the present conventions are to set $T_{F-}^\alpha(z) = \frac{1}{\sqrt{2}} \sum_n z^{-n-3/2} G_n^\alpha$ and $T(z) = \sum_n z^{-n-2} L_n$. On the complex plane, the nonzero two-point functions are

$$\begin{aligned} \langle T_{F-}^\alpha(z) T_{F-\beta}^\dagger(w) \rangle &= \frac{c}{3} \frac{\delta_\beta^\alpha}{(z-w)^3} \\ \langle T_{--}(z) T_{--}(w) \rangle &= \frac{c/2}{(z-w)^4} . \end{aligned} \quad (5.91)$$

Let us also assume that the only nonzero two-point function of F_+ and $F_+^\dagger$ is

$$\langle F(\bar{z}) F^\dagger(\bar{w}) \rangle = \frac{C_F}{(\bar{z} - \bar{w})^{2h_R}} . \quad (5.92)$$

Now we are ready to compute effective string cross-sections. For the dilaton, the operator $\mathcal{O}(t, x)$ entering into the analysis of (5.72)-(5.81) is

$$\mathcal{O}(t, x) = T_{--}(x^-) (F_+(x^+) + F_+^\dagger(x^+)) . \quad (5.93)$$

The fact that T_{--} is not primary does not alter the periodicity properties of its two-point function. Thus the arguments leading from (5.78) to (5.81) still apply, and $C_{\mathcal{O}} = cC_F = 6C_F$. Because of the non-canonical normalization of the dilaton field in the action (5.3), the particle flux in a wave $\phi = e^{-ip \cdot x}$ is $\mathcal{F} = 4\omega$, four times the usual value. Plugging these numbers into (5.81) and taking the large T_L limit, one obtains

$$\sigma_{\text{abs}} = \frac{LC_F}{8T} (2\pi T_L)^{2h_L-1} (2\pi T_R)^3 \frac{\Gamma(h_L)^2}{\Gamma(2h_L)} (1 + 4\lambda^2) . \quad (5.94)$$

The axion of course yields the same result.

The minimal fermions clearly have the same cross-section, so let us consider only Λ_3 . Let the incoming wave be $\Lambda_{3\alpha} = u_\alpha e^{-ip \cdot x}$. With $\mathcal{O}(t, x) = u_\alpha T_{F-}^\alpha(x^-) F_+(x^+)$, the analysis leading to (5.81) goes through as usual, yielding

$$C_{\mathcal{O}} = \bar{u}_{\dot{\alpha}} \delta_{\alpha}^{\beta} u_{\beta} \frac{cC_F}{3} = \bar{u}_{\dot{\alpha}} \sqrt{2} \sigma^{0\dot{\alpha}\beta} u_{\beta} 2C_F . \quad (5.95)$$

In the second equality the equivalence of lower undotted and upper dotted indices has again been used. Recall that $\sqrt{2} \sigma^{0\dot{\alpha}\beta}$ is indeed the identity matrix. The flux is $\mathcal{F} = 4\bar{u}_{\dot{\alpha}} \sqrt{2} \sigma^{0\dot{\alpha}\beta} u_{\beta}$. (As for the dilaton, the 4 here is due to the non-canonical normalization of the fermion field: see the footnote at the beginning of section 5.5.2.) Now (5.81) can be used again to give

$$\sigma_{\text{abs}} = \frac{\pi LC_F}{16} (2\pi T_L)^{2h_L-1} (2\pi T_R)^2 \frac{\Gamma(h_L)^2}{\Gamma(2h_L)} (1 + 16\lambda^2) . \quad (5.96)$$

The effective string cross-sections (5.94) and (5.96) stand in the same ratio as the semi-classical cross-sections for fixed scalars and minimal fermions quoted in (5.54).

Of course, it would be highly desirable to carry out calculations similar to the ones presented here for the black holes in five dimensions that can be modeled using the D1-brane D5-brane bound state. There one can hope that an understanding of the soliton picture can fix overall normalizations; but again one expects the residual supersymmetry to fix relative normalizations between fermionic and bosonic cross-sections.

5.7 Conclusion

One of the main technical results of this chapter has been to show that when the equations of motion for photons or fermions are simple enough to be analyzed by the dyadic index methods of [120, 121, 122], they lead to ordinary differential equations whose near-horizon form is hypergeometric. That fact alone gives their low-energy absorption cross-sections a form which is capable of explanation in the effective string description.

Dyadic index methods are not essential to the analysis of minimally coupled fermions; indeed, the Weyl equation for fermions has been analyzed recently in [71] in arbitrary dimensions using more conventional techniques. However, the dyadic index method provides an efficient, unified treatment of minimal fermions and minimal photons. Indeed, the photon radial equations (5.43) and (5.48) seem difficult to derive by other means. The existence of minimally coupled photons seems to depend essentially on the equal charge condition, which makes the dilaton background constant. Minimally coupled fermions, as suggested in section 5.5.2, may be more common because their existence depends on the vanishing of their supersymmetry variations in the black hole background.

The greybody factors computed in (5.53) and (5.54) are polynomials in the energy ω rather than quotients of gamma functions as found in [75]. This is characteristic of an effective string whose left-moving temperature T_L is much greater than T_R and ω . These polynomial greybody factors are

sufficient to determine the conformal dimension of the right-moving factor in the operator through which a field couples to the effective string, and the number of derivatives in that coupling. For example, the minimal photon couples through its field strength times a $h_R = 2$ operator. But (5.53) and (5.54) do not yield any information regarding the left-movers. To see the effects of left-movers, one might try to generalize the present treatment to black holes far from extremality, as was done recently in [114] for minimally coupled scalars.

The absence of a string soliton description of the equal charge black hole in pure $d = 4$, $\mathcal{N} = 4$ supergravity precludes a precise comparison of cross-sections between the effective string and semi-classical descriptions. However, by assuming that the effective string world-sheet theory is a $(0, 4)$ super-conformal field theory whose right-moving R -symmetry group, $SU(2)$, is identified with the group of spatial rotations, it has been possible to show that the relative normalizations of the dilaton, axion, and minimally coupled fermion cross-sections are correctly predicted by the effective string. The proposed couplings of these fields to the effective string are simple: the scalars couple to the stress-energy tensor while the fermions couple to the supercurrent. One would expect that it possible to extend this picture to a manifestly supersymmetric specification of how all the massless bulk fields couple to the effective string at linear order.

There is a simple point which nevertheless is worth emphasizing: the cross-sections of the dilaton, axion, and minimal fermions are related by supersymmetry despite their different energy dependence. The energy dependence (also known as the greybody factor) arises from finite-temperature kinematics. Unsurprisingly, the kinematic factors are different for particles of different spin; but their form turns out to be fixed by the conformal dimension of the field by which a field couples to the effective string. Supersymmetry acts on the S -matrix, relating the coefficients we have called $C_{\mathcal{O}}$ in section 5.6. The predictions of supersymmetry regarding the absorption cross-sections of different particles in the same multiplet thus have more to do with the relative normalization than the energy dependence.

Appendix: Dyadic index conventions

This appendix presents in a pedestrian fashion the aspects of the Newman-Penrose formalism relevant to the rest of the chapter. A readable introduction can be found in [135]; for an authoritative treatment the reader is referred to [124].

Sign conventions vary by author, and the ones used here are as close as possible to those of the original paper by Newman and Penrose [123] and to those of Teukolsky [120, 121]. First consider flat Minkowski spacetime with mostly minus metric, $\eta_{ab} = \text{diag}(1, -1, -1, -1)$. The conventions on raising and lowering spinor indices are those of “northwest contraction:”

$$\begin{aligned}\psi^\alpha &= \epsilon^{\alpha\beta} \psi_\beta & \psi_\alpha &= \psi^\beta \epsilon_{\beta\alpha} \\ \bar{\psi}^{\dot{\alpha}} &= \epsilon^{\dot{\alpha}\dot{\beta}} \bar{\psi}_{\dot{\beta}} & \bar{\psi}_{\dot{\alpha}} &= \bar{\psi}^{\dot{\beta}} \epsilon_{\dot{\beta}\dot{\alpha}}\end{aligned}\tag{5.97}$$

where the sign of the antisymmetric tensors is fixed by $\epsilon_{01} = \epsilon^{01} = \epsilon_{\dot{0}\dot{1}} = \epsilon^{\dot{0}\dot{1}} = 1$. In flat space, the conventional choice of the matrices $\sigma^a_{\alpha\dot{\beta}}$ which map bispinors to vectors is

$$\sigma^a = \frac{1}{\sqrt{2}} (1, \tau_3, \tau_1, -\tau_2)\tag{5.98}$$

where the matrices τ_i are the standard Pauli matrices. Vector indices are interchanged with pairs of spinor indices using the formulae

$$v^a = \sigma^a_{\alpha\dot{\alpha}} v^{\alpha\dot{\alpha}} \quad v^{\alpha\dot{\alpha}} = \sigma_a^{\alpha\dot{\alpha}} v_a\tag{5.99}$$

where $\sigma_a^{\alpha\dot{\alpha}} = \eta_{ab} \epsilon^{\alpha\beta} \epsilon^{\dot{\alpha}\dot{\beta}} \sigma^b_{\beta\dot{\beta}}$, consistent with our northwest contraction rules. There is no need to define matrices $\bar{\sigma}^{a\dot{\alpha}\beta}$. The metric has a simple form when written with spinor indices:

$$\eta_{ab} \sigma^a_{\alpha\dot{\alpha}} \sigma^b_{\beta\dot{\beta}} = \epsilon_{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}}.\tag{5.100}$$

In curved spacetime, the metric $g_{\mu\nu}$ is again chosen with $+---$ signature. In this chapter, the metric is always of the form

$$ds^2 = e^{2A(r)} dt^2 - e^{2B(r)} dr^2 - e^{2C(r)} (d\theta^2 + \sin^2 \theta d\phi^2) .\tag{5.101}$$

It will turn out to be useful to define not only the standard diagonal vierbein,

$$e^a_\mu = \text{diag}(\sqrt{g_{tt}}, \sqrt{-g_{rr}}, \sqrt{-g_{\theta\theta}}, \sqrt{-g_{\phi\phi}}) ,$$

but also a complex null tetrad⁵

$$\begin{aligned}\ell^\mu &= \frac{e^\mu_t + e^\mu_r}{\sqrt{2}} & n^\mu &= \frac{e^\mu_t - e^\mu_r}{\sqrt{2}} \\ m^\mu &= \frac{e^\mu_\theta + ie^\mu_\phi}{\sqrt{2}} & \bar{m}^\mu &= \frac{e^\mu_\theta - ie^\mu_\phi}{\sqrt{2}} .\end{aligned}\tag{5.102}$$

One of the conveniences of working with spinors is that a spinor is a sort of square root of a null vector: for any spinor ψ^α , $v^\mu = e^\mu_a \sigma^a_{\alpha\dot{\alpha}} \psi^\alpha \bar{\psi}^{\dot{\alpha}}$ is a null vector, and any null vector can be written in this form. It is important to note that in the context of the Newman-Penrose formalism, spinor components are

⁵In the literature it is common to see factors of g_{tt} included in the definitions of ℓ^μ and n^μ so that seven rather than six of the spin coefficients vanish. This however complicates the time-reversal properties of the solutions.

ordinary commuting numbers, not Grassmann numbers. It is possible to introduce a basis (o^α, ι^α) for spinor space with the properties

$$\begin{aligned} o_\alpha \iota^\alpha &= 1 \\ \ell^{\alpha\dot{\alpha}} &= o^\alpha \bar{o}^{\dot{\alpha}} \quad n^{\alpha\dot{\alpha}} = \iota^\alpha \bar{\iota}^{\dot{\alpha}} \quad m^{\alpha\dot{\alpha}} = o^\alpha \bar{\iota}^{\dot{\alpha}} \quad \bar{m}^{\alpha\dot{\alpha}} = \iota^\alpha \bar{o}^{\dot{\alpha}} . \end{aligned} \quad (5.103)$$

A particular choice of (o^α, ι^α) is

$$o^\alpha = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \iota^\alpha = \begin{pmatrix} 0 \\ 1 \end{pmatrix} . \quad (5.104)$$

Dyadic indices are introduced by defining $\xi_0^\alpha = o^\alpha$, $\xi_1^\alpha = \iota^\alpha$ and writing ψ_Γ for the components of the spinor ψ_α with respect to the basis ξ_Γ^α :

$$\psi_\Gamma = \xi_\Gamma^\alpha \psi_\alpha \quad \psi_\alpha = -\xi_\alpha^\Gamma \psi_\Gamma . \quad (5.105)$$

The minus sign in the second equation is the result of insisting on the same raising and lowering conventions for dyadic indices as for spinor indices: $\xi_\alpha^\Gamma = \epsilon^{\Gamma\Delta} \xi_\Delta^\beta \epsilon_{\beta\alpha}$.

It is a familiar story [28] how the minimal $SO(3, 1)$ connection $\omega_\mu{}^a{}_b$ on the local Lorentz bundle is induced from the Christoffel connection: one defines

$$\omega_\mu{}^a{}_b = e_\nu^a \partial_\mu e_\nu^b + e_\nu^a \Gamma_{\mu\rho}^\nu e_b^\rho \quad (5.106)$$

so that

$$\nabla_\mu v^a = \partial_\mu v^a + \omega_\mu{}^a{}_b v^b = e_\nu^a \nabla_\mu v^\nu = e_\nu^a (\partial_\mu v^\nu + \Gamma_{\mu\rho}^\nu v^\rho) . \quad (5.107)$$

The Newman-Penrose spin coefficients are defined in an exactly analogous way. In fact, they are merely special (complex) linear combinations of the $\omega_\mu{}^a{}_b$. It is conventional in the literature to make dyadic indices “neutral” under the covariant derivative ∇_μ : $\nabla_\mu v_\Gamma = \partial_\mu v_\Gamma$. The covariant derivatives of spinors, by contrast, are defined using the connection induced from $\omega_\mu{}^a{}_b$. It is convenient to define a “completely covariant” derivative D_μ and a connection $\gamma_\mu{}^\Sigma{}_\Gamma$ with the defining properties

$$D_\mu \psi_\Gamma = \partial_\mu \psi_\Gamma - \psi_\Sigma \gamma_\mu{}^\Sigma{}_\Gamma = \xi_\Gamma^\alpha \nabla_\mu \psi_\alpha . \quad (5.108)$$

A brief way of characterizing the covariant derivative is to say that under ∇_μ , the quantities $g_{\mu\nu}$, η_{ab} , $\epsilon_{\alpha\beta}$, e_μ^a , and $\sigma_{\alpha\dot{\alpha}}^a$ (together with their alternative incarnations η^{ab} , $\epsilon^{\dot{\alpha}\dot{\beta}}$, etc.) are covariantly constant. Under D_μ , the quantities ξ_Γ^α are covariantly constant as well. From (5.108) it is immediate that⁶

$$\gamma_{m\Sigma\Gamma} = -\xi_{\Gamma\alpha} \nabla_\mu \xi_\Sigma^\alpha . \quad (5.109)$$

The quantities $\gamma_{\Delta\dot{\Delta}\Gamma\Sigma} = \sigma_{\Delta\dot{\Delta}}^\mu \gamma_{m\Gamma\Sigma}$ are the spin coefficients. They have the symmetry $\gamma_{\Delta\dot{\Delta}\Gamma\Sigma} = \gamma_{\Delta\dot{\Delta}\Sigma\Gamma}$. With twelve independent complex components they represent the same information as the forty real components of the Christoffel connection $\Gamma_{\nu\rho}^\mu$. It is useful to note that

$$\sigma_{\Delta\dot{\Delta}}^\mu = e_a^\mu \xi_{\Delta\dot{\Delta}}^\alpha \bar{\xi}_{\dot{\Delta}}^{\dot{\alpha}} \sigma_{\alpha\dot{\alpha}}^a = \begin{pmatrix} \ell^\mu & m^\mu \\ \bar{m}^\mu & n^\mu \end{pmatrix} . \quad (5.110)$$

A useful formula for calculating the spin coefficients can be given in terms of $\sigma_{\Delta\dot{\Delta}}^\mu$:

$$\gamma_{\Delta\dot{\Delta}\Gamma\Sigma} = -\frac{1}{2} \xi_\Gamma^\beta \bar{\xi}_{\dot{\Gamma}}^{\dot{\beta}} \sigma_{\Delta\dot{\Delta}}^\nu \nabla_\nu (\xi_{\Sigma\beta} \bar{\xi}_{\dot{\Sigma}}^{\dot{\beta}}) = -\frac{1}{2} \sigma_{\Gamma\dot{\Gamma}}^\mu \sigma_{\Delta\dot{\Delta}}^\nu \nabla_\nu \sigma_{\mu\Sigma}^{\dot{\Gamma}} . \quad (5.111)$$

⁶Note that the choice of sign for $\gamma_{\Delta\dot{\Delta}\Gamma\Sigma}$ follows the convention of [120] and [123] rather than of [135].

Some further notational definitions are conventional in dyadic index papers:

$$\begin{aligned}\gamma_{00r\Sigma} &= \begin{pmatrix} \kappa & \epsilon \\ \epsilon & \pi \end{pmatrix} & \gamma_{0ir\Sigma} &= \begin{pmatrix} \sigma & \beta \\ \beta & \mu \end{pmatrix} \\ \gamma_{10r\Sigma} &= \begin{pmatrix} \rho & \alpha \\ \alpha & \lambda \end{pmatrix} & \gamma_{1ir\Sigma} &= \begin{pmatrix} \tau & \gamma \\ \gamma & \nu \end{pmatrix}\end{aligned}\tag{5.112}$$

$$D = \ell^\mu \nabla_\mu \quad \Delta = n^\mu \nabla_\mu \quad \delta = m^\mu \nabla_\mu \quad \bar{\delta} = \bar{m}^\mu \nabla_\mu .\tag{5.113}$$

For the metric (5.101), one finds

$$\begin{aligned}\kappa &= \pi = \sigma = \lambda = \tau = \nu = 0 \\ \epsilon = \gamma &= \frac{e^{-B} A'}{2\sqrt{2}} \quad \beta = -\alpha = \frac{e^{-C} \cot \theta}{2\sqrt{2}} \quad \mu = \rho = -\frac{e^{-B} C'}{\sqrt{2}}\end{aligned}\tag{5.114}$$

where primes denote derivatives with respect to r . (5.114) represents a remarkably economical way of describing the connection of an arbitrary spherically symmetric spacetime: there are only three independent nonzero spin coefficients, and they are real.

Chapter 6

Absorption by non-dilatonic branes

6.1 Introduction

Dirichlet branes provide an elegant embedding of Ramond-Ramond charged objects into string theory [4]. The D-brane description of the dynamics of these solitons may be compared with corresponding results in the semi-classical low-energy supergravity. In particular, a counting of degeneracies for certain intersecting D-branes reproduces, in the limit of large charges, the Bekenstein-Hawking entropy of the corresponding geometries [5, 6]. Furthermore, calculations of emission and absorption rates for scalar particles agree with a simple “effective string” model for the dynamics of the intersection [6, 7, 8, 73, 99, 75, 87, 21, 96]. Questions have been raised, however, whether the simplest model is capable of incorporating all the complexities of black holes physics [112, 102, 103]. Indeed, it is possible that the dynamics of intersecting D-branes, which is not yet fully understood, cannot be captured by one simple model.

This chapter, based on the paper [10], is concerned with the simpler configurations which involve parallel D-branes only. Their string theoretic description is well understood in terms of supersymmetric $U(N)$ gauge theory on the world-volume [13]. Perturbative string calculations of scattering [74, 136, 137] and absorption [138] are fairly straightforward for the parallel D-branes, and their low-energy dynamics is summarized in the DBI action.

To leading order in the string coupling, N coincident Dp -branes are described by $O(N^2)$ free fields in $p + 1$ dimensions. This result may be compared with the Bekenstein-Hawking entropy of the near-extremal p -brane solutions in supergravity. In [45, 46] it was found the the scaling of the Bekenstein-Hawking entropy with the temperature agrees with that for a massless gas in p dimensions only for the non-dilatonic p -branes. The only representative of this class which is described by parallel D-branes is the self-dual three-brane. Other representatives include the dyonic string in $D = 6$, and the two-branes and five-branes of M-theory. In [47] a way of reconciling the differing scalings for the dilatonic branes [139] was proposed.¹ Nevertheless, in [9] it was shown that the non-dilatonic branes (and especially the self-dual three-brane) have a number of special properties that allow for more detailed comparisons between semi-classical gravity and the microscopic theory. For example, the string theoretic calculation of the absorption cross-section by three-branes for low-energy dilatons was found to agree *exactly* with the classical calculation in the background of the extremal classical geometry [9].

What are the features that make the three-brane so special? In classical supergravity, the extremal three-brane is the only RR-charged solution that is perfectly non-singular [141, 142, 143]. For N

¹Other ideas on how to find agreement for the dilatonic branes were put forward in [140].

three-branes, the curvature of the classical solution is bounded by a quantity of order

$$\frac{1}{\sqrt{N\kappa_{10}}} \sim \frac{1}{\alpha' \sqrt{Ng_{\text{str}}}} .$$

Thus, to suppress the string scale corrections to the classical metric, we need to take the limit $Ng_{\text{str}} \rightarrow \infty$. This fact seems to lead to a strongly coupled theory on the world-volume and raises questions about the applicability of string perturbation theory to macroscopic three-branes. However, in [9] it was shown that the dimensionless expansion parameter that enters the string theoretic calculation of the absorption cross-section is actually

$$N\kappa_{10}\omega^4 \sim Ng_{\text{str}}\alpha'^2\omega^4 , \quad (6.1)$$

where ω is the incident energy. Thus, we may consider a “double scaling limit,”

$$Ng_{\text{str}} \rightarrow \infty , \quad \omega^2\alpha' \rightarrow 0 , \quad (6.2)$$

where the expansion parameter (6.1) is kept small. Moreover, the classical absorption cross-section is naturally expanded in powers of $\omega^4 \times (\text{curvature})^{-2}$, which is the same expansion parameter (6.1) as the one governing the string theoretic description of the three-branes. The two expansions of the cross-section thus may indeed be compared, and the leading term agrees exactly [9]. This provides strong evidence in favor of absorption by extremal three-branes being a unitary process. While in the classical calculation the information carried by the dilaton seems to disappear down the infinite throat of the classical solution, the stringy approach indicates that the information is not lost: it is stored in the quantum state of the back-to-back massless gauge bosons on the world-volume which are produced by the dilaton. Subsequent decay of the three-brane back to the ground state proceeds via annihilation of the gauge bosons into an outgoing massless state, and there seems to be no space for information loss.

The scenario mentioned above certainly deserves a more careful scrutiny. In this chapter, we carry out further comparisons between string theory and classical gravity of the three-brane. In string theory there is a variety of fields that act as minimally coupled massless scalars with respect to the transverse dimensions. The dilaton, which was investigated in [9], is perhaps the simplest one to study. In this chapter we turn to other such fields: the RR scalar and the gravitons polarized parallel to the world-volume. We show that in classical gravity all these fields satisfy the same equation and, therefore, are absorbed with the same rate as the dilaton. However, their cubic couplings to the massless world-volume modes dictated by string theory [138] are completely different. In particular, the longitudinally polarized gravitons can turn into pairs of gauge bosons, scalars, or fermions. Adding up these rates we find that the total cross-section has the same universal value, in agreement with classical gravity.

Another interesting check concerns the absorption of scalars in partial waves higher than $\ell = 0$. In [9] it was shown that the scaling of the relevant cross-sections with ω and N agrees for all ℓ . For $\ell = 1$ we show that the coefficient agrees as well. For $\ell > 1$ the simplest assumption about the effective action does not yield a coefficient which agrees with the classical calculation. Normalization of the effective action is a subtle matter, however, and we suspect that its direct determination from string amplitudes will yield agreement with the classical cross-sections.

While a microscopic description of RR-charged branes is by now well known in string theory, the situation is not as simple for the p -brane solutions of the 11-dimensional supergravity. There is good evidence that a yet unknown M-theory underlies their fundamental description. Comparisons with semi-classical gravity provide consistency checks on this description. In [9] the scalings of the classical absorption cross-sections were found to agree with the two-brane and five-brane effective action considerations. Here we calculate the absorption cross-section of a longitudinally polarized graviton by a single five-brane. Using the world-volume effective action, we find that the absorption

cross-section is $1/4$ of the rate formally predicted by classical gravity. In fact, one could hardly expect perfect agreement for a single five-brane – the classical description is expected to be valid only for a large number of coincident branes. It is interesting, nevertheless, how close the two calculations come to agreeing with each other. We also carry out a similar comparison for a single two-brane, but find the discrepancy in the coefficient to be far greater than in the five-brane case.

The structure of this chapter is as follows. In section 6.2 we exhibit the spacetime effective actions and derive the classical equations satisfied by various fields. In section 6.3 we present the new three-brane calculations. In section 6.4 we carry out the comparisons of classical 11-dimensional supergravity with the predictions of the M-brane world-volume actions. We conclude in section 6.5.

6.2 Spacetime effective actions and perturbations around non-dilatonic p -brane solutions

6.2.1 Perturbations of three-branes

The bosonic part of the field equations of $d = 10$ type IIB supergravity, which is the low-energy limit of the type IIB superstring [144], can be derived from the following action: [81, 145]

$$S_{10} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \left[\sqrt{-G} (e^{-2\phi} [R + 4(\partial\phi)^2 - \frac{1}{12}(\partial B_2)^2] - \frac{1}{2}(\partial C)^2 - \frac{1}{12}(\partial C_2 - C\partial B_2)^2 - \frac{1}{4\cdot 5!}F_5^2) - \frac{1}{2\cdot 4! \cdot (3!)^2} \epsilon_{10} C_4 \partial C_2 \partial B_2 + \dots \right], \quad (6.3)$$

where²

$$(\partial B_2)_{MNK} \equiv 3\partial_{[M} B_{NK]}, \quad (\partial C_4)_{MNKLP} \equiv 5\partial_{[M} C_{NKL P]}, \\ F_5 = \partial C_4 + 5(B_2 \partial C_2 - C_2 \partial B_2).$$

Following [145] we assume that the self-duality constraint $F_5 = \tilde{F}_5$ may be added at the level of the equations of motion. This action is a useful tool for deriving the dimensionally reduced forms of type IIB supergravity action [145] and also for discussing perturbations near the solitonic p -brane solutions given below. Written in the Einstein frame ($g_{MN} = e^{-\phi/2} G_{MN}$) it takes the following $SL(2, R)$ covariant form:

$$S_{10} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \left(\sqrt{-g_{10}} \left[R - \frac{1}{2}(\partial\phi)^2 - \frac{1}{12}e^{-\phi}(\partial B_2)^2 - \frac{1}{2}e^{2\phi}(\partial C)^2 - \frac{1}{12}e^{\phi}(\partial C_2 - C\partial B_2)^2 - \frac{1}{4\cdot 5!}F_5^2 \right] - \frac{1}{2\cdot 4! \cdot (3!)^2} \epsilon_{10} C_4 \partial C_2 \partial B_2 + \dots \right). \quad (6.4)$$

The extremal three-brane of type IIB theory is represented by the background [146, 141]

$$ds_{10}^2 = H^{-1/2}(-dt^2 + dx_a dx_a) + H^{1/2} dx_i dx_i, \quad (6.5)$$

$$(\partial C_4)_{abcdi} = \epsilon_{abcd} \partial_i H^{-1}, \quad (\partial C_4)_{ijklm} = \epsilon_{ijklmn} \partial_n H, \quad H = 1 + \frac{R^4}{r^4}, \quad r^2 = x_i x_i, \quad (6.6)$$

²We use the following notation. The signature is $(- + \dots +)$. $M, N, \dots$ label the coordinate indices of $d = 10$ or $d = 11$ theory. The indices $\alpha, \beta, \dots (a, b, \dots)$ will label the spacetime (spatial) coordinates parallel to the p -brane world-volume, i.e. $\alpha = (0, a)$, $a = 1, \dots, p$; the indices $i, j, \dots$, will label the coordinates transverse to the p -brane, $i = p+1, \dots, 9(10)$. We shall also use $\mu, \nu, \dots$ for the indices of the dimensionally reduced theory obtained by compactifying the internal directions parallel to the p -brane, i.e. $\mu = (0, i)$. ϵ_d will stand for the totally antisymmetric symbol (density). Contractions over repeated lower-case indices are always performed with the flat metric, and $A_{[ab]} \equiv \frac{1}{2}(A_{ab} - A_{ba})$.

with all other components and fields vanishing (i.e. $\phi = C = B_2 = C_2 = 0$, $F_5 = \partial C_4$). The scalar curvature $\sim (\partial C_4)^2 \sim R^{-2}$ near $r = 0$. In fact, this metric may be extended to a geodesically complete non-singular geometry [142, 143]. This is the only RR-charged p -brane for which this is possible.

Let us consider small perturbations near this background which depend only on x_μ , i.e. on the time t and the transverse coordinates x_i . Our aim is to identify the modes which have simple Klein-Gordon type equations. A guiding principle is to look for (components of) the fields that have trivial background values.

The obvious examples are the dilaton ϕ and the Ramond-Ramond scalar C perturbations which are decoupled from the C_4 -background and thus have the action ($\varphi = (\phi, C)$)

$$S_{scal.} = -\frac{1}{4\kappa_{10}^2} \int d^{10}x \sqrt{-g_{10}} g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi = -\frac{1}{4\kappa_{10}^2} \int d^{10}x [\partial_i \varphi \partial_i \varphi - H(r) \partial_0 \varphi \partial_0 \varphi], \quad (6.7)$$

where we have used the fact that for the three-brane background the Einstein and the string frame metrics are identical, $G_{MN} = g_{MN}$, which implies

$$g_{\mu\nu} = \text{diag}(-H^{-1/2}, H^{1/2} \delta_{ij}), \quad \sqrt{-g_{10}} = H^{1/2}.$$

Perturbations of the metric are, in general, mixed with perturbations of components of C_4 . An important exception is the traceless part of the longitudinal (polarized along the three-brane) graviton perturbations h_{ab} . These perturbations give rise to scalars upon dimensional reduction to $d = 7$. Let us split the metric in the “7+3” fashion

$$ds_{10E}^2 = g_{\mu\nu} dx^\mu dx^\nu + g_{ab} dx^a dx^b,$$

and assume that all the fields depend only on x^μ (this is equivalent to reduction to 7 dimensions). Then the relevant part of (6.4) becomes

$$S_{10} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \sqrt{-g_7} \sqrt{g_3} (R_7 - \frac{1}{4} g^{ab} g^{cd} \partial_\mu g_{ac} \partial^\mu g_{bd} - \frac{1}{48} F_{\mu\nu abc} F^{\mu\nu}_{def} g^{ad} g^{be} g^{cf} + \dots), \quad (6.8)$$

where $g_7 \equiv \det g_{\mu\nu}$, $g_3 \equiv \det g_{ab}$ and we have written down explicitly the only F_5^2 term that could potentially couple h_{ab} to the gauge field strength background. Introducing the “normalized” metric $\gamma_{ab} = g_3^{-1/3} g_{ab}$ which has unit determinant, we find

$$S_{scal.grav.} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \sqrt{-g_7} \sqrt{g_3} (R_7 - \frac{1}{4} \gamma^{ab} \gamma^{cd} \partial_\mu \gamma_{ac} \partial^\mu \gamma_{bd} - \frac{1}{12} g_3^{-2} \partial_\mu g_3 \partial^\mu g_3 - \frac{1}{8} g_3^{-1} F_{\mu\nu} F^{\mu\nu} + \dots), \quad (6.9)$$

where $F_{\mu\nu} = F_{\mu\nu 123}$ is the $d = 7$ vector field strength which, according to (6.6), describes an electrically charged extremal black hole. Thus, only the determinant of g_{ab} , which is related to the 7-dimensional dilaton, couples to the gauge field background. This is consistent with the fact that γ_{ab} has a trivial background value, $\gamma_{ab} = \delta_{ab}$, while the value of $\det g_{ab}$ in (6.5) is $g_3 = H^{-3/2}$.

The fluctuations $h_{ab} = \gamma_{ab} - \delta_{ab}$ (which are traceless, $h_{aa} = 0$, to keep $\det \gamma_{ab} = 1$) thus have the same quadratic part of the action as the scalar fields in (6.7)

$$S_{scal.grav.} = -\frac{1}{8\kappa_{10}^2} \int d^{10}x \sqrt{-g_{10}} \partial_\mu h_{ab} \partial^\mu h_{ab} = -\frac{1}{8\kappa_{10}^2} \int d^{10}x [\partial_i h_{ab} \partial_i h_{ab} - H(r) \partial_0 h_{ab} \partial_0 h_{ab}]. \quad (6.10)$$

Similar conclusions hold for other p -brane solutions because the “normalized” internal space metric γ_{ab} has a flat background value.

6.2.2 Perturbations of two-branes and five-branes

The discussion of perturbations around the two-brane and five-brane solutions [147, 148] of the $d = 11$ supergravity is very similar to the analysis in the preceding subsection. The starting point is the bosonic part of the $d = 11$ supergravity action [149]

$$S_{11} = \frac{1}{2\kappa_{11}^2} \int d^{11}x \left(\sqrt{-g_{11}} \left[R - \frac{1}{2 \cdot 4!} (\partial C_3)^2 \right] + \frac{1}{(12)^4} \epsilon_{11} C_3 \partial C_3 \partial C_3 + \dots \right), \quad (6.11)$$

where $(\partial C_3)_{MNKL} = 4\partial_{[M} C_{NKL]}$. The two-brane and five-brane backgrounds are respectively ($a = 1, \dots, p$, $i = p+1, \dots, 10$, $p = 2, 5$)

$$ds_{11}^2 = H^{-2/3} (-dt^2 + dx_a dx_a) + H^{1/3} dx_i dx_i, \quad (6.12)$$

$$(\partial C_3)_{0abn} = \epsilon_{ab} \partial_n H^{-1}, \quad H = 1 + \frac{R^6}{r^6},$$

$$ds_{11}^2 = H^{-1/3} (-dt^2 + dx_a dx_a) + H^{2/3} dx_i dx_i, \quad (6.13)$$

$$(\partial C_3)_{ijkl} = \epsilon_{ijkln} \partial_n H, \quad H = 1 + \frac{R^3}{r^3}.$$

Both for the two-brane and for the five-brane, the scalar curvature is $\sim (\partial C_3)^2 \sim R^{-2}$ near $r = 0$. There is a subtle difference, however, in that the five-brane metric may be extended to a geodesically complete non-singular geometry, while the two-brane metric cannot [142, 143].

We consider small perturbations near these backgrounds which depend only on $x_\mu = (t, x_i)$. There are no scalars like ϕ or C in the $d = 11$ theory, but it is easy to check that, as in (6.8), (6.9), the “normalized” internal part of the metric $\gamma_{ab} = g_p^{-1/p} g_{ab}$, $g_p \equiv \det g_{ab}$ ($p = 2, 5$), which has a flat background value, is decoupled from the $(\partial C_3)^2$ term. As a result, the gravitons polarized parallel to the brane, $h_{ab} = \gamma_{ab} - \delta_{ab}$, $h_{aa} = 0$, have the minimal action as in (6.10),

$$S_{scal.grav.} = -\frac{1}{8\kappa_{11}^2} \int d^{11}x [\partial_i h_{ab} \partial_i h_{ab} - H(r) \partial_0 h_{ab} \partial_0 h_{ab}]. \quad (6.14)$$

6.3 The self-dual three-brane

6.3.1 Classical absorption

This section focuses on minimally coupled scalars, such as those that satisfy the Klein-Gordon equation following from (6.7), (6.10) with $H(r)$ given in (6.6). Because the classical analysis of absorption for all partial waves of such a scalar was sketched in [9] and is similar to the matching calculations that have appeared in many other places in the literature [7, 8, 73, 99, 75, 87, 21, 96, 37, 150, 71, 70], details will not be presented here. In order to employ analytical rather than numerical techniques, it is necessary as usual to take the Compton wavelength much larger than the typical radii of the black hole. Fortunately, it is precisely in this region where agreement with string theoretic models is expected [111].

Recall from chapter 4 that in d spacetime dimensions, the absorption cross-section σ_{abs}^ℓ of a massless scalar with energy ω in the ℓ -th partial wave is related to the absorption probability $1 - |S_\ell|^2$ by

$$\sigma_{\text{abs}}^\ell = \frac{2^{d-3} \pi^{(d-3)/2}}{\omega^{d-2}} \Gamma((d-3)/2) (\ell + (d-3)/2) \binom{\ell + d - 4}{\ell} (1 - |S_\ell|^2). \quad (6.15)$$

For the three-brane one takes $d = 7$ since the other 3 dimensions are compactified on T^3 . A matching calculation outlined for all partial waves in [9] gives the absorption probability

$$1 - |S_\ell|^2 = \frac{\pi^2 (\omega R)^{8+4\ell}}{[(\ell+1)!]^4 (\ell+2)^2 4^{2\ell+2}} .$$

Therefore, the classical result for the cross-section to absorb the ℓ -th partial wave of a minimally coupled scalar is, to leading order in $(\omega R)^4$,

$$\sigma_{3 \text{ class.}}^\ell = \frac{\pi^4 (\ell+3)(\ell+1)}{24 [(\ell+1)!]^4} \left(\frac{\omega R}{2} \right)^{4\ell} \omega^3 R^8 . \quad (6.16)$$

The scale parameter R of the classical three-brane solution (6.6) is related [45, 9] to the number N of coinciding microscopic three-branes by the equation

$$R^4 = \frac{\kappa_{10}}{2\pi^{5/2}} N , \quad (6.17)$$

which follows from the quantization of the three-brane charge.

There is a number of minimally coupled scalars in the theory: the dilaton, the RR scalar, and off-diagonal gravitons polarized with both indices parallel to the three-brane world-volume. The goal of the next section will be to demonstrate that the universality of leading order cross-section for these scalars, which is so obvious in the semi-classical framework, also follows from the D-brane description.

6.3.2 Universality of the absorption cross-section for minimally coupled scalars

The three-brane is the case where we know the world-volume theory the best: at low energies (and in flat space) it is $\mathcal{N} = 4$ supersymmetric $U(N)$ gauge theory where N is the number of parallel three-branes [13]. Thus, the massless fields on the world-volume are the gauge field, 6 scalars, and 4 Majorana fermions, all in the adjoint representation of $U(N)$. As we will see below, the universality of the cross-section is not trivial in the world-volume description: while for dilatons and RR scalars leading absorption proceeds by conversion into a pair of gauge bosons only, for the gravitons polarized along the brane it involves a summation over conversions into world-volume scalars, fermions, and gauge bosons.

The world-volume action, excluding all couplings to external fields, is

$$S_3 = T_3 \int d^4x \operatorname{tr} \left[-\frac{1}{4} F_{\alpha\beta}^2 + \frac{i}{2} \bar{\psi}^I \gamma^\alpha \partial_\alpha \psi_I - \frac{1}{2} (\partial_\alpha X^i)^2 + \text{interactions} \right] . \quad (6.18)$$

The indices $I = 1, \dots, 4$ and $i = 4, \dots, 9$ label representations of the R-symmetry group, $SO(6)$. It is clear from the $d = 10$ origin of this theory that the R-symmetry group is the group of spatial rotations in the uncompactified dimensions. Under this group, the gauge fields are neutral and the scalars X^i form a **6**. When the fermions are written in chiral components, $\psi_I = \begin{pmatrix} \lambda_I \\ \bar{\lambda}_I \end{pmatrix}$, the fields λ_I form a **4** of $SO(6) = SU(4)$.

In order to properly normalize the amplitudes, one needs the kinetic part of the action for the bulk fields, which is given by (6.7) and (6.10).

We also need to know how the three-brane world-volume fields couple to the bulk fields of type IIB supergravity. For a single three-brane, a κ -symmetric version of the DBI action in a non-trivial type IIB background was constructed in [33, 34] (see also [35, 36]), which, by use of superfields, captures all such couplings to leading order in derivatives of external fields. For the terms necessary to us, the generalization from $U(1)$ to $U(N)$ gauge group is straightforward. One can, in principle,

obtain detailed information about the structure of the non-abelian action by directly computing string amplitudes as done in [138]. The three-brane is well suited to this line of attack because the string theory description is known exactly and is not complicated by the difficulties one encounters in bound states of intersecting branes, such as the D1-D5-brane system [31].

$SO(6)$ invariance and power counting in the string coupling greatly restrict the possible couplings between the bulk and world-volume fields in the interaction part of the action S_{int} . In both the semi-classical and world-volume computations, the dimensionless expansion parameter is not Ng_{str} but rather $(\omega R)^4$. A cross-section which involves n powers of this parameter arises from a coupling of the bulk field to a local operator $\mathcal{O}$ of dimension $n + 4$.

The leading bosonic terms in the action for a single three-brane in a type IIB supergravity background are [32, 151, 152, 153]

$$S_3 = -T_3 \int d^4x \left(\sqrt{-\det(\hat{g} + e^{-\phi/2} \mathcal{F})} + \frac{1}{4!} \epsilon^{\alpha\beta\sigma\rho} \hat{C}_{\alpha\beta\sigma\rho} + \frac{1}{2} \hat{C}_{\alpha\beta} \tilde{\mathcal{F}}^{\alpha\beta} + \frac{1}{4} C \mathcal{F}_{\alpha\beta} \tilde{\mathcal{F}}^{\alpha\beta} \right), \quad (6.19)$$

where

$$\mathcal{F}_{\alpha\beta} = F_{\alpha\beta} + \hat{B}_{\alpha\beta}, \quad \tilde{\mathcal{F}}^{\alpha\beta} = \frac{1}{2} \epsilon^{\alpha\beta\sigma\rho} F_{\sigma\rho}, \quad \hat{g}_{\alpha\beta} = g_{MN} \partial_\alpha X^M \partial_\beta X^N, \text{ etc.},$$

and the background fields are functions of X^M . In the static gauge ($X^\alpha = x^\alpha$, $\alpha = 0, \dots, 3$) one has

$$\hat{g}_{\alpha\beta} = g_{\alpha\beta} + 2g_{i(\alpha} \partial_{\beta)} X^i + g_{ij} \partial_\alpha X^i \partial_\beta X^j.$$

In flat space the fermionic terms may be found by replacing $\hat{g}_{\alpha\beta} + e^{-\phi/2} \mathcal{F}_{\alpha\beta}$ by [33, 34, 35, 36]

$$\hat{g}_{\alpha\beta} + e^{-\phi/2} [\mathcal{F}_{\alpha\beta} - 2\bar{\Psi}(\Gamma_\alpha + \Gamma_i \partial_\alpha X^i) \partial_\beta \Psi + \bar{\Psi} \Gamma^i \partial_\alpha \Psi \bar{\Psi} \Gamma_i \partial_\beta \Psi],$$

where Ψ is the $d = 10$ Majorana-Weyl spinor (which can be split into four $d = 4$ Majorana spinors ψ_I in (6.18)) and Γ_M are the $d = 10$ Dirac matrices.

The leading-order interaction of the dilaton with world-volume fields implied by (6.19) was discussed in [9]. The coupling of the RR scalar C is similar, being related by $SL(2, R)$ duality. The coupling of the gravitons polarized parallel to the brane, $h_{\alpha\beta} = g_{\alpha\beta} - \eta_{\alpha\beta}$, to the bosonic world-volume fields can be deduced by expanding the action (6.19). At the leading order it is given by $\frac{1}{2} h^{\alpha\beta} T_{\alpha\beta}^{\text{bosons}}$ where $T_{\alpha\beta}^{\text{bosons}}$ is the energy-momentum tensor of A_α and X^i . The only possible supersymmetric extension is for $h_{\alpha\beta}$ to couple to the complete energy-momentum tensor corresponding to (6.18).

Generalizing to $U(N)$, we find that the part of S_{int} that is relevant to the leading-order absorption processes we wish to consider is³

$$S_{\text{int}} = T_3 \int d^4x \left[\text{tr} \left(\frac{1}{4} \phi F_{\alpha\beta}^2 - \frac{1}{4} C F_{\alpha\beta} \tilde{F}^{\alpha\beta} \right) + \frac{1}{2} h^{\alpha\beta} T_{\alpha\beta} \right], \quad (6.20)$$

where

$$T_{\alpha\beta} = \text{tr} \left[F_\alpha{}^\gamma F_{\beta\gamma} - \frac{1}{4} \eta_{\alpha\beta} F_{\gamma\delta}^2 - \frac{i}{2} \bar{\psi}^I \gamma_{(\alpha} \partial_{\beta)} \psi_I + \partial_\alpha X^i \partial_\beta X^i - \frac{1}{2} \eta_{\alpha\beta} (\partial_\gamma X^i)^2 \right]. \quad (6.21)$$

Let us first consider an off-diagonal graviton polarized along the brane, say h_{xy} , which is an example of a traceless perturbation h_{ab} whose quadratic action is given in (6.10)). From (6.20) one can read off the invariant amplitudes for absorption into two scalars, two fermions, or two gauge bosons:

$$\begin{aligned} \text{scalars:} \quad & \mathcal{M} = -\sqrt{2\kappa_{10}^2} (p_{1x} p_{2y} + p_{1y} p_{2x}) \\ \text{fermions:} \quad & \mathcal{M} = -\frac{1}{2} \sqrt{2\kappa_{10}^2} \bar{v}(-p_1) (\gamma_x p_{2y} + \gamma_y p_{2x}) u(p_2) \\ \text{gauge bosons:} \quad & \mathcal{M} = -\sqrt{2\kappa_{10}^2} (f_x^{(1)\beta} f_{y\beta}^{(2)} + f_y^{(1)\beta} f_{x\beta}^{(2)}) \end{aligned} \quad (6.22)$$

³We ignore the fermionic couplings like $\phi \bar{\psi}^I \gamma^\alpha \partial_\alpha \psi_I$ and similar ones for C and $h_{\alpha\beta}$ which are proportional to the fermionic equations of motion and thus give vanishing contribution to the S -matrix elements.

where p_1 and p_2 are the momenta of the outgoing particles, and the field strength polarization tensors $f_{\alpha\beta}^{(s)}$ are given by

$$f_{\alpha\beta}^{(s)} = ip_{s\alpha}\epsilon_{\beta}^{(s)} - ip_{s\beta}\epsilon_{\alpha}^{(s)} . \quad (6.23)$$

Summation over the spins of the outgoing particles can be performed using

$$\sum_s u_{(s)}(p)\bar{u}_{(s)}(p) = \not{p} , \quad \sum_s v_{(s)}(p)\bar{v}_{(s)}(p) = -\not{p} , \quad (6.24)$$

$$\sum_s \epsilon_{\alpha}^{(s)} \epsilon_{\beta}^{(s)*} = \eta_{\alpha\beta} .$$

Summing as well over different species of particles available (six different X^i , for example), one obtains

$$\begin{aligned} \text{scalars:} \quad & \overline{|\mathcal{M}|}^2 = 3\kappa_{10}^2 \omega^4 n_x^2 n_y^2 \\ \text{fermions:} \quad & \overline{|\mathcal{M}|}^2 = \kappa_{10}^2 \omega^4 (n_x^2 + n_y^2 - 4n_x^2 n_y^2) \\ \text{gauge bosons:} \quad & \overline{|\mathcal{M}|}^2 = \kappa_{10}^2 \omega^4 (1 - n_x^2 - n_y^2 + n_x^2 n_y^2) \end{aligned} \quad (6.25)$$

where $\vec{n}$ is the direction of one of the outgoing particles. In (6.25) we have anticipated conservation of energy and momentum by setting $\vec{p}_1 + \vec{p}_2 = 0$ and $\omega_1 + \omega_2 = \omega$. It is remarkable that the sum of these three quantities is independent of $\vec{n}$. Thus, if one performs the spin sums not just over all polarizations and species of particles of a given spin, but rather over all the states in the $\mathcal{N} = 4$ super-multiplet, the result is isotropic:

$$\overline{|\mathcal{M}|}^2 = \kappa_{10}^2 \omega^4 . \quad (6.26)$$

The absorption cross-section is evaluated from $\overline{|\mathcal{M}|}^2$ in precisely the same way that decay rates of massive particles are calculated in conventional 4-dimensional field theories:

$$\sigma_{3 \text{ abs}} = \frac{N^2}{2} \frac{1}{2\omega} \int \frac{d^3 p_1}{(2\pi)^3 2\omega_1} \frac{d^3 p_2}{(2\pi)^3 2\omega_2} (2\pi)^4 \delta^4(q - \sum_i p_i) \overline{|\mathcal{M}|}^2 . \quad (6.27)$$

The leading factor of N^2 accounts for the multiple branes; the $1/2$ is present because the outgoing particles are identical (this is true also of the fermions because we are working with Majorana spinors). The cross-section following from (6.26) agrees with the semi-classical $\ell = 0$ result (6.16):

$$\sigma_{3 \text{ abs}} = \frac{\kappa_{10}^2 \omega^3 N^2}{32\pi} = \sigma_{3 \text{ class.}} , \quad (6.28)$$

where we have used the relation (6.17) between N and R .

Compared to the off-diagonal graviton, the calculation of the cross-section for the RR scalar is relatively simple. Inspection of the leading order amplitudes for absorption of a dilaton and a RR scalar makes it obvious that they have the same cross-section:

$$\begin{aligned} \text{dilaton:} \quad & \mathcal{M} = \frac{1}{2} \sqrt{2\kappa_{10}^2} f_{\alpha\beta}^{(1)} f^{(2)\alpha\beta} \\ \text{RR scalar:} \quad & \mathcal{M} = -\frac{1}{2} \sqrt{2\kappa_{10}^2} f_{\alpha\beta}^{(1)} \tilde{f}^{(2)\alpha\beta} . \end{aligned} \quad (6.29)$$

To show that $\overline{|\mathcal{M}|}^2 = \kappa_{10}^2 \omega^2$ in both cases, it suffices to prove the relation

$$\sum_{\text{spins}} f_{\alpha\beta}^{(1)} f_{\gamma\delta}^{(1)*} f^{(2)\alpha\beta} f^{(2)\gamma\delta*} = \sum_{\text{spins}} f_{\alpha\beta}^{(1)} f^{(1)\gamma\delta*} \tilde{f}^{(2)\alpha\beta} \tilde{f}^{(2)\gamma\delta*} = 8(p_1 \cdot p_2)^2 . \quad (6.30)$$

The verification is straightforward algebra. The formula (6.27) applies as written to the RR scalar as well, so the agreement with the semi-classical calculation is clear.

6.3.3 Higher partial waves

A more difficult comparison is the absorption cross-section for higher partial waves. It is possible to argue, based on power counting and group theory, that the interaction term suggested in [9],

$$S_{\text{int}}^\ell = -T_3 \int d^4x \frac{1}{4 \cdot \ell!} \partial_{i_1} \cdots \partial_{i_\ell} \phi \text{tr} (X^{i_1} \cdots X^{i_\ell} F_{\alpha\beta} F^{\alpha\beta}) \quad (6.31)$$

is the only one that could possibly contribute at leading order to a given partial wave. It was shown in [9] that this term predicts a cross-section for the ℓ -th partial wave whose scaling with ω and N is in agreement with classical gravity. Here we show that, if we use the specific normalization given in (6.31), we obtain precise agreement for $\ell = 0, 1$ but disagreement for higher ℓ . Later on we will argue that the disagreement for $\ell > 1$ may be due simply to an incorrect normalization of the necessary effective action terms.

Let us restrict our attention to the dilaton. It will be obvious that all our arguments apply equally well to the RR scalar, and perhaps with a bit more attention to details to the off-diagonal gravitons. The absorption cross-section (6.16) is of order $\kappa_{10}^{\ell+2}$ for the ℓ -th partial wave. Processes which can contribute to this absorption at leading order must be mediated by an operator $\mathcal{O}$ which involves $\ell + 2$ fields. All interaction terms which involve the dilaton must include at least two powers of $F_{\alpha\beta}$ because of the restricted way in which the dilaton enters the DBI action. Since the gauge bosons are neutral under $SO(6)$, the other ℓ fields must be responsible for balancing the $SO(6)$ transformation properties of the ℓ -th partial wave. We want to argue that the only way this can be done is to use ℓ powers of the scalar fields X^i . To do this it is necessary to know something about the addition rules for representations of $SO(6) = SU(4)$. The $SU(4)$ Young tableaux for the fields in question are

$$\lambda_I: \square \quad \bar{\lambda}^I: \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \quad X^i: \begin{array}{|c|} \hline \square \\ \hline \end{array} \quad \phi_\ell: \underbrace{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \end{array} \cdots \begin{array}{|c|} \hline \square \\ \hline \end{array}}_{\ell \text{ columns}} \quad (6.32)$$

where ϕ_ℓ stands for the ℓ -th partial wave. There is a complicated general procedure known as the Littlewood-Richardson rule for taking tensor products of representations of $SU(N)$. It becomes much simpler when one of the factors is a fundamental representation (k boxes in a single column) [154]. In that case, one adds these k boxes to the other factor's Young tableaux in all possible ways, modulo the restriction that not more than one box can be added to any given row. Then one eliminates all columns containing N boxes. Using this rule it is easy to show that with ℓ fundamental representations as the factors in a tensor product, it is impossible to obtain the tableau for ϕ_ℓ unless all the factors are **6** representations.

A shorter argument can be made based on purely dimensional grounds. The dimension of $\mathcal{O}$ must be $\ell + 4$ or the cross-section would be suppressed by extra powers of ω . Since $F_{\alpha\beta}^2$ has dimension 4, the other ℓ fields must add ℓ to the total dimension of $\mathcal{O}$. The only possibility is ℓ scalars.

The upshot is that the only possible term in the action which can contribute at leading order to the absorption of a dilaton in the ℓ -th partial wave is, up to normalization, given by (6.31). The normalization given there is the one arising from Taylor expanding $\phi(X)$.⁴ Actually, this action is not quite the one we want for $\ell > 1$: in addition to mediating the process which contributes at leading order to the absorption of the ℓ -th partial wave, it makes subleading contributions to lower partial waves. In order to isolate the ℓ -th partial wave, the product $X^{i_1} \cdots X^{i_\ell}$ should be replaced by an expression which transforms irreducibly under $SO(6)$. For $\ell = 2$ and $\ell = 3$, the appropriate replacements are

$$\begin{aligned} \ell = 2: \quad X^{i_1} X^{i_2} &\rightarrow X^{i_1} X^{i_2} - \frac{1}{6} \delta^{i_1 i_2} X^2 \\ \ell = 3: \quad X^{i_1} X^{i_2} X^{i_3} &\rightarrow X^{i_1} X^{i_2} X^{i_3} - \frac{3}{8} \delta^{(i_1 i_2} X^{i_3)} X^2 \end{aligned} \quad (6.33)$$

⁴As we argue below, it is possible that this local expansion is not consistent with the string amplitude calculations. This may remove the discrepancy for $\ell > 1$.

One thus subtracts an $\ell = 0$ contribution from the putative $\ell = 2$ term in (6.31), and an $\ell = 1$ contribution from $\ell = 3$. The whole Taylor series can be thus reshuffled.

Because the replacements in (6.33) project out a final state with definite ℓ , it is permissible to let the initial dilaton wave-function be $e^{-i\omega(t-x^1)}$ as usual. The derivatives ∂_i in (6.31) can then be replaced by $i\omega\delta_i^1$. The spin-summed amplitudes squared for absorption of the first few partial waves are

$$\begin{aligned}\ell = 0: \quad & \overline{|\mathcal{M}|}^2 = 4\kappa_{10}^2(p_1 \cdot p_2)^2 \\ \ell = 1: \quad & \overline{|\mathcal{M}|}^2 = \frac{4}{\sqrt{\pi}}\kappa_{10}^3\omega^2(p_1 \cdot p_2)^2 \\ \ell = 2: \quad & \overline{|\mathcal{M}|}^2 = \frac{10}{3\pi}\kappa_{10}^4\omega^4(p_1 \cdot p_2)^2 \\ \ell = 3: \quad & \overline{|\mathcal{M}|}^2 = \frac{5}{2\pi^{3/2}}\kappa_{10}^5\omega^6(p_1 \cdot p_2)^2 .\end{aligned}\tag{6.34}$$

As before, p_1 and p_2 are the momenta of the outgoing gauge bosons. The gauge bosons are identical particles, as are the ℓ outgoing scalars, whose momenta will be labeled $p_3, \dots, p_{\ell+2}$. To avoid over-counting in the integral over phase space, we must include a factor of $1/(2 \cdot \ell!)$ in the expression for the cross-section. The cross-section also includes an explicit factor of $1/(2\omega)$ from the normalization of the incoming dilaton. Altogether,

$$\sigma_D^\ell = \frac{N^{\ell+2}}{2 \cdot \ell!} \frac{1}{2\omega} \int \frac{d^3 p_1}{(2\pi)^3 2\omega_1} \dots \frac{d^3 p_{\ell+2}}{(2\pi)^3 2\omega_{\ell+2}} (2\pi)^4 \delta^4(q - \sum_i p_i) \overline{|\mathcal{M}|}^2 .\tag{6.35}$$

The momentum integrations take the following form:

$$I_\ell = \int \frac{d^3 p_1}{2\omega_1} \dots \frac{d^3 p_{\ell+2}}{2\omega_{\ell+2}} \delta^4(q - \sum_i p_i) (p_1 \cdot p_2)^2 = \frac{3\pi^{\ell+1}}{2^{\ell+1}} \frac{\omega^{2\ell+4}}{(\ell+2)!(\ell+3)!} .\tag{6.36}$$

The quickest way to establish (6.36) is to Fourier transform to position space.

The final results for the first few ℓ are

$$\begin{aligned}\sigma_{D\text{-brane}}^{\ell=0} &= \sigma_{\text{class.}}^{\ell=0} . \\ \sigma_{D\text{-brane}}^{\ell=1} &= \sigma_{\text{class.}}^{\ell=1} . \\ \sigma_{D\text{-brane}}^{\ell=2} &= \frac{9}{5} \sigma_{\text{class.}}^{\ell=2} . \\ \sigma_{D\text{-brane}}^{\ell=3} &= \frac{24}{5} \sigma_{\text{class.}}^{\ell=3} .\end{aligned}\tag{6.37}$$

If the $\ell = 2$ and $\ell = 3$ D-brane cross-sections had been smaller than the corresponding classical results, one might have wondered if the argument around (6.32) might possibly be invalidated by some peculiar interaction. But any such additional interaction could only increase the D-brane cross-section, making the disagreement even worse.

It is worth noting that nowhere in the literature have higher partial waves been compared with complete success between D-brane models and General Relativity. Agreement up to numerical factors was obtained in [100] for effective string models of four- and five-dimensional black holes, and independently in [155] for the five-dimensional case; but in the absence of a well articulated prescription for coupling the effective string to the bulk fields it is difficult to tell much about numerical coefficients.

The agreement of the $\ell = 1$ cross-section certainly encourages us to believe that the D-brane description is capable of handling partial waves correctly. On the level of effective field theory this may seem peculiar because, in this low-energy description, the D-brane has no thickness. How can an object with no extent in transverse dimensions absorb particles with angular momentum? Power counting and group theory alone seem to dictate the answer (6.31), up to normalization.

Fortunately, with three-branes the string theory prescription for the couplings to external fields is more directly accessible than for other solitonic models of black holes. While it does indeed appear that $\ell > 1$ partial waves present a test which the DBI action, supplemented by the prescription of [9] to obtain the normalization of (6.31) via a Taylor expansion, fails to pass, we expect that a proper string theoretic treatment will once again yield agreement. For $\ell = 1$, a full-fledged disk amplitude computation verifies both the form of (6.31) and the normalization shown. The $\ell = 1$ disk amplitude, which involves one bulk insertion and three boundary operators, is easy to deal with because the insertion of one scalar vertex operator on the boundary simply generates a spacetime translation. When more than one such operators are present, one encounters singularities in their mutual collisions that need a careful treatment. The necessary calculations look quite complicated: for example, computing the full $\ell = 2$ amplitude would be equivalent via the prescription of [137] to computing a six point type I disk amplitude. They nevertheless seem highly worthwhile as a means to refine our understanding of the world-volume action.

6.4 Absorption cross-sections for M-branes

A remarkable aspect of the agreement between the string theoretic and the classical results for three-branes is that it holds exactly for any value of N , including $N = 1$. As explained in the introduction, the classical geometry should be trusted only in the limit $g_{\text{str}}N \rightarrow \infty$. Thus, the agreement of the absorption cross-sections for $N = 1$ suggests that our calculations are valid even in the limit $g_{\text{str}} \rightarrow \infty$, provided that $g_{\text{str}}\alpha'^2\omega^4$ is kept small. The supersymmetric non-renormalization theorems are probably at work here, insuring that there are no string loop corrections.

In this section we would like to ask whether the exact agreement between the absorption cross-sections is also found for the two-branes and five-branes of M-theory. These branes have much in common with the self-dual three-brane of type IIB theory: their entropies scale with the temperature in agreement with the scaling for a gas of massless fields on the world-volume [46], while the scaling of their absorption cross-sections with ω agrees with estimates from the world-volume effective theory [9]. While the effective actions for one two-brane [156] and one five-brane [157, 158, 159] are now known in some detail, their generalizations to $N > 1$ remain somewhat obscure.⁵ In this section we compare the cross-sections for $N = 1$ and find that, in contrast to the three-branes, there is no exact agreement in the normalizations. This is probably due to the fact that M-theory has no parameter like g_{str} that can be dialed to make the classical solution reliable.

First we discuss absorption of longitudinally polarized gravitons by an two-brane. The massless fields in the effective action are 8 scalars and 8 Majorana fermions. The longitudinal graviton couples to the energy momentum tensor on the world-volume, $T_{\alpha\beta}$. The terms in the effective action necessary to describe the absorption of h_{xy} are ($i = 3, \dots, 10$; $I = 1, \dots, 8$)

$$S_2 = T_2 \int d^3x \left[-\frac{1}{2} \partial_\alpha X^i \partial^\alpha X^i + \frac{i}{2} \bar{\psi}^I \gamma^\alpha \partial_\alpha \psi^I + \sqrt{2} \kappa_{11} h_{xy} (\partial_x X^i \partial_y X^i - \frac{i}{4} \bar{\psi}^I (\gamma_x \partial_y + \gamma_y \partial_x) \psi^I) \right], \quad (6.38)$$

where h_{xy} is the canonically normalized field which enters the $D = 11$ spacetime action as (cf. (6.14))

$$-\frac{1}{2} \int d^{11}x \partial_M h_{xy} \partial^M h_{xy}.$$

⁵It is believed, for instance, that $N > 1$ coincident five-branes are described by non-trivial conformally invariant theories in $5 + 1$ dimensions. Comparisons with classical gravity of the kind made in [46, 9] and here are among the ways of learning more about this theory.

The absorption cross-section is found using the Feynman rules in a manner analogous to the three-brane calculation of section 6.3. For the 8 scalars, we find that the matrix element squared (with all the relevant factors included) is

$$\frac{\kappa_{11}^2 \omega^4}{2} 4n_x^2 n_y^2 ,$$

where $\vec{n}$ is the unit vector in the direction of one of the outgoing particles. For the 8 Majorana fermions, the corresponding object summed over the final polarizations is

$$\frac{\kappa_{11}^2 \omega^4}{2} (n_x^2 - n_y^2)^2 .$$

Adding them up, we find that the dependence on direction cancels out, just as in the three-brane case. The sum must be multiplied by the phase space factor

$$\frac{1}{2\omega} \frac{1}{2\pi} \int \frac{d^2 p_1}{2\omega_1} \int \frac{d^2 p_2}{2\omega_2} \delta^2(\vec{p}_1 + \vec{p}_2) \delta(\omega_1 + \omega_2 - \omega) = \frac{1}{8\omega^2} ,$$

so that the total cross-section is

$$\sigma_{2 \text{ abs}} = \frac{\kappa_{11}^2 \omega^2}{16} . \quad (6.39)$$

This does not agree with the classical result for N set to 1 [9, 160],

$$\sigma_{2 \text{ class.}} = \frac{\pi^4}{12} \omega^2 R^9 = \frac{1}{6\sqrt{2}\pi} \kappa_{11}^2 \omega^2 N^{3/2} ,$$

where we have used the two-brane charge quantization to express R^9 in terms of N . Notice that even the power of π does not match. This situation is reminiscent of the discrepancy in the near-extremal entropy where the relative factor was a transcendental number involving $\zeta(3)$ [46].

Now we show that, just as for the three-brane, the transversely polarized gravitons have the same absorption cross-section as the longitudinally polarized gravitons. The coupling of h_{67} to scalars is given by

$$-T_2 \int d^3 x \sqrt{2} \kappa_{11} h_{67} \partial_\alpha X^6 \partial^\alpha X^7 , \quad (6.40)$$

while pairs of fermions are not produced because the coupling to them vanishes on shell. The matrix element squared is $\kappa_{11}^2 \omega^4 / 2$. Multiplying this by the phase space factor, we again find the cross-section (6.39).

Now we turn to the five-brane. The massless fields on the five-brane form a tensor multiplet consisting of 5 scalars, 2 Weyl fermions and the antisymmetric tensor $\mathcal{B}_{\alpha\beta}$ with anti-self-dual strength [161, 162]. The transverse gravitons are again the easier case because they produce pairs of scalars only (the coupling to 2 fermions vanishes on shell). The necessary coupling of h_{67} is similar to (6.40):

$$-T_5 \int d^6 x \sqrt{2} \kappa_{11} h_{67} \partial_\alpha X^6 \partial^\alpha X^7 .$$

The matrix element squared is $\kappa_{11}^2 \omega^4 / 2$, while the phase space factor is now

$$\frac{1}{2\omega} \frac{1}{(2\pi)^4} \int \frac{d^5 p_1}{2\omega_1} \int \frac{d^5 p_2}{2\omega_2} \delta^5(\vec{p}_1 + \vec{p}_2) \delta(\omega_1 + \omega_2 - \omega) = \frac{\omega}{2^7 \cdot 3\pi^2} , \quad (6.41)$$

so that the absorption cross-section is

$$\sigma_{5 \text{ class.}} = \frac{\kappa_{11}^2 \omega^5}{2^8 \cdot 3\pi^2} . \quad (6.42)$$

To discuss the absorption of longitudinally polarized gravitons, h_{xy} , we need the action ($i = 6, \dots, 10$; $I = 1, 2$)

$$S = T_5 \int d^6x \left[-\frac{1}{2} \partial_\alpha X^i \partial^\alpha X^i - \frac{1}{12} \mathcal{H}_{\alpha\beta\gamma}^2 + i \bar{\psi}^I \gamma^\alpha \partial_\alpha \psi^I + \sqrt{2} \kappa_{11} h_{xy} \left(\partial_x X^i \partial_y X^i + \frac{1}{2} \mathcal{H}_{x\beta\gamma}^- \mathcal{H}_y^{-\beta\gamma} - \frac{i}{2} \bar{\psi}^I (\gamma_x \partial_y + \gamma_y \partial_x) \psi^I \right) \right]. \quad (6.43)$$

To describe interaction of the anti-self-dual antisymmetric tensor with external field we follow the covariant approach of [163] using the standard unconstrained propagator for $\mathcal{B}_{\alpha\beta}$ and replacing $\mathcal{H}$ by its anti-self-dual part $\mathcal{H}^- = \frac{1}{2}(\mathcal{H} - \tilde{\mathcal{H}})$ in the vertices.

For the 5 scalars, we find that the matrix element squared (with all the relevant factors included) is

$$\frac{\kappa_{11}^2 \omega^4}{2} \frac{5}{2} n_x^2 n_y^2. \quad (6.44)$$

For the 2 Weyl fermions, the corresponding object summed over the final polarizations is⁶

$$\frac{\kappa_{11}^2 \omega^4}{2} (n_x^2 + n_y^2 - 4 n_x^2 n_y^2). \quad (6.45)$$

Finally, the contribution of the anti-self-dual gauge field turns out to be equal to that of the usual, unconstrained $\mathcal{H}_{\alpha\beta\gamma}$ divided by 2. The matrix element squared is, therefore,⁷

$$|\mathcal{M}|^2 = \frac{1}{2} 2 \kappa_{11}^2 \frac{1}{4} \sum [\mathcal{H}_{x\beta\gamma}^{(1)} \mathcal{H}_y^{(2)\beta\gamma} + \mathcal{H}_{x\beta\gamma}^{(2)} \mathcal{H}_y^{(1)\beta\gamma}] \mathcal{H}_{x\alpha\delta}^{(1)*} \mathcal{H}_y^{(2)\alpha\delta*}, \quad (6.46)$$

where we have also included 1/2 because the outgoing particles are identical. Sums over polarizations are to be performed with

$$\sum \epsilon_{\alpha\beta} \epsilon_{\gamma\delta}^* = \eta_{\alpha\gamma} \eta_{\beta\delta} - \eta_{\alpha\delta} \eta_{\beta\gamma}.$$

The entire calculation is lengthy, but the end result is simple,⁸

$$|\overline{\mathcal{M}}|^2 = \frac{\kappa_{11}^2 \omega^4}{2} \left(1 - n_x^2 - n_y^2 + \frac{3}{2} n_x^2 n_y^2 \right). \quad (6.47)$$

Adding up the contributions of the entire tensor multiplet, we find that all the direction-dependent terms cancel out, just as they did for the three-brane and the two-brane. Multiplying by the phase space factor (6.41), we find that the total cross-section for the longitudinally polarized gravitons is again given by (6.42). This turns out to be a factor of 4 smaller than the classical result,

$$\sigma_{5 \text{ class.}} = \frac{2\pi^3}{3} \omega^5 R^9 = \frac{\kappa_{11}^2 N^3 \omega^5}{2^6 \cdot 3\pi^2}, \quad (6.48)$$

evaluated for $N = 1$, i.e. $\sigma_{5 \text{ abs}} = \frac{1}{4} \sigma_{5 \text{ class.}}^{(N=1)}$. This discrepancy is relatively minor and is of a kind that could easily be produced by a calculational error. However, having checked our calculations a number of times, we believe that the factor of 4 discrepancy is real.

⁶It is interesting to observe that, if we consider an $\mathcal{N} = 1$ multiplet consisting of 1 Weyl fermion and 4 scalars, then the “tensor term” $n_x^2 n_y^2$ cancels out in the direction dependence. The same cancelation occurs for the three-brane (both for the $\mathcal{N} = 1$ vector multiplet and for the $\mathcal{N} = 1$ hypermultiplet).

⁷Since the propagator of $\mathcal{B}_{\alpha\beta}$ is taken to be non-chiral, it is sufficient to do the replacement $\mathcal{H}\mathcal{H} \rightarrow \mathcal{H}^- \mathcal{H}^-$ in only one of the two stress tensor factors in $|\mathcal{M}|^2$. The relevant part of $\mathcal{H}^- \mathcal{H}^-$ is $\frac{1}{4}(\mathcal{H}\mathcal{H} + \tilde{\mathcal{H}}\tilde{\mathcal{H}})$ which is equal to $\frac{1}{2}\mathcal{H}\mathcal{H}$ for off-diagonal components of the stress tensor.

⁸This answer passes also a number of heuristic checks. For instance, for the $\mathcal{N} = 1$ multiplet including the gauge field, 1 scalar and 1 Weyl fermion, the $n_x^2 n_y^2$ terms again cancels out.

A conclusion that we may draw from this section is that, although the single M-brane cross-sections scale with the energy in the same way as the classical cross-sections, the normalizations do not agree. The five-brane comes much closer to agreement than the two-brane, which may be connected to the fact that the five-brane supergravity solution is completely non-singular. For $N = 1$, however, the curvature of the solution is of order of the 11-dimensional Planck scale. Obviously, the 11-dimensional supergravity is at best a low-energy approximation to M-theory. The M-theory effective action should contain higher-derivative terms weighted by powers of κ_{11} , by analogy with the α' and g_{str} expansions of the string effective action. Thus, for $N = 1$, the classical solution may undergo corrections of order one which we believe to be the source of the discrepancy. For large N , however, we expect the M-theory cross-section to agree exactly with the classical cross-section. We hope that these considerations will serve as a useful guide in constructing the world volume theory of N coincident five-branes.

6.5 Conclusions

In this chapter we have provided new evidence, furthering the earlier results of [9], that there exists exact agreement between the classical and the D-brane descriptions of the self-dual three-brane of type IIB theory. The specific comparisons that we have carried out involve probing an extremal three-brane with low-energy massless quanta incident from the outside. As argued in [9] and here, the great advantage of the three-brane is that both the perturbative string theory and the classical supergravity calculations are under control and yield expansions in the same dimensionless expansion parameter $(\omega R)^4 \sim N g_{\text{str}} \alpha'^2 \omega^4$, which may be kept small.

While the low-energy physics of the three-brane is described by a $\mathcal{N} = 4$ SYM theory, probing it from the extra dimensions provides a new point of view and allows for a variety of interesting calculations. One example is the absorption of a graviton polarized parallel to the brane which, as discussed in section 6.3, couples to the energy-momentum tensor in 3+1 dimensions. Because of the existence of the transverse momentum, the kinematics for this process is that of a decay of a massive spin-two particle into a pair of massless world-volume modes. Summing over all the states in the $\mathcal{N} = 4$ multiplet we find that the rate is completely isotropic, which is undoubtedly related to the conformal invariance of the theory. We should emphasize, however, that we are exploring the properties of this theory away from the BPS limit; therefore, they are not determined by supersymmetry alone. For this reason, we find it remarkable that the net absorption cross-section has the same value as that of the dilaton and the RR scalar, and which agrees with the cross-section found in classical supergravity.

We have also carried out similar explorations of the physics of M-branes. These studies are hampered, to a large extent, by insufficient understanding of the multiple coincident branes of M-theory. For a single two-brane, a formal application of classical gravity gives an answer which is off by a factor of $4\sqrt{2}/(3\pi)$, while for a single five-brane – off only by a factor of 4. For the five-brane the discrepancy is relatively minor which, we are tempted to speculate, is due to the fact that its classical geometry is completely non-singular, just like that of the three-brane. We should keep in mind, however, that for singly charged branes the quantum effects of M-theory are expected to be important, and the classical reasoning should not be trusted.

While in this chapter we have probed three-branes with massless particles, we may contemplate another interesting use of the three-brane: it may be used as a probe by itself [164]. For example, N coincident three-branes may be probed by a three-brane parallel to them. This situation is described by a $3 + 1$ dimensional $\mathcal{N} = 4$ supersymmetric $U(N + 1)$ gauge theory, with gauge symmetry broken to $U(N) \times U(1)$. A more complicated theory will arise if a three-brane is used to probe a stringy $d = 4$ black hole. As was shown in [26, 165], the $d = 4$ extremal black holes with regular horizons (which are parametrized by 4 charges [61]) can be represented by 1/8 supersymmetric configurations of four intersecting three-branes wrapped over a 6-torus. Following [93, 166] one can find an action for a classical three-brane probe moving in this geometry. If the probe is oriented parallel to one of

the four source three-branes, the resulting moduli space metric is

$$ds_6^2 = H_1 H_2 H_3 (dr^2 + r^2 d\Omega_2^2) + H_1 dy_1^2 + H_2 dy_2^2 + H_3 dy_3^2, \quad H_i = 1 + \frac{R_i}{r},$$

where y_i are toroidal coordinates transverse to the probe. In contrast to the case of the 5-brane-plus-string-plus-momentum configuration describing $d = 5$ regular extremal black holes with 3 charges [82] where one finds [167] that the non-compact part of the moduli space metric is multiplied by the product of two harmonic functions, here we obtain the product of *three* harmonic functions. This is, however, exactly what is needed to get the same near-horizon ($r \rightarrow 0$) behaviour as found in [167], i.e. that the 3-dimensional non-compact part of the moduli space metric becomes flat: $ds_3^2 \rightarrow R_1 R_2 R_3 (d\rho^2 + \rho^2 d\Omega_2^2)$, $\rho \equiv r^{-1/2} \rightarrow \infty$. This close similarity between a $d = 5$ black hole probed by a string and a $d = 4$ black hole probed by a three-brane strongly suggests that one can obtain important information about these black holes by studying the corresponding three-brane world-volume theory. This theory in the presence of intersecting branes is necessarily more intricate than the simpler, parallel brane case discussed in the main part of this chapter.

Clearly, there is much work to be done before we claim a complete understanding of the remarkable duality between the D-brane and the classical descriptions of the three-brane. The complexities that affect the higher-spin classical equations remain to be disentangled. The numerical discrepancies for $\ell > 1$ partial waves suggest an incomplete understanding of the low-energy effective action. As we have argued in section 6.3, a direct string calculation is necessary to check normalizations. And finally, we need to push our methods away from extremality where the physics is necessarily more complicated, involving thermal field theory in 3+1 dimensions. We hope that detailed insight into such a theory will help explain the specific factors appearing in the near-extremal entropy [45].

We feel that further efforts in the directions mentioned above are worthwhile because they offer a promise of building a theory of certain black holes in terms of a manifestly unitary theory – perturbative string theory.

Chapter 7

Cross-sections and Schwinger terms in the world-volume

7.1 Introduction

Extremal black holes with non-vanishing horizon area may be embedded into string theory or M-theory using intersecting p -branes [168, 62, 5, 6, 84, 85, 169, 26, 170]. These configurations are useful for a microscopic interpretation of the Bekenstein-Hawking entropy. The dependence of the entropy on the charges, the non-extremality parameter, and the angular momentum suggests a connection with $1+1$ dimensional conformal field theory [5, 6, 171, 84, 85, 26]. This “effective string” is essentially the intersection of the p -branes. Calculations of emission and absorption rates [6, 7, 8, 99, 75, 87, 21, 96, 112, 102, 103, 83, 101, 104, 22, 113, 114] provide further tests of the “effective string” models of $D = 5$ black holes with three charges and of $D = 4$ black holes with four charges. For minimally coupled scalars the functional dependence of the greybody factors on the frequency agrees exactly with semi-classical gravity, providing a highly non-trivial verification of the effective string idea [75, 87]. Similar successes have been achieved for certain non-minimally coupled scalars, which were shown to couple to higher dimension operators on the effective string. For instance, the fixed scalars [66, 67, 70] were shown in chapter 3 to couple to operators of dimension $(2, 2)$ [21] while the “intermediate” scalars [101] – to operators of dimension $(2, 1)$ and $(1, 2)$. Unfortunately, there is little understanding of the “effective string” from first principles, and some of the more sensitive tests reveal this deficiency. For instance, semi-classical gravity calculations of the fixed scalar absorption rates for general black hole charges reveal a gap in our understanding of higher dimension operators [83]. A similar problem occurs when one attempts a detailed effective string interpretation of the higher partial waves of a minimally coupled scalar, as we saw in chapter 4. Even the s -wave absorption by black holes with general charges is complex enough that it is not reproduced by the simplest effective string model [112, 113, 114]. These difficulties by no means invalidate the general qualitative picture, but they do pose some interesting challenges. In order to gain insight into the relation between supergravity and D-branes it is useful to study, in addition to the intersecting branes, the simpler configurations which involve parallel branes only. This chapter, based on the paper [39], shows how absorption calculations in linearized supergravity are related to certain Schwinger terms in the operator algebra of the D-brane world-volume gauge theory.

A microscopic interpretation of the entropy of near-extremal p -branes was first studied in [45, 46]. It was found that the scaling of the Bekenstein-Hawking entropy with the temperature agrees with that for a massless gas in p dimensions only for the “non-dilatonic p -branes”: namely, the self-dual three-brane of the type IIB theory, and the 2- and 5-branes of M-theory. Their further study was

undertaken in [9, 10], where low-energy absorption cross-sections for certain incident massless particles were compared between semi-classical supergravity and string or M-theory. For three-branes, exact agreement was found for the leading low-energy behavior of the absorption cross-sections for dilatons [9] as well as R-R scalars and for gravitons polarized along the brane [10]. The string-theoretic description of macroscopic three-branes can be given in terms of many coincident D3-branes [172, 4]. Indeed, N parallel D3-branes are known to be described by a $U(N)$ gauge theory in 3+1 dimensions with $\mathcal{N} = 4$ supersymmetry [13]. This theory has a number of remarkable properties, including exact S-duality, and we will be able to draw on a known non-renormalization theorem in explaining the absorption by the three-branes.

From the point of view of supergravity, the three-brane is also special because its extremal geometry,

$$ds^2 = \left(1 + \frac{R^4}{r^4}\right)^{-1/2} (-dt^2 + dx_1^2 + dx_2^2 + dx_3^2) + \left(1 + \frac{R^4}{r^4}\right)^{1/2} (dr^2 + r^2 d\Omega_5^2), \quad (7.1)$$

is non-singular [142], while the dilaton background is constant. Instead of a singularity at $r = 0$ we find an infinitely long throat whose radius is determined by the charge (the vanishing of the horizon area is due to the longitudinal contraction). Thus, for a large number N of coincident branes, the curvature may be made arbitrarily small in Planck units. For instance, for N D3-branes, the curvature is bounded by a quantity of order

$$\frac{1}{\sqrt{N\kappa}} \sim \frac{1}{\alpha' \sqrt{N g_{\text{str}}}}. \quad (7.2)$$

In order to suppress the string scale corrections to the classical metric, we need to take the limit $N g_{\text{str}} \rightarrow \infty$.

The tension of a D3-brane depends on g_{str} and α' only through the ten-dimensional gravitational constant $\kappa = 8\pi^{7/2} g \alpha'^2$:

$$T_{(3)} = \frac{\sqrt{\pi}}{\kappa}. \quad (7.3)$$

This suggests that we can compare the expansions of various quantities in powers of κ between the microscopic and the semi-classical descriptions. Indeed, the dominant term in the absorption cross-section at low energy is (see chapter 6)

$$\sigma = \frac{\pi^4}{8} \omega^3 R^8 = \frac{\kappa^2 \omega^3 N^2}{32\pi}, \quad (7.4)$$

which agrees between semi-classical supergravity and string theory.

It is important to examine the structure of higher power in g_{str} corrections to the cross-section [173]. In semi-classical supergravity the only quantity present is κ , and corrections to (7.4) can only be of the form

$$a_1 \kappa^3 \omega^7 + a_2 \kappa^4 \omega^{11} + \dots \quad (7.5)$$

However, in string theory we could in principle find corrections even to the leading term $\sim \omega^3$, so that

$$\sigma_{\text{string}} = \frac{\kappa^2 \omega^3 N^2}{32\pi} (1 + b_1 g_{\text{str}} N + b_2 (g_{\text{str}} N)^2 + \dots) + \mathcal{O}(\kappa^3 \omega^7). \quad (7.6)$$

Presence of such corrections would spell a manifest disagreement with supergravity because, as we have explained, the comparison has to be carried out in the limit $N g_{\text{str}} \rightarrow \infty$. The main purpose of this chapter is to show that, in fact, $b_i = 0$ due to a non-renormalization theorem in $D = 4$ $\mathcal{N} = 4$ SYM theory.

In section 7.2 we present a detailed argument for the absence of such corrections in the absorption cross-section of gravitons polarized along the brane. These gravitons couple to the stress-energy tensor on the world volume and we will show that the absorption cross-section is, up to normalization, the

central term in the two-point function of the stress-energy tensor. The fact that $b_i = 0$ follows from the fact that the one-loop calculation of the central charge is exact in $D = 4$ $\mathcal{N} = 4$ SYM theory.

The connection between absorption of gravitons polarized along the brane and Schwinger terms in the stress-energy correlators of the world volume theory is a general phenomenon that holds for all branes. In section 7.3 we explore this connection to deduce some properties of the stress-energy tensor OPE's for multiple 2-branes and 5-branes of M-theory, as well as for multiple 5-branes of string theory.

7.2 D-brane approach to absorption

It was probably Callan who first realized that, in terms of D-brane models, absorption cross-sections correspond up to a simple overall factor to discontinuities of two-point functions of certain operators on the D-brane world volume [174]. This realization was exploited in [100] and in chapter 5. Consider massless scalar particles in ten dimensions normally incident upon D3-branes. If the coupling to the brane is given by

$$S_{\text{int}} = \int d^4x \phi(x, 0) \mathcal{O}(x) , \quad (7.7)$$

where $\phi(x, 0)$ is a canonically normalized scalar field evaluated on the brane, and $\mathcal{O}$ is a local operator on the brane, then the precise correspondence is

$$\sigma = \frac{1}{2i\omega} \text{Disc } \Pi(p) \Big|_{\substack{p^0 = \omega \\ \vec{p} = 0}} . \quad (7.8)$$

Here ω is the energy of the incident particle, and

$$\Pi(p) = \int d^4x e^{ip \cdot x} \langle \mathcal{O}(x) \mathcal{O}(0) \rangle . \quad (7.9)$$

When $\mathcal{O}$ is a scalar in the world volume theory, $\Pi(p)$ depends only on $s = p^2$, and $\text{Disc } \Pi(p)$ is computed as the difference of Π evaluated for $s = \omega^2 + i\varepsilon$ and $s = \omega^2 - i\varepsilon$. In the case of the graviton, we shall see that $\Pi(p)$ is a polynomial in p times a function of s , so the evaluation of $\text{Disc } \Pi(p)$ is equally straightforward.

The validity of (7.8) depends on ϕ being a canonically normalized field. In form it is almost identical to the standard expression for the decay rate of an unstable particle of mass ω . The dimensions are different, however: σ is the cross-section of the three-brane per unit world volume, and so has dimensions of (length)⁵. Similar formulas can be worked out for branes of other dimensions and for near-extremal branes, although away from extremality (7.9) would become a thermal Green's function (see chapter 5).

Now let us work through the example of the graviton. The three-brane world volume theory is $\mathcal{N} = 4$ supersymmetric $U(N)$ gauge theory, where N is the number of parallel three-branes [13]. Thus, the massless fields on the world volume are the gauge field, six scalars, and four Majorana fermions, all in the adjoint representation of $U(N)$. To lowest order in κ , and ignoring the couplings to the bulk fields, the world volume-action is ($I = 1, \dots, 4$; $i = 4, \dots, 9$)

$$S_3 = \int d^4x \text{tr} \left[-\frac{1}{4} F_{\alpha\beta}^2 + \frac{i}{2} \bar{\psi}^I \gamma^\alpha \partial_\alpha \psi_I + \frac{1}{2} (\partial_\alpha X^i)^2 + \text{interactions} \right] . \quad (7.10)$$

The interactions referred to here are the standard renormalizable ones of $\mathcal{N} = 4$ super-Yang-Mills (see for example [175] for the complete flat-space action). In equation (6.18), a factor of the three-brane tension $T_{(3)}$ appeared in front of the action. It is convenient to work with canonically normalized fields, and so, relative to the conventions of chapter 6 we have absorbed a factor of $\sqrt{T_{(3)}}$ into A_μ ,

ψ^I , and X^i . Another difference between chapter 6 and our present conventions is that we work here with a mostly minus metric and the spinor conventions of [175].

The full (and as yet unknown) action for multiple three-branes is non-polynomial, and (7.10) includes only the dimension 4 terms. The higher dimension terms will appear with powers of $1/T_{(3)}$, which is to say positive powers of κ . They could give rise to corrections of the form (7.5), but not (7.6).

Despite our ignorance of the full action for multiple coincident three-branes, one can be fairly confident in asserting that the external gravitons polarized parallel to the brane couple via

$$S_{\text{int}} = \int d^4x \frac{1}{2} h^{\alpha\beta} T_{\alpha\beta} , \quad (7.11)$$

where $h_{\alpha\beta} = g_{\alpha\beta} - \eta_{\alpha\beta}$ is the perturbation in the metric, and $T_{\alpha\beta}$ is the stress-energy tensor:

$$\begin{aligned} T_{\alpha\beta} = \text{tr} \big[& -F_\alpha{}^\gamma F_{\beta\gamma} + \frac{1}{4} \eta_{\alpha\beta} F_{\gamma\delta}^2 + \frac{i}{2} \bar{\psi}^I \gamma_{(\alpha} \partial_{\beta)} \psi_I \\ & + \frac{2}{3} \partial_\alpha X^i \partial_\beta X^i - \frac{1}{6} \eta_{\alpha\beta} (\partial_\gamma X^i)^2 - \frac{1}{3} X^i \partial_\alpha \partial_\beta X^i + \frac{1}{12} \eta_{\alpha\beta} X^i \square X^i \\ & + \text{interactions} \big] . \end{aligned} \quad (7.12)$$

This “new improved” form of the stress-energy tensor [176] is chosen so that $\partial_\mu T^{\mu\nu} = 0$ and $T^\mu{}_\mu = 0$ on shell. The scalar terms differ from the canonical form

$$T_{\alpha\beta}^{(\text{can})} = \text{tr} \left[\partial_\alpha X^i \partial_\beta X^i - \frac{1}{2} \eta_{\alpha\beta} (\partial_\gamma X^i)^2 + \dots \right] . \quad (7.13)$$

The difference arises from adding a term $-\frac{1}{12} \sqrt{-g} R \text{tr} X^2$ to the lagrangian so that the scalars are conformally coupled.

Choosing the scalars to be minimally or conformally coupled does not affect the one-loop result for the cross-section. But it is the traceless form of $T_{\alpha\beta}$ presented in (7.12) which has a non-renormalized two-point function. The reason is that the conformal $T_{\alpha\beta}$ is in the same supersymmetry multiplet as the supercurrents and the $SU(4)$ R-currents. $\mathcal{N} = 4$ super-Yang-Mills theory is finite to all orders and anomaly free in flat space. The Adler-Bardeen theorem guarantees that any anomalies of the $SU(4)$ R-currents can be computed exactly at one loop. Because $\partial_\alpha R^\alpha$ and $T^\alpha{}_\alpha$ are in the same supermultiplet (the so-called “multiplet of anomalies”), the one-loop result for the trace anomaly must also be exact.¹ Thus the trace anomaly in a curved background is the same as in the free theory:

$$\langle T^\mu{}_\mu \rangle = -\frac{1}{16\pi^2} \left(cF - \frac{2c}{3} \square R - bG \right) , \quad (7.14)$$

where $F = C_{\alpha\beta\gamma\delta}^2$ is the square of the Weyl tensor and $G = R_{\alpha\beta\gamma\delta}^2 - 4R_{\alpha\beta}^2 + R^2$ is the topological Euler density. Thus the second and third terms in (7.14) are total derivatives. The coefficients c and b are given by [28]

$$\begin{aligned} c &= \frac{12N_1 + 3N_{1/2} + N_0}{120} = \frac{N^2}{4} \\ b &= \frac{124N_1 + 11N_{1/2} + 2N_0}{720} = \frac{N^2}{4} . \end{aligned} \quad (7.15)$$

$N_1 = N^2$, $N_{1/2} = 4N^2$, and $N_0 = 6N^2$ are the numbers of spin-one, Majorana spin-half, and real spin-zero fields in the super-Yang-Mills theory. Note that spin-one, spin-half, and spin-zero particles make contributions to c in the ratio 2 : 2 : 1. The different spins contribute to the cross-section in precisely the same ratio, as was demonstrated in chapter 6.

¹We thank D. Anselmi for pointing out the relevance of the Adler-Bardeen theorem.

It remains to make the connection between $\langle T_\alpha^\alpha \rangle$ and $\langle T_{\alpha\beta}(x)T_{\gamma\delta}(0) \rangle$. Suppressing numerical factors and Lorentz structure, the OPE of $T_{\alpha\beta}$ with $T_{\gamma\delta}$ is [177]

$$T(x)T(0) = \frac{c}{x^8} + \dots + \frac{T(0)}{x^4} + \dots \quad (7.16)$$

We have omitted terms involving the Konishi current as well as terms less singular than $1/x^4$, and we have anticipated the conclusion that the coefficient on the Schwinger term is precisely the central charge c appearing in (7.14). A clean argument to this effect is presented in [178] and summarized briefly below. The Schwinger term is nothing but the two-point function: with all factors and indices written out explicitly² [179],

$$\langle T_{\alpha\beta}(x)T_{\gamma\delta}(0) \rangle = \frac{c}{48\pi^4} X_{\alpha\beta\gamma\delta} \left(\frac{1}{x^4} \right) \quad (7.17)$$

where

$$\begin{aligned} X_{\alpha\beta\gamma\delta} = & 2\Box^2 \eta_{\alpha\beta}\eta_{\gamma\delta} - 3\Box^2 (\eta_{\alpha\gamma}\eta_{\beta\delta} + \eta_{\alpha\delta}\eta_{\beta\gamma}) - 4\partial_\alpha\partial_\beta\partial_\gamma\partial_\delta \\ & - 2\Box (\partial_\alpha\partial_\beta\eta_{\gamma\delta} + \partial_\alpha\partial_\gamma\eta_{\beta\delta} + \partial_\alpha\partial_\delta\eta_{\beta\gamma} + \partial_\beta\partial_\gamma\eta_{\alpha\delta} + \partial_\beta\partial_\delta\eta_{\alpha\gamma} + \partial_\gamma\partial_\delta\eta_{\alpha\beta}) . \end{aligned} \quad (7.18)$$

The argument of [178] starts by obtaining an expression for the flat space three-point function $\langle T_\alpha^\alpha(x)T_{\beta\gamma}(y)T_{\rho\sigma}(z) \rangle$ by expanding (7.14) around flat space to second order. Using Ward identities for the conservation of $T_{\mu\nu}$, one can then derive a relation on two-point functions which can only be satisfied if the coefficients c in (7.17) and (7.14) are identical.

We are finally ready to compute the cross-section. Fourier transforming the two-point function (7.17), one obtains

$$\Pi_{\alpha\beta\gamma\delta}(p) = \int d^4x e^{ip \cdot x} \langle T_{\alpha\beta}(x)T_{\gamma\delta}(0) \rangle = \frac{c}{48\pi^4} \hat{X}_{\alpha\beta\gamma\delta} \int d^4x \frac{e^{ip \cdot x}}{x^4} \quad (7.19)$$

where $\hat{X}_{\alpha\beta\gamma\delta}$ is just the $X_{\alpha\beta\gamma\delta}$ of (7.18) with $\partial \rightarrow -ip$. The integral is evaluated formally as

$$\Pi(s) = \int d^4x \frac{e^{ip \cdot x}}{x^4} = \pi^2 \log(-s) + (\text{analytic in } s) \quad (7.20)$$

where, as before, $s = p^2$. One easily reads off the discontinuity across the positive real axis in the s -plane:

$$\text{Disc } \Pi(s) = \Pi(s + i\varepsilon) - \Pi(s - i\varepsilon) = -2\pi^3 i , \quad (7.21)$$

and so

$$\text{Disc } \Pi_{\alpha\beta\gamma\delta}(p) = -\frac{ic}{24\pi} \hat{X}_{\alpha\beta\gamma\delta} . \quad (7.22)$$

For the sake of definiteness, let us consider a graviton polarized in the x^1 - x^2 direction. $\hat{X}_{1212} = -3\omega^4$ for normal incidence, so

$$\sigma = \frac{2\kappa^2}{2i\omega} \text{Disc } \Pi_{1212}(p) \Big|_{\substack{p^0=\omega \\ \vec{p}=0}} = \frac{c}{8\pi} \kappa^2 \omega^3 , \quad (7.23)$$

in agreement with the classical result when $c = N^2/4$. The extra factor of $2\kappa^2$ in the second expression in (7.23) comes from the fact that $h_{\alpha\beta}$ as defined in the text following (7.11) is not a canonically normalized scalar field; instead, $h_{\alpha\beta}/\sqrt{2\kappa^2}$ is.

In summary, the non-renormalization argument is as follows: the graviton cross-section is read off at leading order in κ , but correct to all orders in g_{str} , from the two-point function of the stress-energy

²The field normalization conventions in [179] differ from those used here.

tensor. The two-point function is not renormalized beyond one loop because the Schwinger term in the OPE (7.16) is similarly non-renormalized. That in turn is due to the fact that the central charge appearing in the Schwinger term is precisely the coefficient of the Weyl tensor squared in the trace anomaly (7.14). The trace anomaly is not renormalized past one loop because T^α_α is related by supersymmetry to the divergence of the $SU(4)$ R -current, which is protected by the Adler-Bardeen theorem against anomalies beyond one loop. Another, more heuristic, reason why the central charge should not be renormalized is that there is a critical line extending from $g_{\text{str}}N = 0$ to $g_{\text{str}}N = \infty$. The central charge is expected to be constant along a critical line. In four dimensions, this expectation is supported by the work of [179].³ Therefore, c can be calculated in the $g_{\text{str}}N \rightarrow 0$ limit where it is given by one-loop diagrams.

We could adopt a different strategy and invert our arguments. Requiring that the world volume theory of N coincident three-branes agrees with semi-classical supergravity tells us that in the $g_{\text{str}}N \rightarrow \infty$ limit its central charge approaches $N^2/4$. In the next section we will similarly deduce the Schwinger terms in the two-point functions of the stress-energy tensor for other branes. Furthermore, we can study two-point functions of other operators by calculating the semi-classical absorption cross-section for particles that couple to them. For instance, the dilaton couples to $\text{tr } F^2$. The dilaton absorption cross-section was calculated in [9]. The semi-classical result implies that the Schwinger term here is again $\sim N^2$ in the $g_{\text{str}}N \rightarrow \infty$ limit. The comparisons with the gauge theory calculation in chapter 6 suggest that the one-loop result is again exact. It will be interesting to extract more results from these connections between gravity and gauge theory.

7.3 Extension to other branes

In this section we further explore the connection between the absorption cross-sections in semi-classical supergravity and central terms in two-point functions calculated in corresponding world volume theories. We proceed in analogy to the three-brane discussion presented in the previous section, and consider absorption of gravitons polarized along the branes. In the world volume theories such gravitons couple to components of stress-energy tensor, $T_{\alpha\beta}$. On the other hand, in supergravity such gravitons satisfy the minimally coupled scalar equation with respect to the coordinates transverse to the brane [10], as we have seen in chapter 6. This establishes a general connection between the absorption cross-section of a minimally coupled scalar and the central term in the algebra of stress-energy tensors. For N coincident three-branes the world volume theory is known, and we have shown that the conformal anomaly is in exact agreement with this principle. There are cases, however, where little is known about the world volume theory of multiple branes. In such cases we can use our method to find the Schwinger term without knowing any details of the world volume theory.

Consider, for instance, the 2-branes and the 5-branes of M-theory. While the world volume theory of multiple coincident branes is not known in detail, the extreme supergravity solutions are well-known. The absorption cross-sections for low-energy gravitons polarized along the brane were calculated in [9, 10, 160], with the results

$$\sigma_2 = \frac{1}{6\sqrt{2}\pi} \kappa_{11}^2 \omega^2 N^{3/2} , \quad \sigma_5 = \frac{1}{3 \cdot 2^6 \pi^2} N^3 \omega^5 . \quad (7.24)$$

For N coincident M2-branes we may deduce that the schematic structure of the stress-energy tensor OPE is

$$T(x)T(0) = \frac{c_2}{x^6} + \dots \quad (7.25)$$

where the central charge behaves as $c_2 \sim N^{3/2}$ in the large N limit. For N coincident M5-branes we

³A more definitive argument has been given in four dimensions for the constancy of flavor central charges along fixed lines [180].

instead have

$$T(x)T(0) = \frac{c_5}{x^{12}} + \dots \quad (7.26)$$

Now the central charge behaves as $c_5 \sim N^3$ in the large N limit. These results have an obvious connection with properties of the near-extremal entropy found in [46]. Indeed, the near-extremal entropy of a large number N of coincident M2-branes is formally reproduced by $\mathcal{O}(N^{3/2})$ massless free fields in 2+1 dimensions, while that of N coincident M5-branes is reproduced by $\mathcal{O}(N^3)$ massless free fields in 5+1 dimensions.

As a final example we consider the 5-branes of string theory. In [76] it was shown that their near-extremal entropy is reproduced by a novel kind of string theory, rather than by massless fields in 5+1 dimensions. For N coincident D5-branes the string tension turns out to be that of a D-string divided by N . This suggests that the degrees of freedom responsible for the near-extremal entropy are those of “fractionated” D-strings bound to the D5-branes. An S-dual of this picture suggests that the entropy of multiple NS-NS 5-branes comes from fractionated fundamental strings bound to them.⁴ We would like to learn more about these theories by probing them with longitudinally polarized gravitons incident transversely to the brane.

The extreme Einstein metric of both the NS-NS and the R-R 5-branes is

$$ds_E^2 = \left(1 + \frac{R^2}{r^2}\right)^{-1/4} (-dt^2 + dx_1^2 + \dots + dx_5^2) + \left(1 + \frac{R^2}{r^2}\right)^{3/4} (dr^2 + r^2 d\Omega_3^2) \quad (7.27)$$

The s -wave Laplace equation in the background of this metric is

$$\left[\rho^{-3} \frac{d}{d\rho} \rho^3 \frac{d}{d\rho} + 1 + \frac{(\omega R)^2}{\rho^2} \right] \phi(\rho) = 0 \quad (7.28)$$

where $\rho = \omega r$. Remarkably, this equation is exactly solvable in terms of Bessel functions. The two possible solutions are

$$\rho^{-1} J_{\pm \sqrt{1 - (\omega R)^2}}(\rho) \quad (7.29)$$

Clearly, there are two physically different regimes. For $\omega R > 1$ the label of the Bessel function is imaginary, and the requirement that the wave is incoming for $\rho \rightarrow 0$ selects the solution

$$\rho^{-1} J_{-i\sqrt{(\omega R)^2 - 1}}(\rho) \quad (7.30)$$

From the large ρ asymptotics we find that the absorption probability is

$$\mathcal{P} = 1 - e^{-2\pi\sqrt{(\omega R)^2 - 1}} \quad (7.31)$$

Hence, for $\omega R \geq 1$ the absorption cross-section is

$$\sigma = \frac{4\pi}{\omega^3} \left(1 - e^{-2\pi\sqrt{(\omega R)^2 - 1}} \right) \quad (7.32)$$

For $\omega R < 1$ the question of how to choose the solution is somewhat more subtle. It is clear that $\rho^{-1} J_{\sqrt{1 - (\omega R)^2}}(\rho)$ is better behaved near $\rho = 0$ than $\rho^{-1} J_{-\sqrt{1 - (\omega R)^2}}(\rho)$. If we approach the extreme 5-brane as a limit of a near-extreme 5-brane, we indeed find that $\rho^{-1} J_{\sqrt{1 - (\omega R)^2}}(\rho)$ is the solution that is selected. Since this solution is real, there is no absorption for $\omega R < 1$.⁵ This result agrees with the extremal limit of the 5-brane absorption cross-section calculated in [112].

⁴New insights into the world volume theory of NS-NS 5-branes were recently obtained in [181, 182, 183, 184].

⁵The fact that an extremal 5-brane in 10 dimensions does not absorb minimally coupled scalars below a certain threshold was noted in [170].

It is not hard to generalize our calculation to higher partial waves. For the ℓ -th partial wave we find that the absorption cross-section vanishes for $\omega R \leq \ell + 1$. Above this threshold it is given by

$$\sigma_\ell = \frac{4\pi(\ell+1)^2}{\omega^3} \left(1 - e^{-2\pi\sqrt{(\omega R)^2 - (\ell+1)^2}}\right). \quad (7.33)$$

From (7.32) we reach the surprising conclusion that

$$\langle T(\omega, \vec{0}) T(-\omega, \vec{0}) \rangle \quad (7.34)$$

vanishes identically for $\omega < 1/R$, which implies that gravity does not couple to the massless modes of the world volume theory! The threshold energy $1/R$ is precisely $1/\sqrt{\alpha'_{\text{eff}}}$, where $\alpha'_{\text{eff}} = 1/(2\pi T_{\text{eff}})$ and

$$T_{\text{eff}} = \frac{1}{2\pi R^2} \quad (7.35)$$

is the tension of the fractionated strings. The ordinary superstring has its first massive excited state at mass $m^2 = 2/\alpha'$. The threshold energy squared is half this value with α' replaced by α'_{eff} . If one imagines producing a single massive string at $\omega = 1/R$, then its mass is $m = 1/\sqrt{\alpha'_{\text{eff}}}$. Perhaps this is the first excited level of the non-critical string living on the 5-brane. Similarly, the higher partial wave thresholds might correspond to higher excited levels of mass $(\ell+1)/\sqrt{\alpha'_{\text{eff}}}$. If instead $\omega \geq 1/R$ corresponds to the pair production threshold of the first massive state of fractionated strings, then the mass would be $m = 1/(2\sqrt{\alpha'_{\text{eff}}})$. Neither picture yields any obvious explanation of the behavior

$$\langle T(\omega, \vec{0}) T(-\omega, \vec{0}) \rangle \sim [(\omega R)^2 - 1]^{1/2} \quad (7.36)$$

just above threshold. Pair production in a weakly interacting theory would predict a $7/2$ power in (7.36). It would be interesting to find an explanation of the observed square-root scaling in (7.36).

We believe that our discussion applies to a large number of coincident D5-branes, as well as to solitonic 5-branes of type IIA and IIB theories. This is because all these solutions have the same Einstein metric. Perhaps in the large N limit some properties of the world volume theories of these different branes become identical.⁶ For N coincident NS-NS 5-branes,

$$R^2 \sim N\alpha'. \quad (7.37)$$

Thus, the absorption cross-section (7.32) is formally independent of g_{str} . Therefore, our formula should be applicable in the $g_{\text{str}} \rightarrow 0$ limit proposed in [182]. There it was argued that in this limit the NS-NS 5-branes decouple from the bulk modes. We indeed find that incident gravitons are not absorbed for sufficiently low energies. However, above a critical energy of order $1/\sqrt{N\alpha'}$ the 5-branes do appear to couple to the bulk modes. As we have commented, this is probably related to the fact that the scale of the string theory living on the 5-brane is

$$\alpha'_{\text{eff}} = N\alpha', \quad (7.38)$$

i.e. the fundamental strings become fractionated [76].

In summary, we note that probing branes with low-energy particles incident from transverse directions is a useful tool for extracting correlation functions in their world volume theory. Here we have given an application of this technique to M2-branes and to 5-branes, but a more general investigation would be worthwhile.

⁶We thank J. Maldacena for suggesting this possibility to us.

Chapter 8

Green's functions from supergravity

8.1 Introduction

The nature of the relation between gauge fields and strings is an old, fascinating, and largely unanswered question. Its full answer is of great importance for theoretical physics. On one hand, it should provide us with a theory of quark confinement by explaining the dynamics of color-electric fluxes. On the other hand, it will perhaps uncover the true “gauge” degrees of freedom of the fundamental string theories, and therefore of gravity. There is still a long way to go before we can hope to give a satisfactory string theory account of the gauge theories observed in the real world. In this chapter, based on the paper [18], we attempt nevertheless to make some forward progress by considering the relation of maximally supersymmetric gauge theory to supergravity via D3-branes.

The Wilson loops of gauge theories satisfy loop equations which translate the Schwinger-Dyson equations into variational equations on the loop space [185, 186]. These equations should have a solution in the form of the sum over random surfaces bounded by the loop. These are the world surfaces of the color-electric fluxes. For the $SU(N)$ Yang-Mills theory they are expected to carry the 't Hooft factor [187], N^χ , where χ is the Euler character. Hence, in the large N limit where $g_{YM}^2 N$ is kept fixed only the simplest topologies are relevant.

Until recently, the action for the “confining string” had not been known. In [188] it was suggested that it must have a rather unusual structure. Let us describe it briefly. First of all, the world surface of the electric flux propagates in at least 5 dimensions. This is because the non-critical strings are described by the fields

$$X^\mu(\sigma) , \quad g_{ij}(\sigma) = e^{\varphi(\sigma)} \delta_{ij} , \quad (8.1)$$

where X^μ belong to 4-dimensional (Euclidean) space and $g_{ij}(\sigma)$ is the world sheet metric in the conformal gauge. The general form of the world sheet lagrangian compatible with the 4-dimensional symmetries is

$$\mathcal{L} = \frac{1}{2}(\partial_i \varphi)^2 + a^2(\varphi)(\partial_i X^\mu)^2 + \Phi(\varphi)^{(2)}R + \text{Ramond} - \text{Ramond backgrounds} , \quad (8.2)$$

where $^{(2)}R$ is the world sheet curvature, $\Phi(\varphi)$ is the dilaton [189], while the field

$$\Sigma(\varphi) = a^2(\varphi)$$

defines a variable string tension. In order to reproduce the zig-zag symmetry of the Wilson loop, the gauge fields must be located at a certain value $\varphi = \varphi_*$ such that $a(\varphi_*) = 0$. We will call this point “the horizon.”

The background fields $\Phi(\varphi)$, $a(\varphi)$ and others must be chosen to satisfy the conditions of conformal invariance on the world sheet [190]. After this is done, the relation between gauge fields and strings can be described as an isomorphism between the general Yang-Mills operators of the type

$$\int d^4x e^{ip \cdot x} \text{tr} (\nabla_{\alpha_1} \dots F_{\mu_1 \nu_1} \dots \nabla_{\alpha_n} \dots F_{\mu_n \nu_n}(x)) \quad (8.3)$$

and the algebra of vertex operators of string theory, which have the form

$$V^{\alpha_1 \dots \alpha_n}(p) = \int d^2\sigma \Psi_p^{i_1 \dots i_n j_1 \dots j_m}(\varphi(\sigma)) e^{ip \cdot X(\sigma)} \partial_{i_1} X^{\alpha_1} \dots \partial_{i_n} X^{\alpha_n} \partial_{j_1} \varphi \dots \partial_{j_m} \varphi, \quad (8.4)$$

where the wave functions $\Psi_p^{i_1 \dots i_n j_1 \dots j_m}(\varphi)$ are again determined by the conformal invariance on the world sheet. The isomorphism mentioned above implies the coincidence of the correlation functions of these two sets of vertex operators.

Another, seemingly unrelated, development is connected with the Dirichlet brane [4] description of black three-branes in [45, 9, 10, 39]. The essential observation is that, on the one hand, the black branes are solitons which curve space [54] and, on the other hand, the world volume of N parallel D-branes is described by supersymmetric $U(N)$ gauge theory with 16 supercharges [13]. A particularly interesting system is provided by the limit of a large number N of coincident D3-branes [45, 9, 10, 39], whose world volume is described by $\mathcal{N} = 4$ supersymmetric $U(N)$ gauge theory in $3 + 1$ dimensions. For large $g_{YM}^2 N$ the curvature of the classical geometry becomes small compared to the string scale [9], which allows for comparison of certain correlation functions between the supergravity and the gauge theory, with perfect agreement [9, 10, 39]. Corrections in powers of α' times the curvature on the string theory side correspond to corrections in powers of $(g_{YM}^2 N)^{-1/2}$ on the gauge theory side. The string loop corrections are suppressed by powers of $1/N^2$.

The vertex operators introduced in [9, 10, 39] describe the coupling of massless closed string fields to the world volume. For example, the vertex operator for the dilaton is

$$\int d^4x e^{ip \cdot x} \text{tr} F_{\mu\nu} F^{\mu\nu}(x) \quad (8.5)$$

while that for the graviton polarized along the three-branes is

$$\int d^4x e^{ip \cdot x} T^{\mu\nu}(x), \quad (8.6)$$

where $T^{\mu\nu}$ is the stress tensor. The low energy absorption cross-sections are related to the two-point functions of the vertex operators, and turn out to be in complete agreement with conformal invariance and supersymmetric non-renormalization theorems, as we saw in chapter 7. An earlier calculation [45] of the entropy as a function of temperature for N coincident D3-branes exhibits a dependence expected of a field theory with $O(N^2)$ massless fields. As we saw in chapter 2, the entropy turns out to be $3/4$ of the free field answer. This is not a discrepancy since the free field result is valid for small $g_{YM}^2 N$, while the result of [45] is applicable as $g_{YM}^2 N \rightarrow \infty$. We now regard this result as a non-trivial prediction of supergravity concerning the strong coupling behavior of $\mathcal{N} = 4$ supersymmetric gauge theory at large N and finite temperature.

The non-critical string and the D-brane approaches to $3+1$ dimensional gauge theory have been synthesized in [16] by rescaling the three-brane metric and taking the limit in which it has conformal symmetry, being the direct product $AdS_5 \times S^5$.¹ This is exactly the confining string ansatz [188] with

$$a(\varphi) = e^{\varphi/R}, \quad (8.7)$$

¹The papers that put an early emphasis on the anti-de Sitter nature of the near-horizon region of certain brane configurations, and its relation with string and M-theory, are [161, 142]. Other ideas on the relation between branes and AdS supergravity were recently pursued in [191, 192, 193].

corresponding to the case of constant negative curvature of order $1/R^2$. The horizon is located at $\varphi_* = -\infty$. The Liouville field is thus related to the radial coordinate of the space transverse to the three-brane. The extra S^5 part of the metric is associated with the 6 scalars and the $O(6)$ R-symmetry present in the $\mathcal{N} = 4$ supersymmetric gauge theory.

The purpose of this chapter is to make the next step and show how the excited states of the “confining string” are related to the anomalous dimensions of the SYM theory. Hopefully this analysis will help future explorations of asymptotically free gauge theories needed for quark confinement.

We will suggest a potentially very rich and detailed means of analyzing the throat-brane correspondence: we propose an identification of the generating function of the Green’s functions of the superconformal world-volume theory and the supergravity action in the near horizon background geometry. We will find it necessary to introduce a boundary of the AdS_5 space near the place where the throat turns into the asymptotically flat space. Thus, the anti-de Sitter coordinate φ is defined on a half-line $(-\infty, 0]$, similarly to the Liouville coordinate of the 2-dimensional string theory [194, 195]. The correlation functions are specified by the boundary terms in the action, again in analogy with the $c = 1$ case. One new prediction that we will be able to extract this way is for the anomalous dimensions of the gauge theory operators that correspond to massive string states. For a state at level n we find that, for large $g_{YM}^2 N$, the anomalous dimension grows as $2\sqrt{n}(2g_{YM}^2 N)^{1/4}$.

8.2 Green’s functions from the supergravity action

The geometry of a large number N of coincident D3-branes is

$$ds^2 = \left(1 + \frac{R^4}{r^4}\right)^{-1/2} (-dt^2 + d\vec{x}^2) + \left(1 + \frac{R^4}{r^4}\right)^{1/2} (dr^2 + r^2 d\Omega_5^2) . \quad (8.8)$$

The parameter R , where

$$R^4 = \frac{N}{2\pi^2 T_3} , \quad T_3 = \frac{\sqrt{\pi}}{\kappa} \quad (8.9)$$

is the only length scale for the geometry. T_3 is the tension of a single D3-brane, and κ is the ten-dimensional gravitational coupling. The near-horizon geometry of N D3-branes is $AdS_5 \times S^5$, as one can see most easily by defining the radial coordinate $z = R^2/r$. Then

$$ds^2 = \frac{R^2}{z^2} (-dt^2 + d\vec{x}^2 + dz^2) + R^2 d\Omega_5^2 . \quad (8.10)$$

The relation to the coordinate φ used in the previous section is

$$z = R e^{-\varphi/R} . \quad (8.11)$$

Note that the limit $z \rightarrow 0$ is far from the brane. Of course, for $z \lesssim R$ the AdS form (8.10) gets modified, and for $z \ll R$ one obtains flat ten-dimensional Minkowski space. We will continue to use the phrase “far from the brane” to emphasize that the all-important boundary conditions at small z are not imposed far down the throat, but rather at the border region where the throat merges into flat space. The geometry is geodesically complete and nonsingular, so there is no sense in which any one point on AdS_5 is the location of the brane.

The basic idea is to identify the generating functional of connected Green’s functions in the gauge theory with the minimum of the supergravity action, subject to some boundary conditions at $z = R$ and $z = \infty$:

$$W[g_{\mu\nu}(x^\lambda)] = K[g_{\mu\nu}(x^\lambda)] = S[g_{\mu\nu}(x^\lambda, z)] . \quad (8.12)$$

W generates the connected Green’s functions of the gauge theory; S is the supergravity action on the AdS space; while K is the minimum of S subject to the boundary conditions. We have kept only the

metric $g_{\mu\nu}(x^\lambda)$ of the world-volume as an explicit argument of W . The boundary conditions subject to which the supergravity action S is minimized are

$$ds^2 = \frac{R^2}{z^2} (g_{\mu\nu} dx^\mu dx^\nu + dz^2) + O(1) \quad \text{as } z \rightarrow R. \quad (8.13)$$

All fluctuations have to vanish as $z \rightarrow \infty$.

A few refinements of the identification (8.12) are worth commenting on. First, it is the generator of connected Green's functions which appears on the left hand side because the supergravity action on the right hand side is expected to follow the cluster decomposition principle. Second, because classical supergravity is reliable only for a large number N of coincident branes, (8.12) can only be expected to capture the leading large N behavior. Corrections in $1/N$ should be obtained as loop effects when one replaces the classical action S with an effective action Γ . This is sensible since the dimensionless expansion parameter $\kappa^2/R^8 \sim 1/N^2$. We also note that, since $(\alpha')^2/R^4 \sim (g_{YM}^2 N)^{-1}$, the string theoretic α' corrections to the supergravity action translate into gauge theory corrections proportional to inverse powers of $g_{YM}^2 N$. Finally, the fact that there is no covariant action for type IIB supergravity does not especially concern us: to obtain n -point Green's functions one is actually considering the n^{th} variation of the action, which for $n > 0$ can be regarded as the $(n-1)^{\text{th}}$ variation of the covariant equations of motion.

In section 8.2.2 we will compute a two-point function of massless vertex operators from (8.12), compare it with the absorption calculations in [9, 10, 39] and find exact agreement. However it is instructive first to examine boundary conditions.

8.2.1 Preliminary: symmetries and boundary conditions

As a preliminary it is useful to examine the appropriate boundary conditions and how they relate to the conformal symmetry. In this discussion we follow the work of Brown and Henneaux [196]. In the consideration of geometries which are asymptotically anti-de Sitter, one would like to have a realization of the conformal group on the asymptotic form of the metric. Restrictive or less restrictive boundary conditions at small z (far from the brane) corresponds, as Brown and Henneaux point out in the case of AdS_3 , to smaller or larger asymptotic symmetry groups. On an AdS_{d+1} space,

$$ds^2 = G_{mn} dx^m dx^n = \frac{R^2}{z^2} (-dt^2 + d\vec{x}^2 + dz^2) \quad (8.14)$$

where now $\vec{x}$ is $d-1$ dimensional, the boundary conditions which give the conformal group as the group of asymptotic symmetries are

$$\delta G_{\mu\nu} = O(1), \quad \delta G_{z\mu} = O(z), \quad \delta G_{zz} = O(1). \quad (8.15)$$

Our convention is to let indices m, n run from 0 to d (that is, over the full AdS_{d+1} space) while μ, ν run only from 0 to $d-1$ (i.e. excluding $z = x^d$). Diffeomorphisms which preserve (8.15) are specified by a vector ζ^m which for small z must have the form

$$\begin{aligned} \zeta^\mu &= \xi^\mu - \frac{z^2}{d} \eta^{\mu\nu} \partial_\nu (\xi^\kappa{}_{,\kappa}) + O(z^4) \\ \zeta^z &= \frac{z}{d} \xi^\kappa{}_{,\kappa} + O(z^3). \end{aligned} \quad (8.16)$$

Here ξ^μ is allowed to depend on t and $\vec{x}$ but not r . (8.16) specifies only the asymptotic form of ζ^m at large r , in terms of this new vector ξ^μ .

Now the condition that the variation

$$\delta G_{mn} = \mathcal{L}_\zeta G_{mn} = \zeta^k \partial_k G_{mn} + G_{kn} \partial_m \zeta^k + G_{mk} \partial_n \zeta^k \quad (8.17)$$

be of the allowed size specified in (8.15) is equivalent to

$$\xi_{\mu,\nu} + \xi_{\nu,\mu} = \frac{2}{d} \xi_{s,\kappa}^\kappa \quad (8.18)$$

where now we are lowering indices on ξ^μ with the flat space Minkowski metric $\eta_{\mu\nu}$. Since (8.18) is the conformal Killing equation in d dimensions ($d = 4$ for the three-brane), we see that we indeed recover precisely the conformal group from the set of permissible ξ^μ .

The spirit of [196] is to determine the central charge of an AdS_3 configuration by considering the commutator of deformations corresponding to Virasoro generators L_m and L_{-m} . This method is not applicable to higher dimensional cases because the conformal group becomes finite, and there is apparently no way to read off a Schwinger term from commutators of conformal transformations. Nevertheless, the notion of central charge can be given meaning in higher dimensional conformal field theories, either via a curved space conformal anomaly (also called the gravitational anomaly) or as the normalization of the two-point function of the stress energy tensor [178]. We shall see in section 8.2.2 that a calculation reminiscent of absorption probabilities allows us to read off the two-point function of stress-energy tensors in $\mathcal{N} = 4$ super-Yang-Mills, and with it the central charge.

8.2.2 The central charge from the stress tensor two-point function

As a preliminary, it is instructive to consider the case of a minimally coupled massless scalar propagating in the anti-de Sitter near-horizon geometry (one example of such a scalar is the dilaton ϕ [9]). As a further simplification we assume for now that ϕ is in the s -wave (that is, there is no variation over S^5). Then the action becomes

$$\begin{aligned} S &= \frac{1}{2\kappa^2} \int d^{10}x \sqrt{G} \left[\frac{1}{2} G^{MN} \partial_M \phi \partial_N \phi \right] \\ &= \frac{\pi^3 R^8}{4\kappa^2} \int d^4x \int_R^\infty \frac{dz}{z^3} \left[(\partial_z \phi)^2 + \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \right] . \end{aligned} \quad (8.19)$$

Note that in (8.19), as well as in all the following equations, we take κ to be the ten-dimensional gravitational constant. The equations of motion resulting from the variation of S are

$$\left[z^3 \partial_z \frac{1}{z^3} \partial_z + \eta^{\mu\nu} \partial_\mu \partial_\nu \right] \phi = 0 . \quad (8.20)$$

A complete set of normalizable solutions is

$$\phi_k(x^\ell) = \lambda_k e^{ik \cdot x} \tilde{\phi}_k(z) \quad \text{where} \quad \tilde{\phi}_k(z) = \frac{z^2 K_2(kz)}{R^2 K_2(kR)} , \quad (8.21)$$

$$k^2 = \vec{k}^2 - \omega^2 .$$

We have chosen the modified Bessel function $K_2(kz)$ rather than $I_2(kz)$ because the functions $K_\nu(kz)$ fall off exponentially for large z , while the functions $I_\nu(kz)$ grow exponentially. In other words, the requirement of regularity at the horizon (far down the throat) tells us which solution to keep. A connection of this choice with the absorption calculations of [9] is provided by the fact that, for time-like momenta, this is the incoming wave which corresponds to absorption from the small z region. λ_k is a coupling constant, and the normalization factor has been chosen so that $\tilde{\phi}_k = 1$ for $z = R$.

Let us consider a coupling

$$S_{\text{int}} = \int d^4x \phi(x^\lambda) \mathcal{O}(x^\lambda) \quad (8.22)$$

in the world-volume theory. If ϕ is the dilaton then according to [9] one would have $\mathcal{O} = \frac{1}{4} \text{tr } F^2$. Then the analogue of (8.12) is the claim that

$$W[\phi(x^\lambda)] = K[\phi(x^\lambda)] = S[\phi(x^\lambda, z)] \quad (8.23)$$

where $\phi(x^\lambda, z)$ is the unique solution of the equations of motion with $\phi(x^\lambda, z) \rightarrow \phi(x^\lambda)$ as $z \rightarrow R$. Note that the existence and uniqueness of ϕ are guaranteed because the equation of motion is just the Laplace equation on the curved space. (One could in fact compactify x^λ on very large T^4 and impose the boundary condition $\phi(x^\lambda, z) = \phi(x^\lambda)$ at $z = R$. Then the determination of $\phi(x^\lambda, z)$ is just the Dirichlet problem for the laplacian on a compact manifold with boundary).

Analogously to the work of [194] on the $c = 1$ matrix model, we can obtain the quadratic part of $K[\phi(x^\lambda)]$ as a pure boundary term through integration by parts,

$$\begin{aligned} K[\phi(x^\lambda)] &= \frac{\pi^3 R^8}{4\kappa^2} \int d^4x \int_R^\infty \frac{dz}{z^3} \left[-\phi \left(z^3 \partial_z \frac{1}{z^3} \partial_z + \eta^{\mu\nu} \partial_\mu \partial_\nu \right) \phi + z^3 \partial_z \left(\phi \frac{1}{z^3} \partial_z \phi \right) \right] \\ &= \frac{1}{2} \int d^4k d^4q \lambda_k \lambda_q (2\pi)^4 \delta^4(k+q) \frac{N^2}{16\pi^2} \int d^4x \mathcal{F} , \end{aligned} \quad (8.24)$$

where we have expanded

$$\phi(x^\lambda) = \int d^4k \lambda_k e^{ik \cdot x} .$$

The “flux factor” $\mathcal{F}$ (so named because of its resemblance to the particle number flux in a scattering calculation) is

$$\mathcal{F} = \left[\tilde{\phi}_k \frac{1}{z^3} \partial_z \tilde{\phi} \right]_R^\infty . \quad (8.25)$$

In (8.24) we have suppressed the boundary terms in the x^λ directions—again, one can consider these compactified on very large T^4 so that there is no boundary. We have also used (8.9) to simplify the prefactor. Finally, we have cut off the integral at R as a regulator of the small z divergence. This is in fact appropriate since the D3-brane geometry is anti-de Sitter only for $z \gg R$. Since there is exponential falloff in $\tilde{\phi}_k$ as $z \rightarrow \infty$, only the behavior at R matters.

To calculate the two-point function of $\mathcal{O}$ in the world-volume theory, we differentiate K twice with respect to the parameters λ_i :²

$$\begin{aligned} \langle \mathcal{O}(k) \mathcal{O}(q) \rangle &= \int d^4x d^4y e^{ik \cdot x + iq \cdot y} \langle \mathcal{O}(x) \mathcal{O}(y) \rangle \\ &= \frac{\partial^2 K}{\partial \lambda_k \partial \lambda_q} = (2\pi)^4 \delta^4(k+q) \frac{N^2}{16\pi^2} \mathcal{F} \\ &= -(2\pi)^4 \delta^4(k+q) \frac{N^2}{64\pi^2} k^4 \ln(k^2 R^2) + (\text{analytic in } k^2) \end{aligned} \quad (8.26)$$

where now the flux factor has been evaluated as

$$\mathcal{F} = \left[\tilde{\phi}_k \frac{1}{z^3} \partial_z \tilde{\phi}_k \right]_{z=R} = \left[\frac{1}{z^3} \partial_z \ln(\tilde{\phi}_k) \right]_{z=R} = -\frac{1}{4} k^4 \ln(k^2 R^2) + (\text{analytic in } k^2) . \quad (8.27)$$

Fourier transforming back to position space, we find

$$\langle \text{tr } F^2(x) \text{tr } F^2(y) \rangle \sim \frac{N^2}{|x-y|^8} . \quad (8.28)$$

²The appearance of the logarithm here is analogous to the logarithmic scaling violation in the $c = 1$ matrix model.

This is consistent with the free field result for small $g_{YM}^2 N$. Remarkably, supergravity tells us that this formula continues to hold as $g_{YM}^2 N \rightarrow \infty$.

Another interesting application of this analysis is to the two-point function of the stress tensor, which with the normalization conventions of chapter 7 is

$$\langle T_{\alpha\beta}(x) T_{\gamma\delta}(0) \rangle = \frac{c}{48\pi^4} X_{\alpha\beta\gamma\delta} \left(\frac{1}{x^4} \right) , \quad (8.29)$$

where the central charge $c = N^2/4$ and

$$\begin{aligned} X_{\alpha\beta\gamma\delta} = & 2\Box^2 \eta_{\alpha\beta} \eta_{\gamma\delta} - 3\Box^2 (\eta_{\alpha\gamma} \eta_{\beta\delta} + \eta_{\alpha\delta} \eta_{\beta\gamma}) - 4\partial_\alpha \partial_\beta \partial_\gamma \partial_\delta \\ & - 2\Box (\partial_\alpha \partial_\beta \eta_{\gamma\delta} + \partial_\alpha \partial_\gamma \eta_{\beta\delta} + \partial_\alpha \partial_\delta \eta_{\beta\gamma} + \partial_\beta \partial_\gamma \eta_{\alpha\delta} + \partial_\beta \partial_\delta \eta_{\alpha\gamma} + \partial_\gamma \partial_\delta \eta_{\alpha\beta}) . \end{aligned} \quad (8.30)$$

For metric perturbations $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ around flat space, the coupling of $h_{\mu\nu}$ at linear order is

$$S_{\text{int}} = \int d^4x \frac{1}{2} h^{\mu\nu} T_{\mu\nu} . \quad (8.31)$$

Furthermore, at quadratic order the supergravity action for a graviton polarized along the brane, $h_{xy}(k)$, is exactly the minimal scalar action, provided the momentum k is orthogonal to the xy plane. We can therefore carry over the result (8.26) to obtain

$$\langle T_{xy}(k) T_{xy}(q) \rangle = -(2\pi)^4 \delta^4(k+q) \frac{N^2}{64\pi^2} k^4 \ln(k^2 R^2) + (\text{analytic in } k^2) , \quad (8.32)$$

which upon Fourier transform can be compared with (8.29) to give $c = N^2/4$. In view of the conformal symmetry of both the supergravity and the gauge theory, the evaluation of this one component is a sufficient test.

The conspiracy of overall factors to give the correct normalization of (8.32) clearly has the same origin as the successful prediction of the minimal scalar s -wave absorption cross-section examined in [9] and chapters 6 and 7. The absorption cross-section is, up to a constant of proportionality, the imaginary part of (8.32). In [9] the absorption cross-section was calculated in supergravity using propagation of a scalar field in the entire three-brane metric, including the asymptotic region far from the brane. Here we have, in effect, replaced communication of the throat region with the asymptotic region by a boundary condition at one end of the throat. The physics of this is clear: signals coming from the asymptotic region excite the part of the throat near $z = R$. Propagation of these excitations into the throat can then be treated just in the anti-de Sitter approximation. Thus, to extract physics from anti-de Sitter space we must introduce a boundary at $z = R$ and take careful account of the boundary terms that contain the dynamical information.

It now seems clear how to proceed to three-point functions: on the supergravity side one must expand to third order in the perturbing fields, including in particular the three point vertices. At higher orders the calculation is still simple in concept (the classical action is minimized subject to boundary conditions), but the complications of the $\mathcal{N} = 8$ supergravity theory seem likely to make the computation of, for instance, the four-point function, rather tedious. We leave the details of such calculations for the future. It may be very useful to compute at least the three point functions in order to have a consistency check on the normalization of fields.

One extension of the present work is to consider what fields couple to the other operators in the $\mathcal{N} = 4$ supercurrent multiplet. The structure of the multiplet (which includes the supercurrents, the $SU(4)$ R -currents, and four spin 1/2 and one scalar field) suggests a coupling to the fields of gauged $\mathcal{N} = 4$ supergravity. The question then becomes how these fields are embedded in $\mathcal{N} = 8$ supergravity. We leave these technical issues for the future, but with the expectation that they are “bound to work” based on supersymmetry.

The main lesson we have extracted so far is that, for certain operators that couple to the massless string states, the anomalous dimensions vanish. We expect this to hold for all massless vertex operators. This may be the complete set of operators that are protected by supersymmetry. As we will see in the next section, other operators acquire anomalous dimensions that grow for large 't Hooft coupling.

8.3 Massive string states and anomalous dimensions

Before we proceed to the massive string states, a useful preliminary is to discuss the higher partial waves of a minimally coupled massless scalar. The action in five dimensions (with Lorentzian signature), the equations of motion, and the solutions are

$$S = \frac{\pi^3 R^8}{4\kappa^2} \int d^4x \int_R^\infty \frac{dz}{z^3} \left[(\partial_z \phi)^2 + (\partial_\mu \phi)^2 + \frac{\ell(\ell+4)}{z^2} \phi^2 \right] \quad (8.33)$$

$$\left[z^3 \partial_z \frac{1}{z^3} \partial_z + \eta^{\mu\nu} \partial_\mu \partial_\nu - \frac{\ell(\ell+4)}{z^2} \right] \phi = 0 \quad (8.34)$$

$$\phi_k(x^\ell) = e^{ik \cdot x} \tilde{\phi}_k(z) \quad \text{where} \quad \tilde{\phi}_k(z) = \frac{z^2 K_{\ell+2}(kz)}{R^2 K_{\ell+2}(kR)} . \quad (8.35)$$

We have chosen the normalization such that $\tilde{\phi}_k(z) = 1$ at $z = R$. The flux factor is evaluated by expanding

$$K_{\ell+2}(kz) = 2^{\ell+1} \Gamma(\ell+2) (kz)^{-(\ell+2)} \cdot \left(1 + \dots + \frac{(-1)^\ell}{2^{2\ell+3} (\ell+1)! (\ell+2)!} (kz)^{2(\ell+2)} \ln kz + \dots \right) , \quad (8.36)$$

where in parenthesis we exhibit the leading non-analytic term. We find

$$\mathcal{F} = \left[\frac{1}{z^3} \partial_z \ln(\tilde{\phi}_k) \right]_{z=R} = \frac{(-1)^\ell}{2^{2\ell+2} [(\ell+1)!]^2} k^{4+2\ell} R^{2\ell} \ln kR . \quad (8.37)$$

As before, we have neglected terms containing analytic powers of k and focused on the leading non-analytic term. This formula indicates that the operator that couples to ℓ -th partial wave has dimension $4 + \ell$. In [9] it was shown that such operators with the $SO(6)$ quantum numbers of the ℓ -th partial wave have the form

$$\int d^4x e^{ik \cdot x} \text{tr} \left[\left(X^{(i_1} \dots X^{i_\ell)} + \dots \right) F_{\mu\nu} F^{\mu\nu}(x) \right] , \quad (8.38)$$

where in parenthesis we have a traceless symmetric tensor of $SO(6)$. Thus, supergravity predicts that their non-perturbative dimensions equal their bare dimensions.

Now let us consider massive string states. Our goal is to use supergravity to calculate the anomalous dimensions of the gauge theory operators that couple to them. To simplify the discussion, let us focus on excited string states which are spacetime scalars of mass m . The propagation equation for such a field in the background of the three-brane geometry is

$$\left[\frac{d^2}{dr^2} + \frac{5}{r} \frac{d}{dr} - k^2 \left(1 + \frac{R^4}{r^4} \right) - m^2 \left(1 + \frac{R^4}{r^4} \right)^{1/2} \right] \tilde{\phi}_k = 0 . \quad (8.39)$$

For the state at excitation level n ,

$$m^2 = \frac{4n}{\alpha'} .$$

In the throat region, $z \gg R$, (8.39) simplifies to

$$\left[\frac{d^2}{dz^2} - \frac{3}{z} \frac{d}{dz} - k^2 - \frac{m^2 R^2}{z^2} \right] \tilde{\phi}_k = 0 . \quad (8.40)$$

Note that a massive particle with small energy $\omega \ll m$, which would be far off shell in the asymptotic region $z \ll R$, can nevertheless propagate in the throat region (i.e. it is described by an oscillatory wave function).

Equation (8.40) is identical to the equation encountered in the analysis of higher partial waves, except the effective angular momentum is not in general an integer: in the centrifugal barrier term $\ell(\ell+4)$ is replaced by $(mR)^2$. Analysis of the choice of wave function goes through as before, with $\ell+2$ replaced by ν , where

$$\nu = \sqrt{4 + (mR)^2} . \quad (8.41)$$

In other words, the wave function falling off exponentially for large z and normalized to 1 at $z = R$ is³

$$\tilde{\phi}_k(z) = \frac{z^2 K_\nu(kz)}{R^2 K_\nu(kR)} . \quad (8.42)$$

Now we recall that

$$K_\nu = \frac{\pi}{2 \sin(\pi\nu)} (I_{-\nu} - I_\nu) ,$$

$$I_\nu(z) = \left(\frac{z}{2}\right)^\nu \sum_{k=0}^{\infty} \frac{(z/2)^{2k}}{k! \Gamma(k + \nu + 1)} .$$

Thus,

$$K_\nu(kz) = 2^{\nu-1} \Gamma(\nu) (kz)^{-\nu} \left(1 + \dots - \left(\frac{zk}{2}\right)^{2\nu} \frac{\Gamma(1-\nu)}{\Gamma(1+\nu)} + \dots \right) , \quad (8.43)$$

where in parenthesis we have exhibited the leading non-analytic term. Calculating the flux factor as before, we find that the leading non-analytic term of the two-point function is

$$\langle \mathcal{O}(k) \mathcal{O}(q) \rangle = -(2\pi)^4 \delta^4(k+q) \frac{N^2}{8\pi^2} \frac{\Gamma(1-\nu)}{\Gamma(\nu)} \left(\frac{kR}{2}\right)^{2\nu} R^{-4} . \quad (8.44)$$

This implies that the dimension of the corresponding SYM operator is equal to $2 + \nu$.

Now, let us note that

$$R^4 = 2N g_{YM}^2 (\alpha')^2 ,$$

which implies

$$(mR)^2 = 4n g_{YM} \sqrt{2N} .$$

Using (8.41) we find that the spectrum of dimensions for operators that couple to massive string states is, for large $g_{YM} \sqrt{N}$,

$$h_n \approx 2 \left(n g_{YM} \sqrt{2N} \right)^{1/2} . \quad (8.45)$$

Equation (8.45) is a new non-trivial prediction of the string theoretic approach to large N gauge theory.⁴

We conclude that, for large 't Hooft coupling, the anomalous dimensions of the vertex operators corresponding to massive string states grow without bound. By contrast, the vertex operators that

³These solutions are reminiscent of the loop correlators calculated for $c \leq 1$ matrix models in [197].

⁴We do not expect this equation to be valid for arbitrarily large n , because application of linearized local effective actions to arbitrarily excited string states is questionable. However, we should be able to trust our approach for moderately excited states.

couple to the massless string states do not acquire any anomalous dimensions. This has been checked explicitly for gravitons, dilatons and RR scalars [9, 10, 39], and we believe this to be a general statement.⁵ Thus, there is an infinite class of operators that do not acquire anomalous dimensions (this is probably due to the fact that they are protected by SUSY). The rest of the operators are not protected and can receive arbitrarily large anomalous dimensions. Such large dimensions may be expected to “freeze out” the operators from the local operator algebra, just as the corresponding massive string states are frozen out of the supergravity description. While we cannot yet write down the explicit form of these operators in the gauge theory, it seems likely that they are conventional local operators, such as (8.3). Indeed, the coupling of a highly excited string state to the world volume may be guessed on physical grounds. Since a string in a D3-brane is a path of electric flux, it is natural to assume that a string state couples to a Wilson loop $\mathcal{O} = \exp\left(i \oint_{\gamma} A\right)$. Expansion of a small loop in powers of F yields the local polynomial operators.

There is one potential problem with our treatment of massive states. The minimal linear equation (8.39) where higher derivative terms are absent may be true only for a particular field definition (otherwise corrections in positive powers of $\alpha' \nabla^2$ will be present in the equation). Therefore, it is possible that there are energy dependent leg factors relating the operators $\mathcal{O}$ in (8.44) and the gauge theory operators of the form (8.3). We hope that these leg factors do not change our conclusion about the anomalous dimensions. However, to completely settle this issue we need to either find an exact sigma-model which incorporates all α' corrections or to calculate the three-point functions of massive vertex operators.

8.4 Conclusions

There are many unanswered questions that we have left for the future. So far, we have considered the limit of large 't Hooft coupling, since we used the one-loop sigma-model calculations for all operators involved. If this coupling is not large, then we have to treat the world sheet theory as an exact conformal field theory (we stress, once again, that the string loop corrections are $\sim 1/N^2$ and, therefore, vanish in the large N limit). This conformal field theory is a sigma-model on a hyperboloid. It is plausible that, in addition to the global $O(2,4)$ symmetry, this theory possesses the $O(2,4)$ Kac-Moody algebra. If this is the case, then the sigma model is tractable with standard methods of conformal field theory.

Throughout this work we detected many formal similarities of our approach with that used in $c \leq 1$ matrix models. These models may be viewed as early examples of gauge theory – non-critical string correspondence, with the large N matrix models playing the role of gauge theories. Clearly, a deeper understanding of the connection between the present work and the $c \leq 1$ matrix models is desirable.

⁵A more general set of such massless fields is contained in the supermultiplet of AdS gauge fields, whose reduction to the boundary was recently studied in [198].

Chapter 9

Concluding Remarks

Because this dissertation represents the author's contribution to a rapidly evolving field, it is perhaps more worthwhile to look ahead to where the field may be moving rather than to simply recapitulate the contents of chapters 2-8. There are certainly a number of avenues for further research, but perhaps the most attractive revolve around the idea of focusing on the near-horizon geometry of p -branes and the relation to gauge theory.

The near-horizon geometry of simple D-brane configurations is in general a homogeneous space: for example, D3-branes have as their near-horizon geometry $AdS_5 \times S^5 = SO(4, 2)/SO(4, 1) \times SO(6)/SO(5)$. Writing the anti-de Sitter factor as a quotient of groups makes the conformal symmetry manifest. There is certainly great interest in investigating further how the properties of conformal field theories can be reflected by supergravity on homogeneous spaces; but from a mainstream physics point of view the most significant extensions of the work in chapter 8 would involve non-conformal gauge theories, such as pure $\mathcal{N} = 2, 1$, or 0 gauge theory in four dimensions. These theories have non-vanishing beta-functions, and for $\mathcal{N} < 2$ there is confinement. Already in [199] a way of studying non-conformal theories and confinement “holographically” has been suggested, using brane configurations at finite temperature. A very clean way to proceed would be to perturb the D3-brane solution by what in the gauge theory are relevant operators (for instance, scalar or fermion masses). The resulting geometry should reflect softly broken $\mathcal{N} = 4$ gauge theory, which in general does confine in the infrared. To put it another way, $\mathcal{N} = 4$ would be the ultraviolet regulator of a confining theory. Several ideas for studying soft breakings in supergravity have been proposed, but the main difficulty lies in finding solutions of the full non-linear supergravity equations with less symmetry than the usual three-brane solution.

The greatest conceptual limitation of the supergravity–gauge theory correspondence is the requirement of large N and strong 't Hooft coupling $g_{YM}\sqrt{N}$. Let us review the reasons for these requirements. The formulas will be specific to the case of the three-brane, but similar observations apply to other brane configurations like the D1-D5 bound state. Supergravity has a scale (the Planck scale) because the gravitational coupling κ is dimensionful. The near-horizon geometry has length scale R and is an exact solution to the tree-level supergravity equations. Loop corrections in supergravity are suppressed only if κ/R^4 is small. But $\kappa/R^4 \sim 1/N$, so we see the origin of the large N requirement. Also, supergravity is the $\alpha' \rightarrow 0$ limit of string theory, and the α' corrections can be neglected only if $\alpha'/R^2 \sim 1/(g_{YM}\sqrt{N})$ is small—hence the requirement of large $g_{YM}\sqrt{N}$.

Gauge theory at large N is dominated by planar graphs, so one way to regard the relation

$$Z_{YM}[\phi_0] \sim \text{maximum}_{\phi=\phi_0 \text{ on } \partial AdS} e^{-S[\phi]} \quad \text{as } g_{YM}^2 N \rightarrow \infty \quad (9.1)$$

is to say that classical supergravity captures the large $g_{YM}\sqrt{N}$ asymptotics of planar graphs.

One would hope to improve (9.1) to an exact equality by replacing the right hand side with a full quantum theory of gravity: namely, type IIB string theory in the D3-brane background. As mentioned in chapter 1, there are formidable difficulties standing in the way of formulating string theory in a background involving Ramond-Ramond fields. Ramond-Ramond fields enter into the world-sheet description of strings as bilinears of spin fields with additional “superghost” dressings which maintain two-dimensional superconformal invariance. At the time of writing we are unaware of any proposals for formulating a ghost-free non-linear sigma-model action.

Notwithstanding the technical difficulties, we can ask what stringy corrections to (9.1) might be like. By including the full tower of massive excited string states, a string theory description would reintroduce the non-chiral gauge theory operators that were argued in chapter 8 to freeze out in the large coupling limit on account of large anomalous dimensions. The ability to make α'/R^2 arbitrary is precisely what we need in order to get down to weak coupling where perturbative gauge theory is reliable. Special quantities, like Schwinger terms in the OPE’s of conserved currents, can be expected to be protected against radiative corrections, as we saw in the case of the central charge in chapter 7. Tests of the gravity–gauge theory correspondence that rely in no way on symmetries, anomalies, or non-renormalizations are much harder to come by. This is perhaps typical of duality conjectures: expansion parameters get shuffled in such a way that what is simple on one side of the duality is very difficult to verify on the other.

One can nevertheless try to see what improvements to (9.1) might be the most accessible to calculation. One idea is to compute the leading $1/(g_{YM}\sqrt{N})$ correction to the entropy from the first α' correction to supergravity (which comes at order α'^3 for type IIB). Another possibility is to consider $1/N$ corrections via low order loop effects in quantized supergravity which are not troubled by divergences. For example, as explained in [19], the $U(1)$ part of the world-volume gauge theory is not visible in the AdS geometry, because the low-dimension conformal fields in the $U(1)$ sector would correspond to tachyons violating the Breitenlohner-Freedman bound. Consequently, supergravity in the throat region should reflect a central charge $c = (N^2 - 1)/4$ rather than the $c = N^2/4$ calculated in chapter 7. This slight difference may be one of the simplest $1/N^2$ effects. To see it in supergravity, one would want to evaluate the two-point function of the stress-tensor via graviton propagation in the AdS geometry, but including one-loop corrections to the propagator. The identification of parameters discussed in the previous paragraph guarantees that this will lead to some $1/N^2$ effect; whether it comes out with the correct coefficient is a check on whether (9.1) can be extended beyond the level of planar graphs.

Another direction, building upon the results in chapters 6 and 8, is to consider what effects in the world-volume theory become visible when one considers the full three-brane metric rather than just its near-horizon limit. Crudely speaking, the AdS space is “cut off” at a radius R by the asymptotically flat portion of the three-brane metric. Modes of the supergravity fields with $\omega R \ll 1$ do not see the effects of this cut-off because they approach a pure power law dependence at radii much larger than ωR^2 . Supergravity effects at these low energies seem to be all that is relevant to the renormalizable gauge theory on the world-volume. The full world-volume theory (presumably some non-abelian version of the DBI action) contains non-renormalizable terms which contribute to Green’s functions with inverse powers of R . By considering the double scaling limit (6.2) with finite ωR , one should be able to probe the complicated dynamics of the full DBI theory. The $\omega R \rightarrow 0$ limit is all that is explored by the analysis of the near-horizon geometry. In [200] the leading corrections in ωR to s -wave minimal scalar absorption were shown to correspond to the effects of breaking conformal invariance in the world-volume theory by retaining the leading non-renormalizable term in the DBI action. Further work in this direction has begun to suggest the appearance of anomalous dimensions at finite ωR for what in the conformal theory are protected operators whose dimensions are fixed by the superconformal algebra.

A better understanding of the double scaling limit may also be interesting from the point of view of black hole physics, for the following reason. In dynamical processes like absorption or emission, one

usually restricts to low energy. The physics picked out by this limit is the same as by the decoupling limit (where gravity is taken arbitrarily weak), the throat limit (where attention is focused on the near-horizon geometry), and the renormalizable gauge theory lagrangian on the world-volume. To put it another way, the “universal” low-energy behavior (1.3) of the cross sections has to do with a renormalizable theory on the brane decoupled from the bulk. If we instead we explore the double scaling limit with $(\omega R)^4 \sim N\kappa\omega^4$ held finite, then the cross-sections retain κ dependence. We may speculate that couplings between the brane and the bulk are being studied in a controlled way. The world-volume gauge theory is now non-renormalizable, and its time evolution cannot be regarded as strictly unitary because of the ultraviolet divergences. The energy scale of the non-renormalizable effects is $1/R$. Is it possible that this divergence-related non-unitarity is related to the apparent lack of unitarity in the evolution of black holes? If so, then string theory resolves information loss by providing a perfectly unitary description of the system at the (much higher) energy scale $1/\sqrt{\alpha'}$. This discussion has focused on the D3-brane, but in principle such considerations should extend to the effective string model. As mentioned before, the string theory description becomes less simple because of the complicated structure of the D1-D5 bound state. However, non-unitarity issues may be more straightforward to analyze in $1+1$ dimensions.

Bibliography

- [1] M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory*. Cambridge University Press, Cambridge, 1987.
- [2] M. Dine, “Supersymmetry phenomenology (with a broad brush),” [hep-ph/9612389](#).
- [3] J. H. Schwarz, “Lectures on superstring and M theory dualities: Given at ICTP Spring School and at TASI Summer School,” *Nucl. Phys. Proc. Suppl.* **55B** (1997) 1, [hep-th/9607201](#).
- [4] J. Polchinski, “Dirichlet Branes and Ramond-Ramond charges,” *Phys. Rev. Lett.* **75** (1995) 4724–4727, [hep-th/9510017](#).
- [5] A. Strominger and C. Vafa, “Microscopic origin of the Bekenstein-Hawking entropy,” *Phys. Lett.* **B379** (1996) 99–104, [hep-th/9601029](#).
- [6] C. G. Callan and J. M. Maldacena, “D-brane approach to black hole quantum mechanics,” *Nucl. Phys.* **B472** (1996) 591–610, [hep-th/9602043](#).
- [7] A. Dhar, G. Mandal, and S. R. Wadia, “Absorption vs. decay of black holes in string theory and T symmetry,” *Phys. Lett.* **B388** (1996) 51–59, [hep-th/9605234](#).
- [8] S. R. Das and S. D. Mathur, “Comparing decay rates for black holes and D-branes,” *Nucl. Phys.* **B478** (1996) 561–576, [hep-th/9606185](#).
- [9] I. R. Klebanov, “World volume approach to absorption by nondilatonic branes,” *Nucl. Phys.* **B496** (1997) 231, [hep-th/9702076](#).
- [10] S. S. Gubser, I. R. Klebanov, and A. A. Tseytlin, “String theory and classical absorption by three-branes,” *Nucl. Phys.* **B499** (1997) 217, [hep-th/9703040](#).
- [11] M. R. Douglas, D. Kabat, P. Pouliot, and S. H. Shenker, “D-branes and short distances in string theory,” *Nucl. Phys.* **B485** (1997) 85–127, [hep-th/9608024](#).
- [12] T. Banks, W. Fischler, S. H. Shenker, and L. Susskind, “M theory as a matrix model: A Conjecture,” *Phys. Rev.* **D55** (1997) 5112–5128, [hep-th/9610043](#).
- [13] E. Witten, “Bound states of strings and p-branes,” *Nucl. Phys.* **B460** (1996) 335–350, [hep-th/9510135](#).
- [14] E. Witten, “Branes and the dynamics of QCD,” *Nucl. Phys.* **B507** (1997) 658, [hep-th/9706109](#).
- [15] A. Giveon and D. Kutasov, “Brane dynamics and gauge theory,” [hep-th/9802067](#).
- [16] J. Maldacena, “The Large N limit of superconformal field theories and supergravity,” [hep-th/9711200](#).

- [17] A. Strominger, “Black hole entropy from near horizon microstates,” [hep-th/9712251](#).
- [18] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, “Gauge theory correlators from noncritical string theory,” [hep-th/9802109](#).
- [19] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” [hep-th/9803131](#).
- [20] J. Maldacena, “Wilson loops in large N field theories,” [hep-th/9803002](#).
- [21] C. G. Callan, S. S. Gubser, I. R. Klebanov, and A. A. Tseytlin, “Absorption of fixed scalars and the D-brane approach to black holes,” *Nucl. Phys.* **B489** (1997) 65–94, [hep-th/9610172](#).
- [22] S. S. Gubser, “Can the effective string see higher partial waves?,” *Phys. Rev.* **D56** (1997) 4984–4993, [hep-th/9704195](#).
- [23] S. S. Gubser, “Absorption of photons and fermions by black holes in four- dimensions,” *Phys. Rev.* **D56** (1997) 7854–7868, [hep-th/9706100](#).
- [24] J. M. Maldacena and A. Strominger, “Statistical entropy of four-dimensional extremal black holes,” *Phys. Rev. Lett.* **77** (1996) 428–429, [hep-th/9603060](#).
- [25] G. T. Horowitz, D. A. Lowe, and J. M. Maldacena, “Statistical entropy of nonextremal four-dimensional black holes and U duality,” *Phys. Rev. Lett.* **77** (1996) 430–433, [hep-th/9603195](#).
- [26] I. R. Klebanov and A. A. Tseytlin, “Intersecting M-branes as four-dimensional black holes,” *Nucl. Phys.* **B475** (1996) 179–192, [hep-th/9604166](#).
- [27] J. Maldacena, A. Strominger, and E. Witten, “Black hole entropy in M theory,” [hep-th/9711053](#).
- [28] P. C. W. Birrell and N. D. Davies, *Quantum Fields in Curved Space*. Cambridge University Press, Cambridge, UK, 1982.
- [29] E. Cremmer, “Supergravities in 5 dimensions,” in *Superspace and Supergravities* (S. W. Hawking and M. Rocek, eds.), p. 267. Cambridge University Press, 1981. Reprinted in *Supergravities in Diverse Dimensions*, (A. Salam and E. Sezgin, eds.), p. 422. World Scientific, 1989.
- [30] M. Cvetič, “Properties of black holes in toroidally compactified string theory,” *Nucl. Phys. Proc. Suppl.* **56B** (1997) 1, [hep-th/9701152](#).
- [31] A. Hashimoto, “Dynamics of Dirichlet-Neumann open strings on D-branes,” *Nucl. Phys.* **B496** (1997) 243, [hep-th/9608127](#).
- [32] R. G. Leigh, “Dirac-Born-Infeld action from Dirichlet sigma model,” *Mod. Phys. Lett.* **A4** (1989) 2767.
- [33] M. Cederwall, A. von Gussich, B. E. W. Nilsson, and A. Westerberg, “The Dirichlet super three brane in ten-dimensional type IIB supergravity,” *Nucl. Phys.* **B490** (1997) 163–178, [hep-th/9610148](#).
- [34] M. Cederwall, A. von Gussich, B. E. W. Nilsson, P. Sundell, and A. Westerberg, “The Dirichlet super p-branes in ten-dimensional type IIA and IIB supergravity,” *Nucl. Phys.* **B490** (1997) 179–201, [hep-th/9611159](#).

- [35] M. Aganagic, C. Popescu, and J. H. Schwarz, “Gauge invariant and gauge fixed D-brane actions,” *Nucl. Phys.* **B495** (1997) 99, [hep-th/9612080](#).
- [36] E. Bergshoeff and P. K. Townsend, “Super D-branes,” *Nucl. Phys.* **B490** (1997) 145–162, [hep-th/9611173](#).
- [37] W. G. Unruh, “Absorption cross section of small black holes,” *Phys. Rev.* **D14** (1976) 3251–3258.
- [38] J. C. Breckenridge, R. C. Myers, A. W. Peet, and C. Vafa, “D-branes and spinning black holes,” *Phys. Lett.* **B391** (1996) 93, [hep-th/9602065](#).
- [39] S. S. Gubser and I. R. Klebanov, “Absorption by branes and Schwinger terms in the world volume theory,” *Phys. Lett.* **B413** (1997) 41–48, [hep-th/9708005](#).
- [40] P. Ginsparg, “Applied conformal field theory.” Lectures given at Les Houches Summer School in Theoretical Physics, Les Houches, France, Jun 28 - Aug 5, 1988.
- [41] G. Moore and N. Seiberg, “Lectures on rcft.” Presented at Trieste Spring School 1989.
- [42] D. Friedan, Z. Qiu, and S. Shenker, “Conformal invariance, unitarity and two-dimensional critical exponents,” *Phys. Rev. Lett.* **52** (1984) 1575–1578.
- [43] C. F. E. Holzhey and F. Wilczek, “Black holes as elementary particles,” *Nucl. Phys.* **B380** (1992) 447–477, [hep-th/9202014](#).
- [44] A. A. Tseytlin, “On non-Abelian generalization of Born-Infeld action in string theory,” *Nucl. Phys.* **B501** (1997) 41, [hep-th/9701125](#).
- [45] S. S. Gubser, I. R. Klebanov, and A. W. Peet, “Entropy and temperature of black 3-branes,” *Phys. Rev.* **D54** (1996) 3915–3919, [hep-th/9602135](#).
- [46] I. R. Klebanov and A. A. Tseytlin, “Entropy of near extremal black p-branes,” *Nucl. Phys.* **B475** (1996) 164–178, [hep-th/9604089](#).
- [47] G. T. Horowitz and J. Polchinski, “A Correspondence principle for black holes and strings,” [hep-th/9612146](#).
- [48] I. R. Klebanov and L. Susskind, “Schwarzschild black holes in various dimensions from matrix theory,” *Phys. Lett.* **B416** (1998) 62, [hep-th/9709108](#).
- [49] N. Itzhaki, J. M. Maldacena, J. Sonnenschein, and S. Yankielowicz, “Supergravity and the large N limit of theories with sixteen supercharges,” [hep-th/9802042](#).
- [50] J. Polchinski, “TASI lectures on D-branes,” [hep-th/9611050](#).
- [51] G. T. Horowitz, J. M. Maldacena, and A. Strominger, “Nonextremal black hole microstates and U duality,” *Phys. Lett.* **B383** (1996) 151–159, [hep-th/9603109](#).
- [52] J. X. Lu, “ADM masses for black strings and p-branes,” *Phys. Lett.* **B313** (1993) 29–34, [hep-th/9304159](#).
- [53] A. Strominger, unpublished.
- [54] G. T. Horowitz and A. Strominger, “Counting states of near extremal black holes,” *Phys. Rev. Lett.* **77** (1996) 2368–2371, [hep-th/9602051](#).
- [55] P. S. Aspinwall, “K3 surfaces and string duality,” [hep-th/9611137](#).

- [56] T. Eguchi, P. B. Gilkey, and A. J. Hanson, “Gravitation, gauge theories, and differential geometry,” *Phys. Rept.* **66** (1980) 213.
- [57] F. Larsen and F. Wilczek, “Internal structure of black holes,” *Phys. Lett.* **B375** (1996) 37–42, [hep-th/9511064](#).
- [58] F. Larsen and F. Wilczek, “Classical hair in string theory. 2: Explicit calculations,” *Nucl. Phys.* **B488** (1997) 261–281, [hep-th/9609084](#).
- [59] R. Kallosh, A. Linde, T. Ortin, A. Peet, and A. V. Proeyen, “Supersymmetry as a cosmic censor,” *Phys. Rev.* **D46** (1992) 5278–5302, [hep-th/9205027](#).
- [60] M. Cvetič and D. Youm, “BPS saturated and nonextreme states in Abelian Kaluza-Klein theory and effective $N=4$ supersymmetric string vacua,” [hep-th/9508058](#).
- [61] M. Cvetič and D. Youm, “Dyonic BPS saturated black holes of heterotic string on a six torus,” *Phys. Rev.* **D53** (1996) 584–588, [hep-th/9507090](#).
- [62] M. Cvetič and A. A. Tseytlin, “Solitonic strings and BPS saturated dyonic black holes,” *Phys. Rev.* **D53** (1996) 5619–5633, [hep-th/9512031](#).
- [63] M. Cvetič and D. Youm, “General rotating five-dimensional black holes of toroidally compactified heterotic string,” *Nucl. Phys.* **B476** (1996) 118–132, [hep-th/9603100](#).
- [64] G. W. Gibbons, “Antigravitating black hole solitons with scalar hair in $N = 4$ supergravity,” *Nucl. Phys.* **B207** (1982) 337.
- [65] P. Breitenlohner, D. Maison, and G. W. Gibbons *Commun. Math. Phys.* **120** (1988) 295.
- [66] S. Ferrara and R. Kallosh, “Supersymmetry and attractors,” *Phys. Rev.* **D54** (1996) 1514–1524, [hep-th/9602136](#).
- [67] S. Ferrara and R. Kallosh, “Universality of supersymmetric attractors,” *Phys. Rev.* **D54** (1996) 1525–1534, [hep-th/9603090](#).
- [68] S. Ferrara, R. Kallosh, and A. Strominger, “ $N = 2$ extremal black holes,” *Phys. Rev.* **D52** (1995) 5412–5416, [hep-th/9508072](#).
- [69] G. Gibbons, R. Kallosh, and B. Kol, “Moduli, scalar charges, and the first law of black hole thermodynamics,” *Phys. Rev. Lett.* **77** (1996) 4992–4995, [hep-th/9607108](#).
- [70] B. Kol and A. Rajaraman, “Fixed scalars and suppression of Hawking evaporation,” *Phys. Rev.* **D56** (1997) 983–986, [hep-th/9608126](#).
- [71] S. R. Das, G. Gibbons, and S. D. Mathur, “Universality of low-energy absorption cross-sections for black holes,” *Phys. Rev. Lett.* **78** (1997) 417–419, [hep-th/9609052](#).
- [72] J. M. Maldacena and L. Susskind, “D-branes and fat black holes,” *Nucl. Phys.* **B475** (1996) 679–690, [hep-th/9604042](#).
- [73] S. R. Das and S. D. Mathur, “Interactions involving D-branes,” *Nucl. Phys.* **B482** (1996) 153–172, [hep-th/9607149](#).
- [74] I. R. Klebanov and L. Thorlacius, “The Size of p-branes,” *Phys. Lett.* **B371** (1996) 51–56, [hep-th/9510200](#).
- [75] J. Maldacena and A. Strominger, “Black hole grey body factors and D-brane spectroscopy,” *Phys. Rev.* **D55** (1997) 861–870, [hep-th/9609026](#).

- [76] J. M. Maldacena, “Statistical entropy of near extremal five-branes,” *Nucl. Phys.* **B477** (1996) 168–174, [hep-th/9605016](#).
- [77] A. A. Tseytlin, “Extremal black hole entropy from conformal string sigma model,” *Nucl. Phys.* **B477** (1996) 431–448, [hep-th/9605091](#).
- [78] E. Halyo, B. Kol, A. Rajaraman, and L. Susskind, “Counting Schwarzschild and charged black holes,” *Phys. Lett.* **B401** (1997) 15–20, [hep-th/9609075](#).
- [79] E. Halyo, “Reissner-Nordstrom black holes and strings with rescaled tension,” [hep-th/9610068](#).
- [80] J. Maharana and J. H. Schwarz, “Noncompact symmetries in string theory,” *Nucl. Phys.* **B390** (1993) 3–32, [hep-th/9207016](#).
- [81] E. Bergshoeff, C. Hull, and T. Ortin, “Duality in the type II superstring effective action,” *Nucl. Phys.* **B451** (1995) 547–578, [hep-th/9504081](#).
- [82] A. A. Tseytlin, “Extreme dyonic black holes in string theory,” *Mod. Phys. Lett.* **A11** (1996) 689–714, [hep-th/9601177](#).
- [83] M. Krasnitz and I. R. Klebanov, “Testing effective string models of black holes with fixed scalars,” *Phys. Rev.* **D56** (1997) 2173–2179, [hep-th/9703216](#).
- [84] J. M. Maldacena and A. Strominger, “Statistical entropy of four-dimensional extremal black holes,” *Phys. Rev. Lett.* **77** (1996) 428–429, [hep-th/9603060](#).
- [85] C. V. Johnson, R. R. Khuri, and R. C. Myers, “Entropy of 4-D extremal black holes,” *Phys. Lett.* **B378** (1996) 78–86, [hep-th/9603061](#).
- [86] I. R. Klebanov and A. A. Tseytlin, “Near extremal black hole entropy and fluctuating three-branes,” *Nucl. Phys.* **B479** (1996) 319–335, [hep-th/9607107](#).
- [87] S. S. Gubser and I. R. Klebanov, “Four-dimensional grey body factors and the effective string,” *Phys. Rev. Lett.* **77** (1996) 4491–4494, [hep-th/9609076](#).
- [88] M. Cvetič and A. A. Tseytlin, “Nonextreme black holes from nonextreme intersecting M-branes,” *Nucl. Phys.* **B478** (1996) 181–198, [hep-th/9606033](#).
- [89] G. Gilbert, “On the perturbations of string theoretic black holes,” [hep-th/9108012](#).
- [90] R. Gregory and R. Laflamme, “Evidence for stability of extremal black p-branes,” *Phys. Rev.* **D51** (1995) 305–309, [hep-th/9410050](#).
- [91] R. Gregory and R. Laflamme, “The Instability of charged black strings and p-branes,” *Nucl. Phys.* **B428** (1994) 399–434, [hep-th/9404071](#).
- [92] R. R. Khuri, “Manton scattering of string solitons,” *Nucl. Phys.* **B376** (1992) 350–364.
- [93] A. A. Tseytlin, “‘No force’ condition and BPS combinations of p-branes in eleven-dimensions and ten-dimensions,” *Nucl. Phys.* **B487** (1997) 141, [hep-th/9609212](#).
- [94] S. R. Das and S. D. Mathur, “Excitations of D strings, entropy and duality,” *Phys. Lett.* **B375** (1996) 103–110, [hep-th/9601152](#).
- [95] J. C. Breckenridge *et. al.*, “Macroscopic and microscopic entropy of near extremal spinning black holes,” *Phys. Lett.* **B381** (1996) 423–426, [hep-th/9603078](#).

- [96] I. R. Klebanov and M. Krasnitz, “Fixed scalar gray body factors in five-dimensions and four-dimensions,” *Phys. Rev.* **D55** (1997) 3250–3254, [hep-th/9612051](#).
- [97] M. Abramowitz and I. A. Stegun, eds., *Handbook of Mathematical Functions*. US Government Printing Office, Washington, DC, 1964.
- [98] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*. Academic Press, San Diego, fifth edition ed., 1994.
- [99] S. S. Gubser and I. R. Klebanov, “Emission of charged particles from four-dimensional and five-dimensional black holes,” *Nucl. Phys.* **B482** (1996) 173–186, [hep-th/9608108](#).
- [100] J. Maldacena and A. Strominger, “Universal low-energy dynamics for rotating black holes,” *Phys. Rev.* **D56** (1997) 4975–4983, [hep-th/9702015](#).
- [101] I. R. Klebanov, A. Rajaraman, and A. A. Tseytlin, “Intermediate scalars and the effective string model of black holes,” *Nucl. Phys.* **B503** (1997) 157, [hep-th/9704112](#).
- [102] S. W. Hawking and M. M. Taylor-Robinson, “Evolution of near extremal black holes,” *Phys. Rev.* **D55** (1997) 7680–7692, [hep-th/9702045](#).
- [103] F. Dowker, D. Kastor, and J. Traschen, “U duality, D-branes and black hole emission rates: Agreements and disagreements,” [hep-th/9702109](#).
- [104] S. D. Mathur, “Absorption of angular momentum by black holes and D- branes,” [hep-th/9704156](#).
- [105] G. N. Watson, *Theory of Bessel Functions*. Cambridge University Press, Cambridge, UK, second edition ed., 1962.
- [106] F. Larsen, “A String model of black hole microstates,” *Phys. Rev.* **D56** (1997) 1005–1008, [hep-th/9702153](#).
- [107] M. Aganagic, C. Popescu, and J. H. Schwarz, “D-brane actions with local kappa symmetry,” *Phys. Lett.* **B393** (1997) 311–315, [hep-th/9610249](#).
- [108] A. A. Tseytlin, “On non-abelian generalization of Born-Infeld action in string theory,” *Nucl. Phys.* **B501** (1997) 41, [hep-th/9701125](#).
- [109] S. S. Gubser, unpublished, April, 1997.
- [110] E. Keski-Vakkuri and P. Kraus, “Microcanonical D-branes and back reaction,” *Nucl. Phys.* **B491** (1997) 249–262, [hep-th/9610045](#).
- [111] J. Maldacena, “D-branes and near extremal black holes at low-energies,” *Phys. Rev.* **D55** (1997) 7645–7650, [hep-th/9611125](#).
- [112] I. R. Klebanov and S. D. Mathur, “Black hole grey body factors and absorption of scalars by effective strings,” *Nucl. Phys.* **B500** (1997) 115, [hep-th/9701187](#).
- [113] M. Cvetič and F. Larsen, “General rotating black holes in string theory: Grey body factors and event horizons,” *Phys. Rev.* **D56** (1997) 4994–5007, [hep-th/9705192](#).
- [114] M. Cvetič and F. Larsen, “Grey body factors for rotating black holes in four- dimensions,” *Nucl. Phys.* **B506** (1997) 107, [hep-th/9706071](#).
- [115] A. Das, “ $SO(4)$ invariant extended supergravity,” *Phys. Rev.* **D15** (1977) 2805.

- [116] E. Cremmer, J. Scherk, and S. Ferrara, “ $U(N)$ invariance in extended supergravity,” *Phys. Lett.* **68B** (1977) 234.
- [117] E. Cremmer and J. Scherk, “Algebraic simplifications in supergravity theories,” *Nucl. Phys.* **B127** (1977) 259.
- [118] E. Cremmer, J. Scherk, and S. Ferrara, “ $SU(4)$ invariant supergravity theory,” *Phys. Lett.* **74B** (1978) 61.
- [119] E. Witten, “Dyons of charge $e\theta/2\pi$,” *Phys. Lett.* **86B** (1979) 283–287.
- [120] S. A. Teukolsky, “Rotating black holes: separable wave equations for gravitational and electromagnetic perturbations,” *Phys. Rev. Lett.* **29** (1972) 1114–1118.
- [121] S. A. Teukolsky, “Perturbations of a rotating black hole I,” *Astrophys. J.* **185** (1973) 635–647.
- [122] A. A. Starobinskiĭ and S. M. Churilov, “Amplification of electromagnetic and gravitational waves scattered by a rotating “black hole”,” *Sov. Phys.-JETP* **38** (1974) 1–5.
- [123] E. T. Newman and R. Penrose, “An approach to gravitational radiation by a method of spin coefficients,” *J. Math. Phys.* **3** (1962) 566–578. erratum **4** (1963), 998.
- [124] R. Penrose and W. Rindler, *Spacetime and Spinors*, vol. 1. Cambridge University Press, Cambridge, UK, 1984.
- [125] W. Rarita and J. Schwinger, “On a theory of particles with half-integral spin,” *Phys. Rev.* **60** (1941) 61.
- [126] S. Weinberg, *The Quantum Theory of Fields*, vol. 1. Cambridge University Press, Cambridge, UK, 1995.
- [127] S. Chandrasekhar, “On the equations governing the perturbations of the Reissner-Nordstrom black hole,” *Proc. R. Soc. Lond.* **A365** (1979) 453–465.
- [128] J. N. Goldberg, A. J. Macfarlane, E. T. Newman, F. Rohrlich, and E. C. G. Sudarshan, “Spin- s Spherical Harmonics and $\not{\partial}$,” *J. Math. Phys.* **8** (1967) 2155–2161.
- [129] H. Onozawa, T. Okamura, T. Mishima, and H. Ishihara, “Perturbing supersymmetric black hole,” *Phys. Rev.* **D55** (1997) 4529–4531, [gr-qc/9606086](#).
- [130] T. Okamura, “On scattering off the extreme Reissner-Nordstrom black hole in $N = 2$ supergravity,” [gr-qc/9703050](#).
- [131] R. Kallosh, J. Rahmfeld, and W. K. Wong, “One loop supergravity corrections to the black hole entropy and residual supersymmetry,” *Phys. Rev.* **D57** (1998) 1063–1067, [hep-th/9706048](#).
- [132] F. Gliozzi, J. Scherk, and D. Olive, “Supersymmetry, supergravity theories and the dual spinor model,” *Nucl. Phys.* **B122** (1976) 253.
- [133] A. L. Fetter, *Quantum Theory of Many-particle Systems*. McGraw-Hill, New York, 1971.
- [134] A. Schwimmer and N. Seiberg, “Comments on the $N = 2$, $N = 3$, $N = 4$ superconformal algebras in two-dimensions,” *Phys. Lett.* **184B** (1987) 191.
- [135] R. M. Wald, *General Relativity*. Chicago University Press, Chicago, 1984.
- [136] S. S. Gubser, A. Hashimoto, I. R. Klebanov, and J. M. Maldacena, “Gravitational lensing by p-branes,” *Nucl. Phys.* **B472** (1996) 231–248, [hep-th/9601057](#).

- [137] M. R. Garousi and R. C. Myers, “Superstring scattering from D-branes,” *Nucl. Phys.* **B475** (1996) 193–224, [hep-th/9603194](#).
- [138] A. Hashimoto and I. R. Klebanov, “Decay of excited D-branes,” *Phys. Lett.* **B381** (1996) 437–445, [hep-th/9604065](#).
- [139] M. J. Duff, H. Lu, and C. N. Pope, “The Black branes of M theory,” *Phys. Lett.* **B382** (1996) 73–80, [hep-th/9604052](#).
- [140] H. Lu, S. Mukherji, C. N. Pope, and J. Rahmfeld, “Loop corrected entropy of near extremal dilatonic p- branes,” *Phys. Lett.* **B389** (1996) 248–254, [hep-th/9604127](#).
- [141] M. J. Duff and J. X. Lu, “The Selfdual type IIB superthreebrane,” *Phys. Lett.* **B273** (1991) 409–414.
- [142] G. W. Gibbons, G. T. Horowitz, and P. K. Townsend, “Higher dimensional resolution of dilatonic black hole singularities,” *Class. Quant. Grav.* **12** (1995) 297–318, [hep-th/9410073](#).
- [143] M. J. Duff, G. W. Gibbons, and P. K. Townsend, “Macroscopic superstrings as interpolating solitons,” *Phys. Lett.* **B332** (1994) 321–328, [hep-th/9405124](#).
- [144] J. H. Schwarz, “Covariant field equations of chiral $N = 2$ $D = 10$ supergravity,” *Nucl. Phys.* **B226** (1983) 269.
- [145] E. Bergshoeff, H. J. Boonstra, and T. Ortin, “S duality and dyonic p-brane solutions in type II string theory,” *Phys. Rev.* **D53** (1996) 7206–7212, [hep-th/9508091](#).
- [146] G. T. Horowitz and A. Strominger, “Black strings and P-branes,” *Nucl. Phys.* **B360** (1991) 197–209.
- [147] M. J. Duff and K. S. Stelle *Phys. Lett.* **B253** (1991) 113.
- [148] R. Guven *Phys. Lett.* **B276** (1992) 49.
- [149] E. Cremmer, B. Julia, and J. Scherk, “Supergravity theory in eleven dimensions,” *Phys. Lett.* **76B** (1978) 409.
- [150] D. Page *Phys. Rev.* **D15** (1976) 3260.
- [151] M. Li, “Boundary states of D-branes and Dy strings,” *Nucl. Phys.* **B460** (1996) 351–361, [hep-th/9510161](#).
- [152] M. R. Douglas, “Branes within branes,” [hep-th/9512077](#).
- [153] A. A. Tseytlin, “Selfduality of Born-Infeld action and Dirichlet 3-brane of type IIB superstring theory,” *Nucl. Phys.* **B469** (1996) 51–67, [hep-th/9602064](#).
- [154] A. Wasserman, Michaelmas, 1995. Part III lectures given at Cambridge University.
- [155] S. S. Gubser, unpublished, November, 1996.
- [156] E. Bergshoeff, E. Sezgin, and P. K. Townsend, “Supermembranes and eleven-dimensional supergravity,” *Phys. Lett.* **B189** (1987) 75–78.
- [157] E. Bergshoeff, M. de Roo, and T. Ortin, “The Eleven-dimensional five-brane,” *Phys. Lett.* **B386** (1996) 85–90, [hep-th/9606118](#).
- [158] M. Aganagic, J. Park, C. Popescu, and J. H. Schwarz, “World volume action of the M theory five-brane,” *Nucl. Phys.* **B496** (1997) 191, [hep-th/9701166](#).

- [159] I. Bandos *et. al.*, “Covariant action for the superfive-brane of M theory,” *Phys. Rev. Lett.* **78** (1997) 4332–4334, [hep-th/9701149](#).
- [160] R. Emparan, “Absorption of scalars by extended objects,” [hep-th/9706204](#).
- [161] G. W. Gibbons and P. K. Townsend, “Vacuum interpolation in supergravity via super p-branes,” *Phys. Rev. Lett.* **71** (1993) 3754–3757, [hep-th/9307049](#).
- [162] D. M. Kaplan and J. Michelson, “Zero modes for the D = 11 membrane and five-brane,” *Phys. Rev.* **D53** (1996) 3474–3476, [hep-th/9510053](#).
- [163] L. Alvarez-Gaume and E. Witten, “Gravitational anomalies,” *Nucl. Phys.* **B234** (1984) 269.
- [164] M. R. Douglas, “Gauge fields and D-branes,” [hep-th/9604198](#).
- [165] V. Balasubramanian and F. Larsen, “On D-branes and black holes in four-dimensions,” *Nucl. Phys.* **B478** (1996) 199–208, [hep-th/9604189](#).
- [166] A. A. Tseytlin, “Composite BPS configurations of p-branes in ten-dimensions and eleven-dimensions,” *Class. Quant. Grav.* **14** (1997) 2085, [hep-th/9702163](#).
- [167] M. Douglas, J. Polchinski, and A. Strominger, “Probing five-dimensional black holes with D-branes,” [hep-th/9703031](#).
- [168] M. Cvetič and A. Tseytlin, “General class of BPS saturated dyonic black holes as exact superstring solutions,” *Phys. Lett.* **B366** (1996) 95–103, [hep-th/9510097](#).
- [169] A. A. Tseytlin, “Harmonic superpositions of M-branes,” *Nucl. Phys.* **B475** (1996) 149–163, [hep-th/9604035](#).
- [170] V. Balasubramanian and F. Larsen, “Extremal branes as elementary particles,” *Nucl. Phys.* **B495** (1997) 206, [hep-th/9610077](#).
- [171] M. Cvetič and D. Youm, “Entropy of nonextreme charged rotating black holes in string theory,” *Phys. Rev.* **D54** (1996) 2612–2620, [hep-th/9603147](#).
- [172] J. Dai, R. G. Leigh, and J. Polchinski, “New connections between string theories,” *Mod. Phys. Lett.* **A4** (1989) 2073–2083.
- [173] S. R. Das, “The Effectiveness of D-branes in the description of near extremal black holes,” *Phys. Rev.* **D56** (1997) 3582–3590, [hep-th/9703146](#).
- [174] C. G. Callan, private communication, April 1997.
- [175] M. F. Sohnius, “Introducing supersymmetry,” *Phys. Rept.* **128** (1985) 39.
- [176] C. G. Callan, S. Coleman, and R. Jackiw, “A New improved energy - momentum tensor,” *Ann. Phys.* **59** (1970) 42–73.
- [177] D. Anselmi, M. Grisaru, and A. Johansen, “A Critical behavior of anomalous currents, electric - magnetic universality and CFT in four-dimensions,” *Nucl. Phys.* **B491** (1997) 221–248, [hep-th/9601023](#).
- [178] J. Erdmenger and H. Osborn, “Conserved currents and the energy momentum tensor in conformally invariant theories for general dimensions,” *Nucl. Phys.* **B483** (1997) 431–474, [hep-th/9605009](#).

- [179] D. Anselmi, D. Z. Freedman, M. T. Grisaru, and A. A. Johansen, “Universality of the operator product expansions of SCFT in four-dimensions,” *Phys. Lett.* **B394** (1997) 329–336, [hep-th/9608125](#).
- [180] D. Anselmi, D. Z. Freedman, M. T. Grisaru, and A. A. Johansen, “Nonperturbative formulas for central functions of supersymmetric gauge theories,” [hep-th/9708042](#).
- [181] R. Dijkgraaf, E. Verlinde, and H. Verlinde, “5-D black holes and matrix strings,” *Nucl. Phys.* **B506** (1997) 121, [hep-th/9704018](#).
- [182] N. Seiberg, “New theories in six-dimensions and matrix description of M theory on T^{*5} and $T^{*5}/Z(2)$,” *Phys. Lett.* **B408** (1997) 98–104, [hep-th/9705221](#).
- [183] O. Aharony, M. Berkooz, S. Kachru, N. Seiberg, and E. Silverstein, “Matrix description of interacting theories in six dimensions,” [hep-th/9707079](#).
- [184] E. Witten, “On the conformal field theory of the Higgs branch,” [hep-th/9707093](#).
- [185] A. M. Polyakov, “Gauge Fields as Rings of Glue,” *Nucl. Phys.* **B164** (1980) 171.
- [186] Y. M. Makeenko and A. A. Migdal, “Exact equation for the loop average in multicolor QCD,” *Phys. Lett.* **88B** (1979) 135.
- [187] G. ’t Hooft, “A planar diagram theory for strong interactions,” *Nucl. Phys.* **B72** (1974) 461.
- [188] A. M. Polyakov, “String theory and quark confinement,” [hep-th/9711002](#).
- [189] E. S. Fradkin and A. A. Tseytlin, “Effective field theory from quantized strings,” *Phys. Lett.* **158B** (1985) 316.
- [190] C. G. Callan, E. J. Martinec, M. J. Perry, and D. Friedan, “Strings in background fields,” *Nucl. Phys.* **B262** (1985) 593.
- [191] S. Hyun, “The Background geometry of DLCQ supergravity,” [hep-th/9802026](#).
- [192] H. J. Boonstra, B. Peeters, and K. Skenderis, “Duality and asymptotic geometries,” *Phys. Lett.* **B411** (1997) 59, [hep-th/9706192](#).
- [193] K. Sfetsos and K. Skenderis, “Microscopic derivation of the Bekenstein-Hawking entropy formula for nonextremal black holes,” [hep-th/9711138](#).
- [194] J. Polchinski, “Critical behavior of random surfaces in one dimension,” *Nucl. Phys.* **B346** (1990) 253–263.
- [195] S. R. Das and A. Jevicki, “String Field Theory and Physical Interpretation of $D = 1$ Strings,” *Mod. Phys. Lett.* **A5** (1990) 1639–1650.
- [196] J. D. Brown and M. Henneaux, “Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity,” *Commun. Math. Phys.* **104** (1986) 207.
- [197] G. Moore, N. Seiberg, and M. Staudacher, “From loops to states in 2-D quantum gravity,” *Nucl. Phys.* **B362** (1991) 665–709.
- [198] S. Ferrara and C. Fronsdal, “Conformal Maxwell theory as a singleton field theory on $adS(5)$, IIB three-branes and duality,” [hep-th/9712239](#).
- [199] E. Witten, “Anti-de Sitter space and holography,” [hep-th/9802150](#).
- [200] S. S. Gubser, A. Hashimoto, I. R. Klebanov, and M. Krasnitz, “Scalar absorption and the breaking of the world volume conformal invariance,” [hep-th/9803023](#).